



NAPPO

North American Plant Protection Organization
Organización Norteamericana de Protección a las Plantas

NAPPO Regional Standards for Phytosanitary Measures (RSPM)

RSPM 35

Guidelines for the Movement of Propagative Plant Material of Stone Fruit, Pome Fruit, and Grapevine into a NAPPO Member Country

The Secretariat of the North American Plant Protection Organization
1730 Varsity Drive, Suite 145
Raleigh, NC 27606-5202
United States of America
March 9, 2022

Arthropod pest tables were added as ink amendments to this standard on May 3, 2023

CONTENTS

Review	4
Approval	4
Virtual approval of NAPPO Products	4
Implementation	5
Amendment Record.....	5
Distribution	5
INTRODUCTION	6
Scope	6
References	6
Definitions.....	7
Outline of Requirements	7
Background	7
1. GENERAL REQUIREMENTS.....	8
1.1 Stone and Pome Fruit and Grapevine Pests.....	8
1.2 Pest Risk Analysis.....	8
1.3 Phytosanitary Measures for Pest Risk Management.....	8
1.3.1 Research Purposes and Subsequent Destruction.....	9
1.3.2 Quarantine, Testing and Treatment at NPPO-authorized Facilities and Subsequent Distribution.....	9
1.3.3 Quarantine at Importers' Premises and Subsequent Distribution	9
1.3.4 Plants originating from an Official Certification Program	9
1.3.5 Plants originating from a Pest-free Area, Pest Free Place of Production, or Pest Free Production Site.....	10
1.3.6 Prohibition	10
1.4 Documentation Requirements	10
2. SPECIFIC REQUIREMENTS	10
2.1 Stone and Pome Fruit and Grapevine Certification Programs.....	10
2.1.1 Program Administration	10
2.1.2 Terminology.....	11
2.1.3 Diagnostics.....	11
2.1.4 Eligibility	12
2.1.5 Generation Levels	12
2.1.6 Agricultural Management.....	12
2.1.7 Phytosanitary Risk Management	12
2.1.8 Inspection and Diagnostics.....	13
2.1.9 Documentation and Identification.....	13
2.1.10 Quality Assurance and Program Review	13
2.1.11 Non-compliance and Corrective Measures	14
3. Post-entry Quarantine	14
4. Evaluation and Approval of a Certification Program.....	15
5. Bilateral Workplans.....	15
Annex 1: Stone and Pome Fruit Tree Pests.....	16
Table 1: Virus pests of stone fruit.....	17
Table 2: Virus pests of pome fruits.....	24
Table 3: Fungal pathogens (incl. Chromista) of stone and pome fruit trees.....	27
Table 4: Bacterial pathogens of stone and pome fruit trees	39

Table 5: Nematode pests of stone and pome fruit trees	45
Table 6: Arthropod pests of stone and pome fruit trees.....	56
Annex 2:Grapevine Pests	68
Table 1: Virus pests of grapevine.....	69
Table 2: Fungal pathogens (incl. Chromista) of grapevine	77
Table 3. Bacterial pathogens of grapevine.....	85
Table 4: Nematode pests of grapevine	89
Table 5: Arthropod pests of grapevines	98

Review

NAPPO Regional Standards for Phytosanitary Measures (RSPMs) are subject to periodic review and amendment. This standard was last reviewed in 2022. A review of a NAPPO standard may be initiated at any time upon request of a NAPPO member country. The next review of RSPM 35 is scheduled for 2026.

Approval

This Standard was approved by the North American Plant Protection Organization (NAPPO) Executive Committee on March 9, 2022, and is effective from this date.

Virtual approval of NAPPO Products

Given the current travel restrictions brought about by the COVID-19 pandemic, the NAPPO Management Team unanimously endorsed a temporary process for virtual approval of its products.

Beginning in January 2021 and until further notice, this statement will be included with each approved NAPPO product in lieu of the Executive Committee original signature page.

Regional standard for phytosanitary measures 35 – ***Guidelines for the Movement of Propagative Plant Material of Stone Fruit, Pome Fruit, and Grapevine into a NAPPO Member Country*** – was approved by the North American Plant Protection Organization (NAPPO) Executive Committee – see approval dates below each signature - and is effective from the latest date below.

Approved by:

NAPPO Executive Committee members for Canada (Greg Wolff, CFIA) on March 8, 2022, the United States (Ibrahim M. Shaqir, APHIS PPQ) on March 08, 2022, and Mexico (Francisco Ramírez y Ramírez, SENASICA-DGSV) on March 9, 2022. Electronic copies of approval emails from each Executive Committee member have been archived by the NAPPO Secretariat.

Stephanie Bloem

Stephanie Bloem
NAPPO Executive Director

Implementation

No implementation plans required.

Amendment Record

Amendments to this Standard will be dated and filed with the NAPPO Secretariat.

Distribution

This standard is distributed by the NAPPO Secretariat to the Industry Advisory Group (IAG), the International Plant Protection Convention (IPPC) Secretariat, and to other Regional Plant Protection Organizations (RPPOs).

INTRODUCTION

Scope

This standard describes guidelines for the importation of propagative plant material of stone fruit, pome fruit and grapevine into NAPPO member countries, and the movement of this material among these countries. Propagative plant material is defined as plants or plant parts for planting or multiplication (RSPM 5 (*NAPPO Glossary of phytosanitary terms*)). For the scope of this standard, propagative plant material includes grafted plants, rooted plants, cuttings, budwood/scion, rootstocks, tissue culture (*in vitro*) and, where applicable, seeds. The pests specifically dealt with in this standard are bacteria, phytoplasmas, viruses, and virus-like agents and, where applicable, arthropods, fungi, and nematodes. These pests include those that pose a direct risk to their host, as well as those which act as potential vectors of a secondary pest. This standard does not address abiotic disorders, varietal trueness-to-type, quality grades and standards, or soil or fruit associated with the material.

References

- Good plant protection practice – EPPO PP 2/1(2), 2003.
Good plant protection practice – Grapevine EPPO PP 2/23(1), 2002.
Good plant protection practice – Pome fruits EPPO PP 2/18(1), 1999.
Good plant protection practice – Stone fruits EPPO PP 2/33(1), 2004.
Jelkmann, W. 2004. *International Working Group on Fruit Tree Viruses: Detection of virus and virus-like diseases of fruit trees*. Acta Horticulturae 657:575-596.
Thompson, D.A. 1998. The Role of NAPPO in Fruit Crop Virus-Testing and Certification. Acta Horticulturae 472:100.
- ISPM 2.** 2016. *Framework for pest risk analysis*. Rome, IPPC, FAO.
ISPM 4. 2017. *Requirements for the establishment of pest free areas*. Rome, IPPC, FAO.
ISPM 5. 2021. *Glossary of phytosanitary terms*. Rome, IPPC, FAO.
ISPM 6. 2018. *Surveillance*. Rome, IPPC, FAO.
ISPM 7. 2016 *Phytosanitary certification system*. Rome, IPPC, FAO.
ISPM 8. 2021. *Determination of pest status in an area*. Rome, IPPC, FAO.
ISPM 10. 2016. *Requirements for the establishment of pest free places of production and pest free production sites*. Rome, IPPC, FAO.
ISPM 11. 2017. *Pest risk analysis for quarantine pests*. Rome, IPPC, FAO.
ISPM 12. 2017. *Phytosanitary certificates*. Rome, IPPC, FAO.
ISPM 14. 2017. *The use of integrated measures in a systems approach for pest risk management*. Rome, IPPC, FAO.
ISPM 29. 2017. *Recognition of pest free areas and areas of low pest prevalence*. Rome, IPPC, FAO.
ISPM 36. 2012. *Integrated measures for plants for planting*. Rome, IPPC, FAO.
- RSPM 3.** 2017. *Requirements for importation of potatoes into a NAPPO member country on hold*. Raleigh, NC, USA.
RSPM 5. 2021. *NAPPO glossary of phytosanitary terms*. Raleigh, NC, USA. NAPPO.
RSPM 9. 2021. *Authorization of laboratories for phytosanitary testing*. Raleigh, NC, USA. NAPPO.
RSPM 19. 2012. *Guidelines for bilateral workplans*. Raleigh, NC, USA. NAPPO.
RSPM 24. 2013 *Integrated pest risk management measures for the importation of plants for planting into NAPPO member countries*. Raleigh, NC, USA. NAPPO.

Definitions

Definitions of phytosanitary terms used in this standard can be found in RSPM 5 (NAPPO *glossary of phytosanitary terms*) and in ISPM 5 (*Glossary of phytosanitary terms*).

Stone fruits: Species belonging to the genus *Prunus*, including all its varieties, cultivars, and hybrids.

Pome fruits: Species belonging to the genera of *Malus*, *Pyrus*, *Cydonia*, and *Chaenomeles*, including all their varieties, cultivars, and hybrids.

Grapevines: Species of the genus *Vitis*, including all varieties, cultivars, and hybrids,

SPFTG: propagative plant material of **Stone and Pome Fruit Trees and Grapevines**.

Outline of Requirements

This standard outlines a systems approach for mitigating the risk of pest introductions associated with the international movement of SPFTG, without undue restriction of trade. A systems approach is achieved through a combination of phytosanitary measures, two or more of which act independently of each other, to prevent the entry, establishment and spread of associated pests. Section 1 (General Requirements) of this standard addresses the pest risk assessment and phytosanitary measures for pest risk management. Section 2 (Specific Requirements) identifies and describes the components of a SPFTG certification program primarily designed to control phytoplasmas, viruses, and virus-like agents spread by infected propagative material. Certification programs may also be applicable to other types of pests.

Background

This standard deals specifically with bacteria, phytoplasmas, viruses, and virus-like agents and, where applicable, arthropods, fungi, and nematodes of SPFTG. Economic impacts of these pests on commodities covered by this standard include delayed maturity; increased agricultural inputs; reduced growth, yield, and fruit quality; graft incompatibility; plant mortality; as well as potential impacts on trade. The pests addressed in this standard may also affect other commodities or natural ecosystems, with varying economic impacts. In addition to direct effects on the plants, certain nematodes and arthropods are vectors of SPFTG diseases. SPFTG including cuttings, rootstocks, and tissue cultures pose a high risk for introducing pests. Traditional phytosanitary measures used to reduce the risk of pest introduction include prohibition, quarantine restrictions, entry point inspections, appropriate treatment and therapies, and post-entry quarantines. Competitiveness of these commodities in the worldwide market requires development and exchange of new varieties and germplasm. The increased volume of trade in SPFTG now occurring between and within countries has increased the risk of pest introduction and spread.

Systems approaches for pest risk management should be developed in accordance with ISPM 14 (*The use of integrated measures in a systems approach for pest risk management*), RSPM 24 (*Integrated pest risk management measures for the importation of plants for planting into NAPPO member countries*) and ISPM 36 (*Integrated measures for plants for planting*).

Systems approaches can provide alternatives to the application of single-point mitigations such as phytosanitary treatments or more restrictive measures such as prohibition. Systems approaches provide the opportunity to consider both pre-harvest and post-harvest procedures and measures that

may contribute to the effective management of pest risk.

A systems approach requires two or more measures that are independent of each other and may also include any number of measures that are dependent on each other. An advantage of a systems approach is its ability to address variability and uncertainty in the system by modifying the number and strength of phytosanitary measures to meet the appropriate level of phytosanitary protection.

Certification programs used to control virus diseases are good examples of a systems approach at work. Various independent measures such as virus-testing, field inspection, isolation distances, and vector control all work together to minimize the introduction and spread of pests.

The objectives of this standard are to:

- prevent the entry, establishment, and spread of quarantine pests into NAPPO member countries
- mitigate the impact of regulated non-quarantine pests within NAPPO member countries
- facilitate trade into and within the NAPPO region
- promote the use of systems approaches and good plant protection practices as the basis for the development of certification programs for international exchange of SPFTG.

1. GENERAL REQUIREMENTS

1.1 Stone and Pome Fruit and Grapevine Pests

A comprehensive list of pests associated with SPFTG along with their phytosanitary status in each NAPPO member country can be found in Annexes 1 and 2. While not all these pests are regulated by NAPPO member countries, they may be considered as pests of concern to SPFTG production. These pest lists are useful references for the targeted testing, therapy, and transfer of clean germplasm among NAPPO countries.

1.2 Pest Risk Analysis

All Pest Risk Analyses (PRAs) should be performed in accordance with ISPM 2 (*Framework for pest risk analysis*) and ISPM 11 (*Pest risk analysis for quarantine pests*). Application of phytosanitary measures should be based on the results of a PRA. The pests in Annexes 1 and 2 may be classified as regulated pests, depending on their presence or absence and their official status in each country.

1.3 Phytosanitary Measures for Pest Risk Management

Phytosanitary measures for pest risk management should be used to prevent the entry, establishment, and spread of quarantine pests and mitigate the economic impact of regulated non-quarantine pests.

The importation and movement of SPFTG is subject to the application of integrated measures in a systems approach in accordance with ISPM 14 (*The use of integrated measures in a systems approach for pest risk management*).

Following a PRA, phytosanitary measures to mitigate the risks associated with importing propagative plant material into a NAPPO member country are identified. The phytosanitary measures described

below may be combined to reach the appropriate level of phytosanitary protection of the importing country. For the purposes of this standard “small quantities” of plant material are considered to be 100 plants or less, but the definition of “small quantities” will ultimately be left to the discretion of the National Plant Protection Organizations (NPPOs).

Phytosanitary measures and procedures such as inspection, fumigation, chemical sprays, hot water dips, biological control, cold treatment, and others, may be applied to plant material, growing media, or packaging. Section 3.0 “Post-entry Quarantine” identifies the criteria appropriate for the post-entry quarantine of these plants.

Appropriate phytosanitary measures for pest risk management may also depend on the end use of the material. The various categories of use of SPFTG are outlined below:

1.3.1 *Research Purposes and Subsequent Destruction*

This option is appropriate for SPFTG that do not necessarily come from an official certification program or that may be infested with pests. Plants are maintained under quarantine conditions authorized by the NPPO to prevent the establishment and spread of regulated pests. Plant material may be inspected, tested, or treated for pests of concern before importation or after entry, at the discretion of the NPPO. Plant material, growing media, or packaging must be disposed of as instructed by the NPPO. This option is only suitable for small quantities of plant material.

1.3.2 *Quarantine, Testing and Treatment at NPPO-authorized Facilities and Subsequent Distribution*

This option is appropriate for SPFTG that do not come from an official certification program or that may be infested with pests. Plants are imported for quarantine, testing and treatment in a post-entry quarantine station approved by the NPPO of the importing country. Detected regulated pests should be eliminated from the plants before their release from quarantine conditions. This option is only suitable for small quantities of plant material.

1.3.3 *Quarantine at Importers’ Premises and Subsequent Distribution*

This option is appropriate for SPFTG that do not come from an official certification program recognized by the NPPO of the importing country. Plants are planted under quarantine conditions on the importer’s premises, as required by the NPPO. The NPPO or authorized entity should test and/or visually examine or treat, as appropriate, for regulated pests before release from the quarantine conditions. This option may only be suitable for small quantities of plant material, at the discretion of the NPPO.

1.3.4 *Plants originating from an Official Certification Program*

SPFTG may be imported into a NPPO member country if produced under an official certification program that has been evaluated using this standard and authorized by the NPPO of the importing country. The NPPO of the importing country may perform inspections either in the country of origin or on imported plants, including testing samples for the presence of pests. The NPPO of the importing country may require post-entry quarantine conditions.

1.3.5 Plants originating from a Pest-free Area, Pest Free Place of Production, or Pest Free Production Site

This option is usually not appropriate for pests such as viruses, viroids, and bacteria, where inspection is inadequate and comprehensive sampling and monitoring are impractical for determining pest-free status. SPFTG may be certified free from specific pests for entry into NAPPO member countries based on absence of these pests in the exporting site or area as outlined in ISPM 4 (*Requirements for the establishment of pest free areas*), ISPM 10 (*Requirements for establishment of pest-free places of production and pest-free production sites*), and ISPM 29 (*Recognition of pest-free areas and areas of low pest prevalence*). The NPPO of the importing country may perform inspections and may take samples to test for the presence of pests. The NPPO of the importing country may require post-entry quarantine conditions.

1.3.6 Prohibition

If no satisfactory phytosanitary measure to reduce risk to an acceptable level can be found, the final option may be to prohibit importation of SPFTG.

1.4 Documentation Requirements

A phytosanitary certificate or an equivalent official document should be issued by the exporting country according to the requirements of the NPPO of the importing country. A permit to import must be obtained by the importer, if required by their NPPO.

2. SPECIFIC REQUIREMENTS

2.1 Stone and Pome Fruit and Grapevine Certification Programs

This standard outlines the essential elements of a certification program to mitigate the risk of introduction and spread of SPFTG pests listed in the Annexes.

The certification program should be carried out by the NPPO, or an entity authorized by the NPPO. The certification program should clearly define phytosanitary requirements such as terminology, testing, eligibility, the nomenclature of generation levels, agricultural management, isolation and sanitation requirements, inspection and re-testing, documentation and reporting, identification and labeling, quality controls and monitoring, non-compliance and remedial measures, and criteria for post-entry quarantine.

2.1.1 Program Administration

The certification program should be administered by an NPPO, or an entity recognized by an NPPO to administer certification programs (herein referred to as an “authorized entity”).

The program should clearly and comprehensively specify the roles and responsibilities of the NPPO, the program participants, any authorized entities (e.g., laboratories involved in testing), and any organizations involved in certification activities.

The NPPO and any authorized entities should ensure that all administration, inspection, certification, and laboratory personnel meet appropriate training, experience, educational and proficiency requirements. Authorized entities should be prepared to supply, upon request, this information to their NPPO.

Authorized entities should notify their NPPO prior to making changes to the certification program or deviating from program requirements, and before plants produced under such modifications are exported.

The exporting country's NPPO or authorized entity must inform the importing country's NPPO of changes to the certification program or diagnostics before plants produced under the modified program are exported.

2.1.2 Terminology

The certification program should define all terminology specific to the certification program using sufficient detail to ensure clear understanding of the certification requirements. The terminology used by NAPPO member countries for similar purposes should be harmonized to the greatest extent possible.

2.1.3 Diagnostics

Diagnostics for pests in this standard include but are not limited to:

- sample processing for recovery or isolation and identification
- identification utilizing morphological characteristics
- determinations utilizing indicator plants
- serological tests such as Enzyme Linked Immuno-Sorbent Assays (ELISA) and,
- assays based on nucleic acid amplification by various Polymerase Chain Reaction (PCR) methodologies.

Determinations based on High Throughput Sequencing (HTS) methodologies may also be considered but should follow the recommendations outlined in CPM-14; 2019 Recommendation R-08 (*Preparing to use high-throughput sequencing (HTS) technologies as a diagnostic tool for phytosanitary purposes*).

Two or more of the aforementioned types should be considered for a more accurate diagnosis.

The NPPO or authorized entity will deliver diagnostic services. If laboratories are used, they should be authorized by the NPPO in accordance with RSPM 9 (*Authorization of laboratories for phytosanitary testing*).

Upon request, the NPPO or authorized entity of the exporting country must provide the NPPO of the importing country with the diagnostic results, methodology and a list of pests regulated in the certification program in the exporting country.

The NPPO or authorized entity of the exporting country must notify the NPPO of the importing country of proposed changes to diagnostics being used by the NPPO or authorized entity in the exporting country. Such changes must be communicated and approved by the NPPO or authorized entity of the importing country before plants produced under the modified program are exported.

2.1.4 Eligibility

Potential certification program participants should file an application with their NPPO or authorized entity. Eligibility is conferred by the NPPO or authorized entity if the conditions of the certification program have been met.

The certification program should specify eligibility for plant material used in the program. Eligibility requirements also must be met when plants are purchased for export. Plant brokers must ensure the traceability of exported consignments to approved places of production.

2.1.5 Generation Levels

Generation levels represent successive generations of propagation from the original tested material and as such, act as a measure of the phytosanitary risk associated with plants in a certification program. Additional phytosanitary measures may be applied depending on the generation. The certification program should clearly define generation levels. Eligibility criteria should be established at each level, including nomenclature, propagation and pest management measures, and the number of generations removed from the original tested material. It is strongly recommended that the levels within a certification program be identified as Generation 1, 2, 3, 4, etc.

2.1.6 Agricultural Management

The certification program should define agricultural management requirements for hosts of pests or pest vectors within the field and buffer zones.

All stone and pome fruit trees and grapevines in the certification program should be kept in good horticultural condition by following good agricultural practices for their region. Good agricultural practices are explained in the Food and Agriculture Organization (FAO) website at <http://www.fao.org/3/a-i6677e.pdf>.

Where they exist, it is recommended to employ best management practices, for example:

- principles of good plant protection practice – EPPO PP 2/1(2), 2003
- good plant protection practice – Grapevine EPPO PP 2/23(1), 2002
- good plant protection practice – Pome fruits EPPO PP 2/18(1), 1999 and,
- good plant protection practice – Stone fruits EPPO PP 2/33(1), 2004.

2.1.7 Phytosanitary Risk Management

The isolation and containment requirements of the certification program will vary according to the generation level of the propagative plant material and should be based on the epidemiology and biology of the pests and their vectors present in the certification area. The certification program should specify the minimum distance from non-certified hosts, and acceptable cover crops and weed control measures required to reduce alternate pest hosts to acceptable levels.

The certification program should specify pest management measures including vector suppression and the control of pollen-borne viruses required to adequately protect plants produced under the program from exposure to pests.

The certification program should specify the measures to mitigate the risks associated with the movement of soil or water, other growing media or plant products potentially infested with vectors or pests to acceptable levels.

The certification program should specify crop rotation and fallow requirements between host crops, and chemical control requirements for a site being used to produce plants under the certification program.

2.1.8 *Inspection and Diagnostics*

The certification program should specify the inspection and diagnostics requirements throughout all levels of the program.

Plants in the certification program should be inspected during the growing season at times appropriate for detecting disease symptoms and determining the presence of insects or other pest vectors using appropriate methods.

The certification program should specify:

- procedures to be undertaken upon suspicion of pest infestation
- procedures to be undertaken upon confirmation of pest infestation
- notification and inspection requirements when selling or purchasing certified material
- sampling procedures and diagnostics for pests at each generation level
- the diagnostic methodology(ies) to be used, and
- inspection requirements including review of maps of places of production and production sites, variety labelling practices, approval of new places of production and production sites, and any deviations between inventory, sales, and purchases.

2.1.9 *Documentation and Identification*

The NPPO or authorized entity must document all inspection, certification, and diagnostic activities to ensure the eligibility of the program participants, their places of production and production sites, and the plants produced under this certification program. These documents must be available, upon request, to the NPPOs of the importing and exporting country for audit, traceback, trace forward and other regulatory purposes.

The certification program should include a system approved by the NPPO or authorized entity to document and identify plants during growth, post-harvest, and at sale to ensure traceability. The system should at least record the generation level, the year of propagation, the program participant, geographic location of the production field, location of certified plants within the production field, the variety and rootstock, and the purchaser's identity.

Purchases and sales of plants produced under the certification program, previous cropping history for production sites, and production site maps should be retained by the program participants for a period specified by their NPPO or authorized entity.

2.1.10 *Quality Assurance and Program Review*

The NPPO or authorized entity of the exporting country should ensure the validity and reliability of their certification program through periodic program audits and reviews. Records and any other

supporting documentation must be maintained for any such audits or reviews. In addition, the place of production should adhere to the program's quality management requirements as indicated by the NPPO or authorized entity. Fundamental changes to program delivery or deviations on agreed-upon phytosanitary measures must be communicated in advance and approved by the NPPO of the importing country.

The NPPO of the importing country should review and/or audit the certification program of the exporting country to ensure it continues to meet the certification standards and their import requirements. This may be done on a periodic basis, or in response to certification program or pest status changes, or because of evidence of non-compliance. It should include testing of imported plant material, site visits and/or review of the entire export certification program. Detection of pests or vectors controlled under the program or deficiencies in documentation may indicate that the integrity of the export certification program is compromised.

2.1.11 Non-compliance and Corrective Measures

The certification program should specify the consequences of non-compliance. In addition, the corrective measures should be specified to enable a suspended or de-certified program participant, production area or variety to become eligible for re-certification or reinstatement.

3. Post-entry Quarantine

The NPPO or authorized entity of the importing country may require post-entry quarantine for imported SPFTG. Post-entry quarantine may take place at an NPPO-approved facility and should follow the guidelines outlined in ISPM 34 (*Design and operation of post-entry quarantine stations for plants*). The post-entry requirements should be based on the level of risk determined by the biology of the pests of concern, including their host range, their means of natural spread, and the likelihood of transmission by local vectors.

Post-entry quarantine criteria should specify:

- the roles and responsibilities of the NPPO of the importing country, authorized entities, and the importer
- agricultural practices to promote plant growth and the detection of pests
- isolation and suppression measures to control pest vectors and prevent the movement of pests within and outside the post-entry quarantine area
- measures within the post-entry quarantine area to reduce alternate hosts and pest vectors, such as weed control and buffer zones
- soil and plant treatment, vector surveillance and suppression, facility design and other criteria to be met before a facility, production site, or production area becomes suitable for post-entry quarantine
- the requirements for movement and sanitation of agricultural equipment and personnel into and from the post-entry quarantine area
- containment, security, and access restrictions to the imported plants
- handling and disposal of pruning waste and all other articles capable of harboring or spreading pests
- inspection, sampling, and diagnostics to determine the presence of pests in the imported plants
- conditions under which the imported plants would be moved, removed, or released from post-entry quarantine, and

- sanitation and subsequent use restrictions of a post-entry quarantine area.

4. Evaluation and Approval of a Certification Program

Prior to allowing importation of SPFTG, the NPPO of the importing country should evaluate the certification program of the exporting country, including a documentation review, a site visit, and/or testing of plants by the NPPO of the importing country to ensure they meet the certification program standards.

Following approval of the certification program additional temporary restrictions such as pre-clearance testing, and post-entry quarantine may be used.

5. Bilateral Workplans

The NPPOs of the exporting and importing countries may decide that a bilateral workplan is necessary to elaborate on these guidelines. Guidelines for the development of bilateral workplans are provided in RSPM 19 (*Guidelines for Bilateral Workplans*). Modifications to these guidelines should be technically justified.

Annex 1: Stone and Pome Fruit Tree Pests

Note: Synonyms for virus names in Tables 1 and 2 can be found in Appendix 2 of RSPM 25 archived in the NAPPO website.

LEGEND FOR SYMBOLS USED IN TABLES

Presence or absence unless otherwise noted conform to the categories listed in ISPM 8: 2021 (*Determination of pest status in an area*). For ease of reference, alphanumeric designations have been added here.

Ab1: Absent: pest not recorded
Ab2: Absent: the entire country is pest-free
Ab3: Absent: pest records invalid
Ab4: Absent: pest no longer present
Ab5: Absent: pest eradicated

P1: Present: widely distributed
P2: Present: not widely distributed and not under official control
P3: Present: not widely distributed and under official control
P4: Present: at low prevalence
P5: Present: except in specified pest-free areas
P6: Present: transient
P7: Present: not associated with host (NAPPO category)

Table 1: Virus pests of stone fruit

Table 2: Virus pests of pome fruit

Table 3: Fungal pathogens (incl. Chromista) of stone and pome fruit trees

Table 4: Bacterial pathogens of stone and pome fruit trees

Table 5: Nematode pests of stone and pome fruit trees

Table 6: Arthropod pests of stone and pome fruit trees

Table 1: Virus pests of stone fruit.

Last updated December 2021.

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
						CAN	USA	MEX
Amasya cherry disease associated virus	ACDaV	<i>Chrysoviridae</i>	<i>Chrysovirus</i>	<i>P. avium</i>	Covelli <i>et al.</i> , 2004	Ab1	Ab1	Ab1
American plum line pattern virus	APLPV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i> , <i>P. salicina</i>	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Apple chlorotic leaf spot virus	ACLSV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Apple mosaic virus	ApMV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P7	P2	Ab3
Apricot latent ringspot virus	ALRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i>	Gentit <i>et al.</i> , 2001	Ab1	Ab1	Ab1
Apricot latent virus	ApLV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. salicina</i> , <i>P. persica</i>	Grimová and Rysanek, 2012	Ab1	Ab1	Ab1
Apricot pseudo-chlorotic leaf spot virus	APCLSV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i> , <i>P. salicina</i>	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Apple scar skin viroid	ASSVd	<i>Pospiviroidae</i>	<i>Apscaviroid</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. persica</i>	Kaponi <i>et al.</i> , 2013	P7	P2	Ab1
Apricot vein clearing associated virus	AVCaV	<i>Betaflexiviridae</i>	<i>Prunevirus</i>	<i>P. armeniaca</i>	Elbeaino <i>et al.</i> , 2014	Ab1	Ab1	Ab1
Arabis mosaic virus	ArMV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. avium</i> , <i>P. persica</i>	Hadidi <i>et al.</i> , 2011	P7	P2	Ab3
Asian <i>Prunus</i> virus 1	APV1	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>Prunus</i> spp.	Marini <i>et al.</i> , 2009	P2	Ab1	Ab1
Asian <i>Prunus</i> virus 2	APV2	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>Prunus</i> spp.	Marais <i>et al.</i> , 2016	P2	Ab1	Ab1
Asian <i>Prunus</i> virus 3	APV3	<i>Unassigned</i>	<i>Unassigned</i>	<i>Prunus</i> spp.	Marais <i>et al.</i> , 2016	P2	Ab1	Ab1
Carnation Italian ringspot	CIRV	<i>Tombusviridae</i>	<i>Tombusvirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	Ab1	Ab3	Ab1

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
						CAN	USA	MEX
virus								
Caucasus prunus virus	CPrV	<i>Betaflexiviridae</i>	<i>Prunevirus</i>	<i>Prunus</i> spp.	Marais <i>et al.</i> , 2015b	Ab1	Ab1	Ab1
Cherry associated luteovirus	ChALV	<i>Luteoviridae</i>	<i>Luteovirus</i>	<i>P. avium</i>	Lenz <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Cherry green ring mottle virus	CGRMV	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P1	P2	Ab1
Cherry leaf roll virus	CLRV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P3	P2	Ab1
Cherry mottle leaf virus	CMLV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. persica</i>	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Cherry necrotic rusty mottle virus	CNRMV	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i>	Villamor <i>et al.</i> , 2015	P1	P2	Ab1
Cherry rasp leaf virus	CRLV	<i>Secoviridae</i>	<i>Cheravirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Cherry robiovirus 5	CRV-5	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>P. avium</i>	Wu <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Cherry rusty mottle associated virus	CRMV or CRMaV	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P1	P2	Ab1
Cherry twisted leaf associated virus	CTLaV	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Cherry virus A	CVA	<i>Betaflexiviridae</i>	<i>Capillovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i>	Hadidi <i>et al.</i> , 2011	P1	P1	Ab1
Cherry virus F	CVF	<i>Secoviridae</i>	<i>Fabavirus</i>	<i>P. avium</i>	Koloniuk <i>et al.</i> , 2020; Maliogka <i>et al.</i> , 2018	P2	Ab1	Ab1
Cherry virus Trakiya	CVT	Unassigned <i>Picornavirales</i> (proposed)	Unassigned	<i>P. avium</i>	Milusheva <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Cherry virus Turkey	CVTR	<i>Betaflexiviridae</i>	<i>Robiogovirus</i>	<i>P. avium</i>	Çağlayan <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Cherry yellow spot-associated virus	CYSaV	Unassigned in Order <i>Tymovirales</i>	<i>Gratylivirus</i>	<i>P. davidiana</i>	Hou <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Cucumber green mottle mosaic virus	CGMMV	<i>Virgaviridae</i>	<i>Tobamovirus</i>	<i>P. armeniaca</i>	Chitambar, 2018; Tian <i>et al.</i> , 2014	P7	P2	Ab1
Cucumber mosaic virus	CMV	<i>Bromoviridae</i>	<i>Cucumovirus</i>	<i>P. armeniaca</i>	Hadidi <i>et al.</i> , 2011	P7	P2	P2

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
						CAN	USA	MEX
Hop stunt viroid	HSVd	<i>Popsiviroidae</i>	<i>Hostuviroid</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. salicina</i>	Hadidi <i>et al.</i> , 2011	P2	P2	P2
Little cherry virus 1	LChV-1	<i>Closteroviridae</i>	<i>Velarivirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P1 (P3 in BC)	P2	Ab1
Little cherry virus 2	LChV-2	<i>Closteroviridae</i>	<i>Ampelovirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P1 (P3 in BC)	P2	Ab1
Mume virus A	MuVA	<i>Betaflexiviridae</i>	<i>Capillovirus</i>	<i>Prunus</i> spp.	Marais <i>et al.</i> , 2018	Ab1	Ab1	Ab1
Nectarine stem pitting associated virus	NSPaV	<i>Luteoviridae</i>	<i>Luteovirus</i>	<i>P. persica</i> var. <i>nectarina</i>	Bag <i>et al.</i> , 2015	Ab1	Ab4	Ab1
Nectarine virus M	NeVM	<i>Tymoviridae</i>	<i>Marafavirus</i>	<i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i>	Villamor <i>et al.</i> , 2016	Ab1	P2	Ab1
Peach associated luteovirus	PaLV	<i>Luteoviridae</i>	<i>Luteovirus</i>	<i>P. persica</i>	Wu <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Peach chlorotic mottle virus	PcCMV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>P. persica</i>	James <i>et al.</i> , 2007	P2	Ab1	Ab1
Peach latent mosaic viroid	PLMVd	<i>Avsunviroidae</i>	<i>Pelamoviroid</i>	<i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i>	Hadidi <i>et al.</i> , 2011	P1	P2	P2
Peach leaf pitting-associated virus	PLPaV	<i>Secoviridae</i>	<i>Fabavirus</i>	<i>P. persica</i>	He <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Peach mosaic virus	PcMV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarine</i> , <i>P. salicina</i>	Hadidi <i>et al.</i> , 2011; James <i>et al.</i> , 2006	Ab1	P2	Ab1
Peach rosette mosaic virus	PRMV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. domestica</i> , <i>P. persica</i> , <i>P. salicina</i>	Hadidi <i>et al.</i> , 2011	P2	P2	Ab1
Peach virus D	PeVD	<i>Tymoviridae</i>	<i>Marafavirus</i>	<i>P. persica</i>	Igori <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Peach virus M	PeVM	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>P. persica</i>	De La Torre-	Ab1	Ab1	P2

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
						CAN	USA	MEX
					Almaráz <i>et al.</i> , 2019			
Petunia asteroid mosaic virus	PeAMV	<i>Tombusviridae</i>	<i>Tombusvirus</i>	<i>P. avium</i>	Hadidi <i>et al.</i> , 2011	P2	Ab1	Ab1
Plum bark necrosis stem pitting-associated virus	PBNPaV	<i>Closteroviridae</i>	<i>Ampelovirus</i>	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i>	Hadidi <i>et al.</i> , 2011	Ab1	P2	Ab1
Plum pox virus	PPV	<i>Potyviridae</i>	<i>Potyvirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P3 (Ab5 for NS)	Ab5	Ab3
Prune dwarf virus	PDV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P1	P2	Ab1
<i>Prunus</i> geminivirus A	PrGVA	<i>Geminiviridae</i>	<i>Grablovirus</i>	<i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. salicina</i>	Al-Rwahnih <i>et al.</i> , 2018	Ab1	P2	Ab1
<i>Prunus</i> necrotic ringspot virus	PNRSV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P1	P2	Ab3
<i>Prunus</i> virus F	PrVF	<i>Secoviridae</i>	<i>Fabavirus</i>	<i>P. avium</i>	Villamor <i>et al.</i> , 2017	P2	P2	Ab1
<i>Prunus</i> virus T	PrVT	<i>Betaflexiviridae</i>	<i>Tepovirus</i>	<i>Prunus</i> spp.	Marais <i>et al.</i> , 2015a	Ab1	Ab1	Ab1
Raspberry ringspot virus	RRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. domestica</i>	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Sowbane mosaic virus	SoMV	<i>Solemoviridae</i>	<i>Sobemovirus</i>	<i>P. domestica</i>	Németh, 1986	P7	P2	Ab1
Stocky prune virus	StPV	<i>Secoviridae</i>	<i>Cheravirus</i>	<i>P. domestica</i> , <i>P. persica</i>	Candresse <i>et al.</i> , 2006	Ab1	Ab1	Ab1
Strawberry latent ringspot virus	SLRSV	<i>Secoviridae</i>	<i>unassigned</i>	<i>P. avium</i> , <i>P. persica</i> , <i>P. salicina</i>	Tang <i>et al.</i> , 2013	P2	P2	Ab3
Tobacco mosaic virus	TMV	<i>Virgaviridae</i>	<i>Tobamovirus</i>	<i>P. avium</i> , <i>P. cerasus</i> , <i>P. domestica</i>	Németh, 1986	P7	P2	P2
Tobacco necrosis virus A	TNVA	<i>Tombusviridae</i>	<i>Alphanecrovirus</i>	<i>P. armeniaca</i> , <i>P. cerasus</i> , <i>P. domestica</i>	Németh, 1986	P7	P2	Ab1
Tobacco necrosis virus D	TNVD	<i>Tombusviridae</i>	<i>Bentanecrovirus</i>	<i>P. armeniaca</i> , <i>P. cerasus</i> , <i>P. domestica</i>	Németh, 1986	P7	P7	Ab1
Tobacco ringspot virus	TRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. avium</i> , <i>P. incisa</i> , <i>P.</i>	Hadidi <i>et al.</i> , 2011	P2	P2	P2

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
						CAN	USA	MEX
				<i>persica, P. serrula, P. serrulata</i>				
Tomato black ring virus	TBRV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>Prunus</i> spp.	Hadidi <i>et al.</i> , 2011	P7	Ab1	Ab1
Tomato bushy stunt virus	TBSV	<i>Tombusviridae</i>	<i>Tombusvirus</i>	<i>P. avium, P. domestica, P. salicina</i>	Ogawa, 1995	P2	P2	Ab3
Tomato ringspot virus	ToRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>P. avium, P. dulcis, P. persica</i>	Ogawa, 1995	P1	P2	Ab3

REFERENCES

- Al-Rwahnih, M., Alabi, O. J., Westrick, N. M. & Golino, D.** 2018. *Prunus* Geminivirus A: A novel Grablovirus infecting *Prunus* spp. *Plant Disease* 102: 1246-1253.
- Bag, S., Al Rwahnih, M., Li, A., Gonzalez, M., Rowhani, A., Uyemoto, J. & Sudarshana, M.** 2015. Detection of a new Luteovirus in imported nectarine trees: A case study to propose adoption of metagenomics in post-entry quarantine. *Phytopathology* 105(6): 840-846.
- Çağlayan, K., Roumi, V., Gazel, M., Elçi, E., Acioğlu, M., Plesko, I. M., Reynard, J. S., Maclot, F. & Massart, S.** 2019. Identification and characterization of a novel Robigovirus species from sweet cherry in Turkey. *Pathogens*, 8(2): 57.
- Chitambar, J.** 2018. *Cucumber green mottle mosaic virus*. California Department of Food and Agriculture, Davis, California. Last accessed 5/6/2020, <https://blogs.cdfa.ca.gov/Section3162/?p=4314>.
- Covelli, L., Coutts, R. H. A., Serio, F. D., Citir, A., Açıkgöz, S., Hernández, C., Ragozzino, A. & Flores, R.** 2004. Cherry Chlorotic Rusty Spot and Amasya Cherry Diseases are associated with a complex pattern of Mycoviral-Like Double-Stranded Rnas. I. Characterization of a new species in the genus *Chrysovirus*. 85: 3389-3397.
- De La Torre-Almaráz, R., V. Pallás, & Sánchez-Navarro, J. A.** 2019. Molecular characterization of a new trichovirus from peach in Mexico. *Archives of Virology* 164:2617-2620.
- Elbeaino, T., Giampetruzzi, A., De Stradis, A. & Digiario, M.** 2014. Deep-sequencing analysis of an apricot tree with vein clearing symptoms reveals the presence of a novel *Betaflexivirus*. *Virus Research*, 181: 1-5.
- Gentit, P., Delbos, R.-P., Candresse, T. & Dunez, J.** 2001. Characterization of a new Nepovirus infecting apricot in Southeastern France: Apricot Latent Ringspot Virus. *European Journal of Plant Pathology* 107: 485-494.
- Grimová, L. & Rysanek, P.** 2012. Apricot Latent Virus – Review. *Horticultural Science*, 39(3): 144-148.
- Hadidi, A., Barba, M., Candresse, T. & Jelkmann, W.** 2011. Virus and Virus-Like diseases of Pome and Stone Fruits, American Phytopathological Society, St. Paul, MN, USA. 429p.
- He, Y., Cai, L., Zhou, L., Yang, Z., Hong, N., Wang, G., Li, S. & Xu, W.** 2017. Deep sequencing reveals the first Fabavirus infecting Peach. *Scientific Reports*, 7(11329): 1-11.
- Hou, Q., Han, T., Li, L., Wang, J., Yu, M., Zhang, S., Cao, M. & Yang, C.** 2019. The complete nucleotide sequence and genome organization of a novel virus of the Order Tymovirales isolated from *Prunus davidiana* (Carr.) Franch. In Liaoning, China. *Archives of Virology* 164: 1245-1248.
- Igori, D., Lim, S., Baek, D., Kim, S. Y., Seo, E., Cho, I. S., Choi, G. S., Lim, H. S. & Moon, J. S.** 2017. Complete nucleotide sequence and genome organization of Peach Virus D, a putative new member of the genus *Marafivirus*. *Archives of Virology* 162: 1769-1772.
- James, D., Varga, A. & Croft, H.** 2007. Analysis of the complete genome of Peach Chlorotic Mottle Virus: Identification of non-Aug start codons, In Vitro Coat Protein Expression, and Elucidation of Serological Cross-Reactions. *Archives of Virology* 152: 2207-2215.
- James, D., Varga, A., Croft, H., Rast, H., Thompson, D. & Hayes, S.** 2006. Molecular characterization, phylogenetic relationships, and specific detection of Peach Mosaic Virus. *Phytopathology* 96: 137-44.
- Kaponi, M., Sano, T. & Kyriakopoulou, P.** 2013. Natural Infection of sweet cherry trees with Apple Scar Skin Viroid. *Journal of Plant Pathology* 95: 429-433.
- Koloniuk, I., Přebilová, J., Fránová, J. & Špak, J.** 2020. Genomic characterization of *Malus domestica* Virus A (Mdova), a novel Velarivirus infecting apple. *Archives of Virology*. 165(2): 479-482.
- Lenz, O., Přebilová, J., Fránová, J., Koloniuk, I. & Špak, J.** 2017. Identification and characterization of a new member of the genus *Luteovirus* from cherry. *Archives of Virology* 162: 587-590.
- Maliogka, V.I., A. Minafra, P. Saldarelli, A. B. Ruíz-García, M. Glasa, N. Katis & A. Olmos.** 2018.

Recent advances on detection and characterization of fruit tree viruses using high-throughput sequencing technologies. *Viruses* 10(436): 1-23.

- Marais, A., Faure, C. & Candresse, T.** 2016. New insights into Asian *Prunus* viruses in the light of Ngs-Based full genome sequencing. *Plos One*, 11, E0146420.
- Marais, A., Faure, C., Mustafayev, E., Barone, M., Alioto, D. & Candresse, T.** 2015a. Characterization by deep sequencing of Prunus Virus T, a novel Tepovirus infecting *Prunus* species. *Phytopathology* 105: 135-140.
- Marais, A., Faure, C., Mustafayev, E. & Candresse, T.** 2015b. Characterization of new isolates of apricot Vein Clearing-Associated Virus and of a new *Prunus*-Infecting Virus: Evidence for recombination as a driving force in Betaflexiviridae Evolution. *Plos One*, 10, E0129469.
- Marais, A., Faure, C., Theil, S. & Candresse, T.** 2018. Molecular characterization of a novel species of *Capillovirus* from Japanese apricot (*Prunus mume*). *Viruses* 10(144): 1-8.
- Marini, D. B., Gibson, P. G. & Scott, S. W.** 2009. The complete nucleotide sequence of an isolate of Asian *Prunus* Virus 1 from peach [*Prunus persica* (L) Batch]. *Archives of Virology* 154: 1375-1377.
- Milusheva, S., Phelan, J., Piperkova, N., Nikolova, V., Gozmanova, M. & James, D.** 2019. Molecular analysis of the complete genome of an unusual virus detected in sweet cherry (*Prunus avium*) in Bulgaria. *European Journal of Plant Pathology* 153: 197-207.
- Németh, M. V.** 1986. *The Virus, Mycoplasma and Rickettsia Diseases of Fruit Trees*, Dordrecht, Martinus Nijhoff Publishers.
- Ogawa, J. M.** 1995. *Compendium of Stone Fruit Diseases*, American Phytopathological Society, St. Paul, MN, USA.
- Tang, J., Ward, L. I. & Clover, G. R. G.** 2013. The diversity of strawberry latent ringspot Virus in New Zealand. 97, 662-667.
- Tian, T., Posis, K., Maroon-Lango, C. J., Mavrodieva, V., Haymes, S., Pitman, T. L., & Falk, B.W.** 2014. First report of *Cucumber green mottle mosaic virus* on melon in the United States. *Plant Disease* 98(8):1163-1164.
- Villamor, D. E., Mekuria, T. A., Pillai, S. S. & Eastwell, K. C.** 2016. High-throughput sequencing identifies novel viruses in nectarine: Insights to the etiology of stem-pitting disease. *Phytopathology* 106: 519-27.
- Villamor, D. E., Susaimuthu, J. & Eastwell, K. C.** 2015. Genomic analyses of Cherry Rusty Mottle Group and Cherry Twisted Leaf-associated viruses reveal a possible new genus within the family Betaflexiviridae. *Phytopathology* 105: 399-408.
- Villamor, D. E. V., Pillai, S. S. & Eastwell, K. C.** 2017. High throughput sequencing reveals a novel Fabavirus infecting sweet cherry. *Archives of Virology* 162: 811-816.
- Wu, L., Liu, H., Bateman, M., Komorowska, B. & Li, R.** 2019. First identification and molecular characterization of a novel cherry Robigovirus. *Archives of Virology*, 164, 3103-3106.
- Wu, L. P., Liu, H. W., Bateman, M., Liu, Z. & Li, R.** 2017. Molecular Characterization of a novel Luteovirus from peach identified by high-throughput sequencing. *Archives of Virology* 162: 2903-2905.

Table 2. Virus pests of pome fruits

Last updated December 2021.

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE / ABSENCE		
						CAN	USA	MEX
Apple associated luteovirus	AaLV	<i>Luteoviridae</i>	<i>Luteovirus</i>	<i>Malus</i> spp.	Shen <i>et al.</i> , 2018	Ab1	P2	Ab1
Apple chlorotic leaf spot virus	ACLSV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>Cydonia</i> spp., <i>Chaenomeles</i> spp.	Hadidi <i>et al.</i> , 2011	P1	P2	Ab1
Apple dimple fruit viroid	ADFVd	<i>Pospiviroidae</i>	<i>Apscaviroid</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Apple fruit crinkle viroid	AFCVd	<i>Pospiviroidae</i>	<i>Apscaviroid</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Apple geminivirus	AGV	<i>Geminiviridae</i>	<i>Geminivirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Apple green crinkle associated virus	AGCaV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>Malus</i> spp., <i>Cydonia</i> spp.	Morelli <i>et al.</i> , 2017	P2	P2	Ab1
Apple hammerhead viroid-like RNA	AHVd	<i>Avsunviroidae</i>	<i>Pelamoviroid</i>	<i>Malus</i> spp.	Szostek <i>et al.</i> , 2018; Hadidi <i>et al.</i> , 2017	P2	P1	Ab1
Apple latent spherical virus	ALSV	<i>Secoviridae</i>	<i>Cheravirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Apple luteovirus 1	ALV	<i>Luteoviridae</i>	<i>Luteovirus</i>	<i>Malus</i> spp.	Lim <i>et al.</i> , 2019	P2	P2	Ab1
Apple mosaic virus	ApMV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Malus</i> spp., <i>Pyrus</i> spp.	Grimová <i>et al.</i> , 2016; Liang <i>et al.</i> , 2015	P1	P2	Ab3
Apple necrotic mosaic virus	ApNMV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Malus</i> spp.	Noda <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Apple rootstock virus A	ApRVA	<i>Rhabdoviridae</i>	<i>Nucleorhabdovirus</i>	<i>Malus</i> spp.	Morelli <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Apple rubbery wood virus 1	ARWV-1	<i>Phenuiviridae</i>	<i>Rubodvirus?</i>	<i>Malus</i> spp., <i>Pyrus</i> spp.	Rott <i>et al.</i> , 2018	P1	P2	Ab1
Apple rubbery wood virus 2	ARWV-2	<i>Phenuiviridae</i>	<i>Rubodvirus?</i>	<i>Malus</i> spp., <i>Pyrus</i> spp.	Rott <i>et al.</i> , 2018	P1	P2	Ab1
Apple scar skin viroid	ASSVd	<i>Pospiviroidae</i>	<i>Apscaviroid</i>	<i>Malus</i> spp., <i>Pyrus</i> spp.	Hadidi <i>et al.</i> , 2017	P2	P2	Ab1
Apple stem grooving	ASGV	<i>Betaflexiviridae</i>	<i>Capillovirus</i>	<i>Malus</i> spp.,	Hadidi <i>et al.</i> , 2011	P1	P2	Ab1

PEST	ABBREVIATION	FAMILY	GENUS	MAIN HOST(S)	REFERENCES	PRESENCE / ABSENCE		
						CAN	USA	MEX
virus				<i>Pyrus</i> spp.				
Apple stem pitting virus	ASPV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>Malus</i> spp., <i>Pyrus</i> spp.	Hadidi <i>et al.</i> , 2017	P1	P2	Ab1
Apricot latent virus	ApLV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	Ab1	Ab1	Ab1
Cherry leaf roll virus	CLRV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>Malus</i> spp.	Woo <i>et al.</i> , 2012	P7	P2	Ab1
Cherry rasp leaf virus	CRLV	<i>Secoviridae</i>	<i>Cheravirus</i>	<i>Malus</i> spp.	Noda <i>et al.</i> , 2017	P2	P2	Ab1
Citrus concave gum-associated virus	CCGaV	<i>Bunyaviridae?</i>	<i>Bunya-like virus</i>	<i>Malus</i> spp.	Wright <i>et al.</i> , 2018	Ab1	P2	Ab1
Citrus virus A	CiVA	<i>Bunyaviridae?</i>	<i>Bunya-like virus</i>	<i>Pyrus</i> spp.	Baek <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Hop stunt viroid	HSVd	<i>Pospiviroidae</i>	<i>Hostuviroid</i>	<i>Pyrus</i> spp.	Hadidi <i>et al.</i> , 2017	P7	P2	P2
<i>Malus domestica</i> virus A	MdoVA	<i>Closteroviridae</i>	<i>Velarivirus</i>	<i>Malus</i> spp.	Rott <i>et al.</i> , 2018	Ab1	Ab1	Ab1
Peach latent mosaic viroid	PLMVd	<i>Avsunviroidae</i>	<i>Pelamoviroid</i>	<i>Pyrus</i> spp.	Hadidi <i>et al.</i> , 2017	P7	P2	P2
Pear blister canker viroid	PBCVd	<i>Pospiviroidae</i>	<i>Apscaviroid</i>	<i>Pyrus</i> spp.	Rott <i>et al.</i> , 2018	P2	P2	Ab1
<i>Prunus</i> necrotic ringspot virus	PNRSV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2017	P7	P2	Ab3
<i>Pyrus pyrifolia</i> cryptic virus	PpCV	<i>Partitiviridae</i>	<i>Deltapartivirus</i>	<i>P. pyrifolia</i>	Osaki <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Raspberry bushy dwarf virus	RBDV	<i>Unassigned, proposed Bromoviridae</i>	<i>Idaeovirus</i>	<i>Cydonia</i> spp.	Hadidi <i>et al.</i> , 2011	P7	P2	Ab1
Tobacco ringspot virus	TRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	P2	P2	P2
Tomato ringspot virus	ToRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011	P2	P2	Ab3

REFERENCES

- Baek, D., Lim, S., Ju, H. J., Kim, H. R., Lee, S. H. & Moon, J. S.** 2019. The complete genome sequence of Apple Rootstock Virus A, a novel Nucleorhabdovirus identified in apple rootstocks. *Archives of Virology* 164: 2641-2644.
- Grimová, L., Winkowska, L., Konrady, M. & P. Rysánek, P.** 2016. *Apple mosaic virus*. *Phytopathologia Mediterranea* 55(1):1-19
- Hadidi, A., Barba, M., Candresse, T. & Jelkmann, W.** 2011. *Virus and Virus-Like diseases of Pome and Stone Fruits*, American Phytopathological Society, St. Paul, MN, USA. 429p.
- Hadidi, A., Flores, R., Randles, J. W. & Palukaitis, P.** 2017. *Viroids and Satellites*, Elsevier Science.
- Koloniuk, I., Přibylková, J., Fránová, J. & Špak, J.** 2020. Genomic characterization of *Malus domestica* Virus A (Mdova), a novel Velarivirus infecting apple. *Archives of Virology* 165(2): 479-482.
- Liang, P., Navarro, B., Zhang, Z., Wang, H., Lu, M., Xiao, H., Wu, Q., Zhou, X., Di Serio, F. & Li, S.** 2015. Identification and characterization of a novel *Geminivirus* with a monopartite genome infecting apple trees. *Journal of General Virology* 96(8), 2411-2420.
- Lim, S., Baek, D., Cho, I.-S., Do, Y. S., Lee, D. H. & Lee S. H.** 2019. First report of *Apple luteovirus 1* and *Apple rubbery wood virus 1* on apple tree rootstocks in Korea. *Plant Disease* 103(3): 591.
- Morelli, M., Giampetruzzi, A., Laghezza, L., Catalano, L., Savino, V. N. & Saldarelli, P.** 2017. Identification and characterization of an isolate of Apple Green Crinkle Associated Virus involved in a severe disease of quince (*Cydonia oblonga*, Mill.). *Archives of Virology* 162: 299-306.
- Noda, H., Yamagishi, N., Yaegashi, H., Xing, F., Xie, J., Li, S., Zhou, T., Ito, T. & Yoshikawa, N.** 2017. Apple Necrotic Mosaic Virus, a novel *Ilarvirus* from mosaic-diseased apple trees in Japan and China. *Journal of General Plant Pathology* 83: 83-90.
- Osaki, H., Sasaki, A., Nakazono-Nagaoka, E., Ota, N. & Nakaune, R.** 2017. Genome segments encoding capsid protein-like variants of *Pyrus pyrifolia* Cryptic Virus. *Virus Research* 240: 64-68.
- Rott, M. E., Kesanakurti, P., Berwarth, C., Rast, H., Boyes, I., Phelan, J. & Jelkmann, W.** 2018. Discovery of negative-sense RNA viruses in trees infected with apple rubbery wood disease by next-generation sequencing. *Plant Disease* 102: 1254-1263.
- Shen, P., Tian, X., Zhang, S., Ren, F., Li, P., Yu, Y. Q., Li, R., Zhou, C. & Cao, M.** 2018. Molecular characterization of a novel Luteovirus infecting apple by next-generation sequencing. *Archives of Virology* 163: 761-765.
- Szostek, S. A., Wright, A. A. & Harper S. J.** 2018. First report of *Apple hammerhead viroid* in the United States, Japan, Italy, Spain, and New Zealand. *Plant Disease* 102(12):2670.
- Woo, E. N. Y., Clover, G. R. G. & Pearson, M. N.** 2012. First report of Cherry leaf roll virus (CLRV) In *Malus domestica*. *Australasian Plant Disease Notes* 7: 151-156.
- Wright, A. A., Szostek, S. A., Beaver-Kanuya, E. & Harper, S. J.** 2018. Diversity of three Bunya-Like viruses infecting apple. *Archives of Virology* 163: 3339-3343.

Table 3: Fungal pathogens (incl. Chromista) of stone and pome fruit trees

Last updated December 2021.

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Alternaria alternata</i> (Fr.: Fr.) Keissl. Syn.: <i>Alternaria mali</i> Roberts	<i>Malus</i> spp.	Alternaria blotch	Jones and Aldwinckle, 1990	P2	P2	Ab1
<i>Alternaria gaisen</i> Nagano ex Hará. Syn.: <i>Alternaria kikuchiana</i> Tanaka	<i>Pyrus</i> spp.	Japanese pear black spot	Dickens and Cook, 1995; Sutton <i>et al.</i> , 2014	Ab1	Ab3	Ab1
<i>Apiognomonium erythrostoma</i> (Fuckel) Höhn (Pers.) V. Hohnel.	<i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus cerasus</i>	Red spot, leaf scorch, gnomoniosis	Chen <i>et al.</i> , 2002; Hecht and Zinkernagel, 2006; Ogawa <i>et al.</i> , 1995	Ab1	Ab3	Ab1
<i>Athelia rolfsii</i> (Curzi) C.C. Tu and Kimbr. Syn.: <i>Sclerotium rolfsii</i> Sacc	<i>Prunus</i> spp., <i>Malus</i> spp.	Sclerotium Stem Rot, Southern blight	Ogawa <i>et al.</i> , 1995; Ohlendorf, 1999; Sutton <i>et al.</i> , 2014	P7	P2	P2
<i>Berkeleyomyces basicola</i> (Berk. and Broome) W.J. Nel, Z.W. de Beer, T.A. Duong and M.J. Wingf. Syn.: <i>Thielaviopsis basicola</i> (Berk. and Broome) Ferraris, <i>Chalara elegans</i> Nag Raj and W.B. Kendr	<i>Prunus avium</i>	Black root rot	Sewell and Wilson, 1975; Sønderhausen, 1970	P1	P2	Ab3
<i>Blumeriella jaapii</i> (Rehm) Arx	<i>Prunus</i> spp.	Leaf spot, Shot hole	Joshua and Mmbaga, 2015; Ogawa <i>et al.</i> , 1995; USDA, 2020	P1	P2	Ab1
<i>Botryosphaeria dothidea</i> (Moug.:Fr.) Ces. and De Not. (Moug. ex Fr.) Ces. and De Not. Syn.: <i>Fusicoccum aesculi</i> Corda	<i>Prunus</i> spp., <i>Pyrus</i> spp., <i>Malus</i> spp.	Fungal gummosis (Prunus), white rot (Malus)	Ko <i>et al.</i> , 2010; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	P2	P2	Ab3
<i>Botryosphaeria kuwatsukai</i> (Hara) G.Y. Sun and E. Tanaka. Syn.: <i>Guignardia piricola</i> (nose) W. Yanam, <i>Macrophoma kuwatsukai</i> Hara, <i>Botryosphaeria berengeriana</i> f. sp. <i>pyricola</i> (Nose) Kogan. and	<i>Prunus americana</i> , <i>Pyrus pyrifolia</i> , <i>Malus domestica</i>	Apple ring rot, Plum wilt	Deng <i>et al.</i> , 2004; Sutton <i>et al.</i> , 2014; Xu <i>et al.</i> , 2015	Ab1	Ab1	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
Sakuma, <i>Guignardia pyricola</i> (Nose) W. Yamam						
<i>Botrytis cinerea</i> Pers.: Fr. Syn.: <i>Botryotinia fuckeliana</i> (de Bary) Whetzel	<i>Chaenomeles</i> spp., <i>Prunus</i> spp., <i>Pyrus</i> spp. and <i>Malus</i> spp.	Gray mold, dry eye rot blossom end rot green fruit rot	Norina and Rumpunen, 2003; Ferrada <i>et al.</i> , 2016; Strand, 1999; Sutton <i>et al.</i> , 2014	P2	P2	P2
<i>Cadophora malorum</i> (Kidd and Beaumont) W. Gams Syn.: <i>Phialophora malorum</i> (Kidd and Beaumont) McColloch	<i>Malus</i> spp., <i>Pyrus</i> spp.	Side rot	McColloch, 1944; Sugar and Spotts, 1992; Sutton <i>et al.</i> , 2014	P2	P2	Ab3
<i>Ceratobasidium ochroleucum</i> (F. Noack) Ginns and M.N. Lefebvre Syn.: <i>Corticium stevensii</i> Burt	<i>Malus</i> spp., <i>Pyrus</i> spp.	Thread blight	Jones and Sutton, 2019; Sutton <i>et al.</i> , 2014	Ab1	P2	Ab3
<i>Colletotrichum gloeosporioides</i> (Penz.) Penz. and Sacc. Syn.: <i>Glomerella cingulata</i> (Stoneman) Spauld. and H. Schrenk	<i>Prunus dulcis</i> , <i>Prunus</i> spp., <i>Malus</i> spp., <i>Pyrus</i> spp.	Anthracnose, Fruit rot, Bitter rot, Black spot	Haviland <i>et al.</i> , 2019; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	P1	P2	P2
<i>Cylindrocarpon didymum</i> (Harting) Wollenweb.	<i>Malus x domestica</i>	Twig blight, Black foot	Dugan and Grove, 1994; Littke and Brwning 2001	P7	P2	Ab3
<i>Cytospora ceratosperma</i> (Tode) G.C. Adams and Rossman. Syn.: <i>Valsa ceratosperma</i> (Tode) Maire, <i>Cytospora sacculus</i> (Schwein.) Gvritschvili	<i>Cydonia oblonga</i> , <i>Malus</i> spp., <i>Pyrus</i> spp.	Valsa canker	CABI 2021; Sutton <i>et al.</i> , 2014	P7	P2	Ab1
<i>Cytospora cincta</i> Sacc. Syn.: <i>Leucostoma cinctum</i> (Fr.: Fr.) Höhn., <i>Leucostoma cincta</i> (Fr.: Fr.) Höhn.; <i>Valsa cincta</i> (Fr.: Fr.) Fr.	<i>Prunus avium</i> , <i>Prunus</i> spp., <i>Malus</i> spp.	Dieback, Perennial canker	Barakat and Johnson, 1997; Brown-Rytlewski and McManus, 2000; Jones, 1989; Ogawa <i>et al.</i> , 1995; Proffer and Jones, 1989; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Cytospora leucostoma</i> (Pers.) Sacc. Syn.: <i>Leucostoma persoonii</i> (Nitschke) Höhn., <i>Valsa leucostoma</i> (Pers.: Fr.) Fr.	<i>Cydonia</i> , <i>Prunus domestica</i> , <i>Prunus</i> spp.	Dieback, canker	Biggs and Grove, 2005; Ogawa <i>et al.</i> , 1995; Strand, 1999	P1	P2	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Diaporthe ambigua</i> Nitschke Syn.: <i>Phomopsis ambigua</i> (Sacc.) Traverso	<i>Malus domestica</i> , <i>Prunus salicina</i> , <i>Pyrus communis</i> , <i>Malus</i> spp.	Constriction canker, Phomopsis fruit rot, Branches dieback	Havenga <i>et al.</i> , 2019; Smit, 1996	P2	P2	Ab1
<i>Diaporthe amygdali</i> (Delacr.) Udayanga, Crous and K.D. Hyde Syn.: <i>Phomopsis amygdali</i> (Del.) Tuset and Portilla, <i>Fusicoccum amygdali</i> Del.	<i>Prunus dulcis</i> , <i>Prunus persica</i> , <i>Malus</i> spp, <i>Pyrus pyrifolia</i>	Constriction canker, Phomopsis fruit rot, Branches dieback	Bai <i>et al.</i> , 2015; Havenga <i>et al.</i> , 2019; Haviland <i>et al.</i> , 2019; Ogawa <i>et al.</i> , 1995	P2	P2	Ab1
<i>Diaporthe eres</i> Nitschke Syn.: <i>Phomopsis fukushii</i> Tanaka and S. Endô, <i>Phomopsis oblonga</i> (Desm.) Traverso	<i>Cydonia oblonga</i> , <i>Malus</i> spp., <i>Prunus avium</i> , <i>P. persica</i> , <i>Pyrus pyrifolia</i>	Bark canker	Abramczyk <i>et al.</i> , 2018; Ali <i>et al.</i> , 2019; Bai <i>et al.</i> , 2015; Prencipe <i>et al.</i> , 2017; Ristić <i>et al.</i> , 2016	P2	P2	Ab1
<i>Diaporthe perniciosa</i> Marchal and É.J. Marchal. Syn.: <i>Phomopsis mali</i> Roberts	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus</i> spp.	Canker and fruit decay	Ogawa, 1995; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Diaporthe tanakae</i> Ts. Kobay. and Sakuma Kobayashi and Sakuma Syn.: <i>Phomopsis tanakae</i> Kobayashi and Sakuma	<i>Malus</i> spp., <i>Pyrus</i> spp.	Canker	Sutton <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Didymosphaeria rubi-ulmifoli</i> Ariyaw., Camporesi and K.D. Hyde	<i>Prunus pérsica</i> , <i>Malus</i> spp.		Havenga <i>et al.</i> , 2019; Liang <i>et al.</i> , 2020	Ab1	Ab1	Ab1
<i>Diplodia mutila</i> (Fr.: Fr.) Mont. Syn.: <i>Botryosphaeria stevensii</i> Shoemaker, <i>Physalospora malorum</i> Shear, N. Stevens, and M.S. Wilcox	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus salicina</i> , <i>Prunus laurocerasus</i>	Black rot, Dieback, Diplodia Canker.	Damm <i>et al.</i> , 2007; Jones and Aldwinckle, 1990; Sutton <i>et al.</i> , 2014; Úrbez-Torres <i>et al.</i> , 2016; Zlatković <i>et al.</i> , 2018	P1	P2	Ab1
<i>Diplodia seriata</i> De Not. Syn.: <i>Botryosphaeria obtusa</i> (Schwein.) Shoemaker, <i>Sphaeropsis malorum</i> Berk.	<i>Cydonia oblonga</i> , <i>Prunus dulcis</i> , <i>Malus</i> spp., <i>Pyrus pyrifolia</i>	Dieback, Black rot, Fungal gummosis	Haviland <i>et al.</i> , 2019; Mohammadi and Sharifi, 2016; Sutton <i>et al.</i> , 2014	P1	P2	P2

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Ellisembia asterinum</i> (Cooke) Shoemaker and Hambl. Syn.: <i>Helminthosporium papulosum</i> A. Berg	<i>Malus</i> spp., <i>Pyrus</i> spp.	Black pox of apple, Blister canker of pear	Sutton <i>et al.</i> , 2014	Ab1	P2	Ab1
<i>Erythricium salmonicolor</i> (Berk. and Broome) Burdsall	<i>Malus</i> spp., <i>Pyrus</i> spp.	Pink disease	Momol <i>et al.</i> , 2017; Prasad, 2013	Ab1	P2	Ab3
<i>Erythricium salmonicolor</i> (Berk. and Broome) Burds. Syn.: <i>Corticium salmonicolor</i> Berk. and Broome	<i>Malus</i> spp., <i>Pyrus</i> spp.	Pink disease, Limb blight	Momol <i>et al.</i> , 2007; Prasad, 2013	Ab1	P2	Ab3
<i>Eutypa lata</i> (Pers.) Tul. and C. Tul.	<i>Cydonia oblonga</i> , <i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus domestica</i>	Eutypa dieback, Gummosis disease	Moyo <i>et al.</i> , 2017; Ogawa <i>et al.</i> , 1995; Strand, 1999	P2	P2	Ab3
<i>Gloeodes pomigena</i> (Schwein.) Colby. Syn.: <i>Phyllachora pomigena</i> (Schwein.) Sacc.	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus</i> spp.	Sooty blotch, flyspeck	Estafne, 2015; Jones and Aldwinckle, 1990; USDA, 2020; Wilcox, 1994; Ogawa <i>et al.</i> , 1995	P2	P2	Ab1
<i>Gloeocystidiellum sacratum</i> (G. Cunn.) Stalpers and P.K. Buchanan. Syn.: <i>Peniophora sacrata</i> G. Cunn.	<i>Malus sylvestris</i> , <i>Malus</i> spp.	Peniophora root canker	Sutton <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Gloeosporium laeticolor</i> Berk.	<i>Prunus persica</i>	Anthraco nose	Ogawa <i>et al.</i> , 1995	Ab1	Ab1	Ab1
<i>Gymnosporangium asiaticum</i> Miyabe ex G. Yamada. Syn.: <i>Roestelia koreaënsis</i> Henn.	<i>Pyrus</i> spp., <i>Chaenomeles</i> spp., <i>Cydonia oblonga</i>	Japanese pear rust	Kikuhara <i>et al.</i> , 2019; Sutton <i>et al.</i> , 2014; Yun <i>et al.</i> , 2009	Ab1	P2	Ab1
<i>Gymnosporangium clavipes</i> Cooke and Peck (Cooke and Peck) Cooke and Peck en Peck	<i>Malus</i> spp., <i>Chaenomeles</i> spp., <i>Cydonia</i> spp. <i>Pyrus calleryana</i> ;	Quince rust	Bergdahl <i>et al.</i> , 2016; Creswell <i>et al.</i> , 2015; Ellis, 2016; Gauthier and Heisdorffer, 2015; MBG, 2020	P1	P2	P2
<i>Gymnosporangium globosum</i> (Farl.) Farl.	<i>Malus</i> spp., <i>Pyrus</i> spp.	American hawthorn rust	Jones and Aldwinckle, 1990	P1	P1	P2

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Gymnosporangium juniperi-virginianae</i> Schwein.	<i>Malus</i> spp.	Cedar-apple rust	Bergdahl <i>et al.</i> , 2016; Gauthier and Heisdorffer, 2015; Morton Arboretum, 2020	P2	P2	Ab1
<i>Gymnosporangium kernianum</i> Bethel	<i>Pyrus</i> spp.	Kern's pear rust	Jones and Aldwinckle, 1990	Ab1	P2	Ab3
<i>Gymnosporangium libocedri</i> (Henn.) F. Kern	<i>Pyrus</i> spp., <i>Cydonia</i> , <i>Chaenomeles</i>	Pacific coast pear rust	Windbiel-Rojas <i>et al.</i> , 2020; Worrall, 2020	Ab1	P2	Ab1
<i>Gymnosporangium nelsonii</i> Arthur.	<i>Pyrus</i> spp.	Rocky mountain pear rust	Jones and Aldwinckle, 1990	P2	P2	Ab3
<i>Gymnosporangium sabinae</i> (Dicks.) G. Winter Syn.: <i>Gymnosporangium fuscum</i> R. Hedw. in DC	<i>Pyrus calleryana</i> <i>Pyrus</i> spp.	European pear rust, Pear trellis Rust	Hansen <i>et al.</i> , 2016; Lim <i>et al.</i> , 1978; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Gymnosporangium yamadae</i> Miyabe ex G. Yamada	<i>Malus</i> spp.	Japanese apple rust	Gregory <i>et al.</i> , 2010, Sutton <i>et al.</i> , 2014	Ab1	P2	Ab1
<i>Helicobasidium mompa</i> Tanaka Syn.: <i>Septobasidium mompa</i> (Tanaka) Racib.	<i>Prunus</i> spp., <i>Malus</i> spp.	Violet root rot	Nakamura <i>et al.</i> , 2004; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Monilinia fructigena</i> Honey in Whetzel (Aderh. and Ruhland) Honey	<i>Prunus</i> spp., <i>Malus</i> spp., <i>Pyrus</i> spp.	Brown rot, Blossom and twig blight	Lesik, 2013; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	Ab1	Ab3	Ab1
<i>Monilinia kusanoi</i> (Henn. ex Takah.) (Takah.) W. Yamamoto. Syn.: <i>Monilia kusanoi</i> P. Henn.	<i>Prunus</i> spp., <i>Prunus avium</i> , <i>Prunus grayana</i>	Leaf blight, green fruit rot	Ogawa <i>et al.</i> , 1995	Ab1	Ab1	Ab1
<i>Monilinia laxa</i> (Aderhold and Ruhland) Honey. Syn.: <i>Sclerotinia laxa</i> Aderh and Ruhland	<i>Prunus</i> spp., <i>Malus</i> spp. <i>Chaenomeles</i> spp., <i>Pyrus</i> spp.	Brown rot	Haviland <i>et al.</i> , 2019; Lesik 2013; Norina and Rumpunen, 2003; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Monilinia linhartiana</i> (Prill. and Delacr.) N.F. Buchw. Syn.: <i>Monilia cydoniae</i> Schellenb.	<i>Cydonia oblonga</i>	Leaf blotch, Shoot blight	Lantos <i>et al.</i> , 2017	Ab1	Ab1	Ab1
<i>Monilinia mali</i> (Takah.) Whetzel. Syn.: <i>Sclerotinia mali</i> Takah.	<i>Malus</i> spp.	Blossom blight, leaf blight	Sutton <i>et al.</i> , 2014; Webster <i>et al.</i> , 2020	Ab1	Ab1	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Monilia polystroma</i> G. van Leeuwen et al. Syn.: <i>Monilinia polystroma</i> (G.C.M. Leeuwen) Kohn	<i>Malus</i> spp., <i>Pyrus</i> spp.	Brown rot	Petróczy and Palkovic, 2009; Van Leeuwen et al., 2002	Ab1	Ab1	Ab1
<i>Neofabraea malicorticis</i> (Cordley) H. Jacks. Syn.: <i>Cryptosporiopsis curvispora</i> (Peck) Gremmen, <i>Pezicula malicorticis</i> (Cordley) Nannf.	<i>Malus</i> spp.	Anthracnose canker and Perennial canker	Brun and Bush, 2016; Sutton et al., 2014	P2	P2	Ab3
<i>Neofusicoccum ribis</i> (Slippers, Crous and M.J. Wingf.) Crous, Slippers and A.J.L. Phillips Syn.: <i>Botryosphaeria ribis</i> Grossenb. and Duggar, <i>Dothiorella ribis</i> (Fuckel) Sacc	<i>Malus</i> spp., <i>Pyrus</i> spp.	Fruit rot, gummosis	Brown II and Briton, 1986; Elliot, 2015; Vovey Jr. 1967; Jurick et al., 2013; Urbez-Torres et al., 2012; WPFUS, 2020	P7	P1	Ab3
<i>Neonectria ditissima</i> (Tul. and C. Tul.) Samuels and Rossman Syn.: <i>Nectria galligena</i> Bres., <i>Neonectria galligena</i> (Bres.) Rossman and Samuels, <i>Cylindrocarpon heteronema</i> (Berk. and Broome) Wollenw.	<i>Malus</i> spp., <i>Pyrus</i> spp.	European canker, Thread blight	Ohlendorf, 1999; Sutton et al., 2014; Wunderlich et al., 2017	P2	P2	Ab3
<i>Phacidiopycnis pyri</i> (Fuckel) Weindlm J. Weindlymayr. Syn.: <i>Discula pyri</i> (Fuckel) Höhn, <i>Potebniomyces pyri</i> (Berk. and Broome) Dennis, <i>Potebniomyces discolor</i> (Mouton and Sacc.) Smerlis	<i>Pyrus</i> spp., <i>Malus</i> spp.	Canker, stem end rot	Amiri, 2020; Xiao and Boal, 2005	P2	P2	Ab1
<i>Phacidiopycnis washingtonensis</i> C. L. Xiao and J. D. Rogers	<i>Pyrus</i> spp., <i>Malus</i> spp.	Speck rot	Amiri, 2020; Xiao et al. 2005	Ab1	P2	Ab1
<i>Phaeoacremonium parasiticum</i> (Ajello, Georg and C.J.K. Wang) W. Gams, Crous and M.J. Wingf. Syn.: <i>Phialophora parasitica</i> Ajello, Georg and Wang	<i>Cydonia oblonga</i> , <i>Prunus avium</i> , <i>Prunus armeniaca</i> , <i>Prunus salicina</i> , <i>Malus</i> spp., <i>Pyrus</i> spp.	Dieback	Damm et al., 2008; Groenewald et al., 2001; Mohammadi and Sharifi, 2016; Ogawa et al., 1995; Rumbos, 1986; Sami et al., 2014	Ab1	P2	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Phaeoacremonium scolyti</i> L. Mostert, Summerb. and Crous	<i>Cydonia oblonga</i> , <i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus</i> spp.	Esca and Petri disease	Damm <i>et al.</i> , 2008, Havenga <i>et al.</i> , 2019; Sami <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Phaeoacremonium parasiticum</i> (Ajello, Georg and C.J.K. Wang) W. Gams, Crous and M.J. Wingf. Syn.: <i>Phialophora parasitica</i> Ajello, Georg and Wang	<i>Cydonia oblonga</i> ; <i>Prunus avium</i> , <i>Prunus armeniaca</i> , <i>Prunus salicina</i> ; <i>Malus</i> spp., <i>Pyrus</i> spp.	Dieback	Damm <i>et al.</i> , 2008; Mohammadi and Sharifi, 2016; Ogawa <i>et al.</i> , 1995; Rumbos 1986; Sami <i>et al.</i> , 2014	Ab1	P2	Ab1
<i>Phyllosticta arbutifolia</i> Ellis and G. Martin Syn.: <i>Phyllosticta solitaria</i> Ellis and Everh.	<i>Malus</i> spp.	Blotch	Jones and Aldwinckle, 1990; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Phymatotrichopsis omnivora</i> (Duggar) Hennebert	<i>Malus</i> spp., <i>Prunus</i> spp., <i>Pyrus</i> spp., <i>Cydonia</i> spp.	Texas root rot	Jones and Aldwinckle, 1990; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	Ab1	P2	Ab3
<i>Phytophthora cambivora</i> (Petri) Buisman	<i>Malus</i> spp., <i>Prunus</i> spp.	Ink disease, Collar rot, Root rot, Trunk canker	Erwin and Ribeiro, 1996; Ogawa <i>et al.</i> , 1995; Rooney-Latham <i>et al.</i> , 2018; Sutton <i>et al.</i> , 2014	P2	P2	Ab1
<i>Phytophthora cryptogea</i> Pethybr. and Lafferty	<i>Prunus</i> spp., <i>Malus</i> spp.	Collar rot	Erwin and Ribeiro, 1996; Jones and Aldwinckle, 1990; Latorre <i>et al.</i> , 2001; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	P5	P2	P2
<i>Phytophthora drechsleri</i> Tucker	<i>Malus</i> spp., <i>Prunus</i> spp.	Root rot, Trunk canker	Erwin and Ribeiro, 1996; Sutton <i>et al.</i> , 2014	P7	P2	P2
<i>Phytophthora megasperma</i> Dreschsler	<i>Malus</i> spp., <i>Prunus</i> spp.	Collar rot, crown rot	Erwin and Ribeiro, 1996, Jones and Aldwinckle, 1990; Kurbetli and Yilmaz, 2015; Ogawa <i>et al.</i> , 1995	P7	P2	Ab1
<i>Phytophthora syringae</i> (Kleb.) Kleb.	<i>Malus</i> spp., <i>Prunus</i> spp., <i>Pyrus</i> spp.	Fruit rot	Grigel <i>et al.</i> , 2019; Jones and Aldwinckle, 1990; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014	P2	P2	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCE	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Phytophthium vexans</i> (de Bary) Abad, de Cock, Bala, Robideau, Lodhi and Lévesque Syn.: <i>Pythium vexans</i> de Bary	<i>Prunus persica</i>	Damping-off, root rot	Biesbrock and Hendrix Jr., 1970; Hendrix <i>et al.</i> , 1966	P7	P2	Ab1
<i>Podosphaera pannosa</i> (Wallr.: Fr.) de Bary. Syn.: <i>Sphaerotheca pannosa</i> (Wallr.: Fr.) Lév	<i>Prunus</i> spp., <i>Pyrus</i> spp.	Powdery mildew	Ogawa <i>et al.</i> , 1995; Pscheidt and Ocamb, 2020; Strand, 1999	P1	P2	P2
<i>Pythium irregulare</i> Buisman	<i>Malus</i> spp.	Replant disease	Braun, 1991	P1	P1	P2
<i>Rosellinia necatrix</i> Berl. ex Prill. Syn.: <i>Dematophora necatrix</i> R. Hartig	<i>Cydonia oblonga</i> , <i>Pyrus</i> spp., <i>Prunus</i> spp., <i>Malus</i> spp.	Root rot	Ogawa <i>et al.</i> , 1995; Pastorelli, 2012; Strand, 1999; Sutton <i>et al.</i> , 2014; Thomas <i>et al.</i> , 1953; Van den Berg <i>et al.</i> , 2018	Ab1	P2	P2
<i>Tranzschelia discolor</i> (Fuckel) Tranzschel and M.A. Litv. Syn.: <i>Tranzschelia pruni-spinosae</i> var. <i>discolor</i> (Pers. :Pers.) Dietel var. <i>discolor</i> (Fuckel) Dunegan	<i>Prunus</i> spp.	Rust	Haviland <i>et al.</i> , 2019; Ogawa <i>et al.</i> , 1995; Soto-Estrada <i>et al.</i> , 2005; Strand, 1999	P2	P2	Ab3
<i>Venturia carpophila</i> E.E. Fisher Syn.: <i>Fusicladium amygdali</i> Ducomet, <i>Cladosporium carpophilum</i> Thüm.	<i>Prunus</i> spp.	Peach scab	Haviland <i>et al.</i> , 2019; Ogawa <i>et al.</i> , 1995	P1	P2	Ab1
<i>Venturia nashicola</i> S. Tanaka and S. Yamamoto Syn.: <i>Fusicladium nashicola</i> K. Schub. and U. Braun	<i>Pyrus pyrifolia</i> , <i>Pyrus ussuriensis</i>	Asian Pear scab	Li <i>et al.</i> , 2003; Sutton <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Verticillium albo-atrum</i> Reinke and Berthier Berthold	<i>Prunus</i> spp.	Verticillium wilt	Ogawa <i>et al.</i> , 1995	P1	P2	P2
<i>Verticillium dahliae</i> Kleb.	<i>Prunus</i> spp.	Verticillium wilt	Pscheidt and Ocamb, 2020; Ogawa <i>et al.</i> , 1995	P1	P2	P2

References

- Ali, S., Renderos, W., Bevis, E., Hebb, J. & Abbasi, P. A.** 2019. *Diaporthe eres* causes stem cankers and death of young apple rootstocks in Canada. Canadian Journal of Plant Pathology:1-10.
- Amiri, A.** 2020. Phacidiopycnis Rot. Washington State University.
- Abramczyk, B. A., Król, E. D., Zalewska, E. D. & Zimowska B.** 2018. Morphological characteristics and pathogenicity of *Diaporthe eres* isolates to the fruit tree shoots. Acta Scientiarum Polonorum Hortorum Cultus 17(6):125-133.
- Barakat, R. M. & Johnson, D. A.** 1997. Expansion of cankers caused by *Leucostoma cincta* on sweet cherry trees. Plant Disease 81(12):1391-1394.
- Bai, Q., Zhai, L., Chen, X., Hong, N., Xu, W. & Wang G.** 2015. Biological and molecular characterization of five *Phomopsis* species associated with pear shoot canker in China. Plant Disease 99(12):1704-1712.
- Bergdahl, A. D., A. Hill,** and tech. coords. 2016. Diseases of trees in the Great Plains (General Technical Report RMRS-GTR-335). U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO. 229 pp.
- Biesbrock, J. A. & Hendrix Jr., F. F.** 1970. Influence of soil water and temperature on root necrosis of peach caused by *Pythium spp.* Phytopathology 60:880-882.
- Biggs, A. R. & Grove, G. G.** 2005. *Leucostoma* canker of stone fruits. American Phytopathological Society, St. Paul, MN, USA.
- Braun, P. G.** 1991. The combination of *Cylindrocarpon lucidum* and *Pythium irregulare* as a possible cause of ample replant disease in Nova Scotia. Can. J. Plant Pathol. 13:291-297.
- Brown II, E. A., & K. O. Britton.** 1986. *Botryosphaeria* diseases of apple and peach in the Southeastern United States. Plant Disease 70(5):480-484.
- Brown-Rytlewski, D. E. & McManus, P. S.** 2000. Outbreak of leucostoma canker caused by *Leucostoma cincta* on McIntosh apple trees in Wisconsin. Plant Disease 84(8):923-923.
- Brun, C. & Bush, M.** 2016. Pest management guide for apples in Washington home orchards. (EM101E). Washington State University Extension 21 pp.
- CABI.** 2021. Crop Protection Compendium. Centre for Agriculture and Bioscience International (CABI). <https://www.cabi.org/cpc/>. (Archived at PERAL).
- Chen, W. Q., Ntahimpera, N., Morgan, D. P. & Michailides T. J.** 2002. Mycoflora of *Pistacia vera* in the Central Valley, California. Mycotaxon 83:147-158.
- Damm, U., Crous, P. W. & Fourie P. H.** 2007. Botryosphaeriaceae as potential pathogens of *Prunus* species in South Africa, with descriptions of *Diplodia africana* and *Lasiodiplodia plurivora* sp. nov. Mycologia 99(5):664-680.
- Damm, U., Mostert, L., Crous, P. W. & Fourie, P. H.** 2008. Novel *Phaeoacremonium* species associated with necrotic wood of *Prunus* trees. Persoonia 20:87-102.
- Deng, X., Guo, L., Lin, Z., & Zhang, T.** 2004. Identification of the pathogen of *Prunus americana* and its medicinal sensitivity (Chinese). Plant Protection 30(5):29-32.
- Dickens, J. S. W., & R. T. A. Cook.** 1995. Japanese pear black spot and apple Alternaria blotch. Bulletin OEPP/EPPO Bulletin Bulletin 25.
- Dugan, F. M. & Grove, G. G.** 1994. *Cylindrocarpon didymium* pathogenic on apple seedlings in Washington. Plant Dis. 78:1219.
- Elliott, M.** 2015. Tree Disease identification stem and branch 1: cankers & *Phytophthora* diseases. WSU Puyallup Research and Extension Center, Washington State University.
- Estafne.** 2015. Reported fruit problems in Mississippi 2014. Mississippi State University Extension Service. Last accessed 4/30/2020, <https://blogs.extension.msstate.edu/fruit/2015/02/03/reported-fruit-problems-in-mississippi-2014/>.
- Erwin, D. C., & Ribeiro O. K.** 1996. *Phytophthora* Diseases Worldwide. American Phytopathological Society, St. Paul, MN, USA. 562 pp.

- Ferrada, E., B. Latorre, J. Zoffoli, & A. Castillo.** 2016. Identification and characterization of *Botrytis* blossom blight of Japanese plums caused by *Botrytis cinerea* and *B. prunorum* sp. nov. in Chile. *Phytopathology* 106(2):155-165.
- Gauthier, N. W., & A. M. Heisdorffer.** 2015. Apple rust diseases (PPFS-FR-T-05). University of Kentucky.
- Gregory, N. F., Bischoff, J. F., Dixon, L. J. & Ciurlino R.** 2010. First report of the telial stage of Japanese apple rust on *Juniperus chinensis* in North America and the aecial stage on *Malus domestica*. *Plant Disease* 94(9):1169-1169.
- Grígel, J., K. Černý, M. Mrázková, L. Havrdová, D. Zahradník, B. Jílková, & M. Hrabětová.** 2019. *Phytophthora* root and collar rots in fruit orchards in the Czech Republic. *Phytopathologia Mediterranea* 58(2):261-275.
- Groenewald, M., J.-C. Kang, P. W. Crous, & W. Gams.** 2001. ITS and β -tubulin phylogeny of *Phaeoacremonium* and *Phaeomoniella* species. *Mycological Research* 105(6):651-657.
- Hansen, M. A., Demers, J., Sutphin, M., Yoder, K., Bush, E., & L. Castlebury, L.** 2016. First report of European pear rust (pear trellis rust) caused by *Gymnosporangium sabiniae* on ornamental pear (*Pyrus calleryana*) in Virginia. *Plant Disease* 100(10):2166-2166.
- Havenga, M., Gatsi, G. M., Halleen, F., Spies, C. F. J., van der Merwe, R. & Mostert L.** 2019. Canker and wood rot pathogens present in young apple trees and propagation material in the Western Cape of South Africa. *Plant Disease* 103(12):3129-3141.
- Haviland, D. R., E. J. Symmes, J. E. Adaskaveg, R. A. Duncan, J. A. Roncoroni, W. D. Gubler, B. Hanson, K. J. Hembree, B. Holtz, J. J. Stapleton, K. E. Tollerup, F. P. Trouillas, & F. G. Zalom.** 2019. UC IPM Pest Management Guidelines: Almond (UC ANR Publication 3431). UC Statewide Integrated Pest Management Program, Oakland, CA.
- Hecht, D., & Zinkernagel, V.** 2006. Epidemiological studies of cherry leaf scorch (*Gnomonia erythrostoma* (Pers.) Auerswald). *Journal of Plant Diseases and Protection* 113:68-73.
- Hendrix, E. F. Jr., Powell, W. M. & Owen, J. H. H.** 1966. Relation of root necrosis caused by *Pythium* species to peach tree decline. *Phytopathology* 56:1229-1232.
- Jones, A. L. & Aldwinckle, H.S.** 1990. Compendium of Apple and Pear Diseases. American Phytopathological Society, St. Paul, MN, USA. 100 pp.
- Jones, A. L. & Sutton, T. B.** 2019. Thread blight of apple. Cooperative Extension, USDA.
- Joshua, J., Mmbaga, M. T. & Mackasmiel, L. A.** 2017. Cherry leaf spot disease management in ornamental cherries in mid-Tennessee. *Canadian Journal of Plant Science* 97:110-118.
- Jurick, W. M., I. Vico, V. L. Gaskins, W. J. Janisiewicz, & K. A. Peter.** 2013. First report of *Neofusicoccum ribis* causing postharvest decay of apple fruit from cold storage in Pennsylvania. *Plant Disease* 97(7):999-999.
- Kikuhara, K., Iiyama, K. Matsumoto, M. & Furuya, N.** 2019. First report on DMI fungicide resistance in *Gymnosporangium asiaticum*, the causal agent of Japanese pear rust, in Japan. *Journal of General Plant Pathology* 85(1):49-56.
- Ko, Y., C. W. Liu, S. S. Chen, K. Y. Chiu, Y. W. Sun, & S. Maruthasalam.** 2010. First report of gummosis disease of Japanese apricot caused by *Botryosphaeria dothidea* in Taiwan. *Plant Disease* 95(1):77-77.
- Kurbetli, İ., & A. Yilmaz.** 2015. First report of *Phytophthora megasperma* causing crown and root rot of almond in Turkey. *New Disease Reports* 32. <http://dx.doi.org/10.5197/j.2044-0588.2015.032.025>
- Lantos, A., Petróczy, M., Oláh, R. & Palkovics L.** 2017. Characterization of *Monilinia linhartiana* isolates. *Journal of Plant Pathology* 99(3):583-591.
- Lesik, K.** 2013. *Monilinia* species causing fruit brown rot, blossom and twig blight in apple orchards in Belarus. *Proceedings of the Latvian Academy of Sciences. Section B* 67:192-194.
- Liang, K., Huang, W., He, Q., Lan, J. Lu, & Q. & Pu L.** 2020. First report of *Didymosphaeria rubi-ulumifolii* causing wood rot of peach (*Prunus persica*) in China. *Plant Disease*.
- Li, B., Zhao, H. Li, B. & Xu X.-M.** 2003. Effects of temperature, relative humidity and duration of wetness period on germination and infection by conidia of the pear scab pathogen (*Venturia nashicola*). *Plant Pathology* 52(5):546-552.

- Lim, W. L., S. H. Wang, & O. C. Ng. 1978. Overwintering of pear trellis rust in pear. *Plant Disease Reporter* 62:659-660.
- Littke, W., & Browning J. 2001. *Cylindrocarpon didymum*-a new pathogen or secondary root colonizer? Western Nursery Pathology Workshop,
- McColloch, L. P. 1944. A study of the apple rot fungus *Phialophora malorum*. *Mycologia* 36(6):576-590.
- Mohammadi, H. & Sharifi, S. 2016. Association of Botryosphaeriaceae and Phaeoacremonium species with insect-damaged quince shoots. *Journal of Plant Pathology* 98(1):35-42.
- Momol, T., Ritchie, L., & Dankers, H. 2007. 2007 Florida plant disease management guide: pear (*Pyrus communis*) (PDMG-V3-27). University of Florida, IFAS Extension.
- Morton Arboretum. 2020. Cedar-apple rust. The Morton Arboretum. <https://www.mortonarb.org/trees-plants/tree-and-plant-advice/help-diseases/cedar-apple-rust>.
- Moyo, P., Halleen, F., Dedekind, R., van Jaarsveld, W. J., Pierron, R. & Mostert, L. 2017. First report of twig dieback and cankers caused by *Eutypa lata* on quince in South Africa. *Plant Disease* 102(7):1446.
- Nakamura, H., Ikeda, K.-i., Arakawa, M., Akahira, T. & Matsumoto N. 2004. A comparative study of the violet root rot fungi, *Helicobasidium brebissonii* and *H. mompa*, from Japan. *Mycological Research* 108(6):641-648.
- Norina, I. & Rumpunen K. 2003. Pathogens on Japanese quince (*Chaenomeles japonica*) plants. *Biology*:37-54.
- Ogawa, J. M., Zehr, E. I., Bird, G. W., Ritchie, D. F., Uriu, K., & Uyemoto, J. K. 1995. Compendium of Stone Fruit Diseases. American Phytopathological Society, St. Paul, MN, USA. 98 pp.
- Ohlendorf, L. P. 1999. Integrated Pest Management for Apples & Pears (Second Edition). University of California, Agriculture and Natural Resources, Oakland, CA. 231 pp.
- Pastorelli, S. 2012. *Rosellinia necatrix*: an increasing problem in apple orchards Fondazione Edmund Mach.
- Petróczy, M., & L. Palkovic. 2009. First report of *Monilia polystroma* on apple in Hungary. *European Journal of Plant Pathology* 125:343-347.
- Prasad, D. 2013. Studies on epidemiology and management of pink canker (*Corticium salmonicolor* berk. & br.) in apple, Dr. Y. S. Parmar University of Horticulture and Forestry, Nauni, Solan.
- Prencipe, S., Nari, L., Vittone, G. & Spadaro, D. 2017. First report of *Diaporthe eres* causing stem canker on peach (*Prunus persica*) in Italy. *Plant Disease* 101(6):1052-1052.
- Proffer, T. J. & Jones, A.L. 1989. A new canker disease of apple caused by *Leucostoma cincta* and other fungi associated with cankers on apple in Michigan. *Plant Disease* 73:508-514.
- Pscheidt, J. W. & Ocamb, C. M. 2020. Pacific Northwest Plant Disease Management Handbook. Oregon State University. <https://pnwhandbooks.org/>. (Archived at PERAL).
- Ristić, D., Stevanović, M., Stošić, S., Vučurović, I., Gašić, K., Gavrilović, V. & Živković, S. 2016. *Diaporthe eres* as a pathogen of quince fruit (*Cidonia oblonga*) in Serbia. Pages 1270-1276 in D. Kovačević, (ed.). VII International Scientific Agriculture Symposium, "Agrosym 2016", 6-9 October 2016, Jahorina, Bosnia and Herzegovina. Proceedings.
- Rumbos, I. C. 1986. *Phialophora parasitica*, causal agent of cherry dieback. *Journal of Phytopathology* 117:283-287.
- Sami, S., Mohammadi, H. & Heydarnejad, J. 2014. *Phaeoacremonium* species associated with necrotic wood of pome fruit trees in Iran. *Journal of Plant Pathology* 96(3):487-495.
- Sessa, L., Abreo, E., Bettucci, L. & Lupo, S. 2016. *Botryosphaeriaceae* species associated with wood diseases of stone and pome fruits trees: symptoms and virulence across different hosts in Uruguay. *European Journal of Plant Pathology* 146(3):519-530.
- Sewell, G. W. F. & Wilson, J. F. 1975. The role of *Thielaviopsis basicola* in the specific replant disorders of cherry and plum. *Ann. Appl. Biol.* 79:149-169.
- Sønderhausen, E. 1970. Forekomst af *Thielaviopsis basicola* (Berk. og Br.) Ferr. i jord og rødder af kirsebærfrøplanter (*Prunus avium* L.) og andre træagtige planter. *Planteavl*:214-223.
- Smit, W. A. 1996. A new canker disease of apple, pear, and plum rootstocks-caused by *Diaporthe ambigua* in South Africa. *Plant Disease* 80:1331-1335.

- Soto-Estrada, A., H. Förster, D. A. DeMason, & J. E. Adaskaveg.** 2005. Initial infection and colonization of leaves and stems of cling peach by *Tranzschelia discolor*. *Phytopathology*® 95(8):942-950.
- Strand, L. L.** 1999. Integrated Pest Management for Stone Fruits (Publication 3389). University of California, Agriculture and Natural Resources. 264 pp.
- Sugar, D., & R. A. Spotts.** 1992. Sources of inoculum of *Phialophora malorum*, causal agent of side rot of pear. *Phytopathology* 82:735-738.
- Sutton, T. B., Alswinkle, H. S., Agnello, A.M. & Walgenbach, J. F.** 2014. Compendium of apple and pear diseases and pests (Second Edition). American Phytopathological Society, St. Paul, MN, USA. 218 pp.
- Thomas, H. E., S. Wilhelm, & N. A. MacLean.** 1953. Two root rots of fruit trees. *Yearbook of Agriculture*:702-705.
- Úrbez-Torres, J. R., Boulé, J. & O’Gorman, D. T.** 2016. First report of *Diplodia seriata* and *D. mutila* causing apple dieback in British Columbia. *Plant Disease* 100(6):1243.
- USDA.** 2020. USDA Crop Profiles. United States Department of Agriculture (USDA), National Institute of Food and Agriculture. https://ipmdata.ipmcenters.org/source_list.cfm. (Archived at PERAL).
- Van den Berg, N., Hartley, J., Engelbrecht, J., Mufamadi, Z., van Rooyen, Z. & Mavuso, Z.** 2018. First report of white root rot caused by *Rosellinia necatrix* on *Persea americana* in South Africa. *Plant Disease* 102(9):1850.
- Van Leeuwen, G. C. M., Baayen, R. P., Holb, I. J. & Jeger, M. J.** 2002. Distinction of the Asiatic brown rot fungus *Monilia polystroma* sp. nov. from *M. fructigena*. *Mycol. Res.* 106:444-451.
- Vovey Jr., R. P.** 1967. *Botryosphaeria* canker on apple in Washington. *Plant Disease Reporter* 51:593-594.
- Webster, T., Berrie, A., Johnson, D., Biddlecombe, T., Pennell, D., Luton, M. & Guest, J.** 2020. Apple best practice guide: Blossom wilt (*Monilinia laxa* f.sp. *malí*). AHDB Horticulture.
- Wilcox, W. F.** 1994. Sooty Blotch *Gloeodes pomigena* (Schwein.) Colby and Flyspeck *Schizothyrium pomi* (Mont. and Fr.) Arx (Disease Identification Sheet No. D11). Cornell Cooperative Extension.
- Windbiel-Rojas, K., C. Laning, L. Blecker, & M. Alfaro.** 2020. Pests in Gardens and Landscapes. Agriculture and Natural Resources, University of California. <http://ipm.ucanr.edu/PMG/>.
- Worrall, J.** 2020. Forest Pathology: Incense cedar rust. 2020 forestpathology.org, Vancouver, Washington, USA. Last accessed 7/15/2020, <https://forestpathology.org/rusts/incense-cedar-rust/>.
- WPFUS.** 2020. Distribution by State. Widely Prevalent Fungi of the United States (WPFUS). <https://www.prevalentfungi.org/>.
- Wunderlich, L. R., Varela, L. G., Gubler, W. D., Westerdahl, B. B., Caprile, J. L., Grant, J. A., Johnson, R. S. & Vossen, P. M.** 2017. UC IPM Pest Management Guidelines: Apple (UC ANR Publication 3432). UC Statewide Integrated Pest Management Program, Oakland, CA.
- Xiao, C. L., & Boal, R. J.** 2005. Distribution of *Potebniamyces pyri* in the Pacific Northwest and its association with a canker and twig disease of pear trees. *Plant Dis.* 89: 920-925.
- Xiao, C. L., Rogers, J. D., Kim, Y. K. & Liu, Q.** 2005. *Phacidiopycnis washingtonensis*—a new species associated with pome fruits from Washington State. *Mycologia* 97(2):464-473.
- Xu, C., C. Wang, L. Ju, R. Zhang, A. R. Biggs, E. Tanaka, B. Li, & G. Sun.** 2015. Multiple locus genealogies and phenotypic characters reappraise the causal agents of apple ring rot in China. *Fungal Diversity* 71(1):215-231.
- Yun, H. Y., Hong, S. G., Rossman, A. Y., Lee, S. K., Lee, & K. J. & Bae, K. S.** 2009. The rust fungus *Gymnosporangium* in Korea including two new species, *G. monticola* and *G. unicornae*. *Mycologia* 101(6):780-809.
- Zlatković, M., Wingfield, M.J., Jami, F. & Slippers, B.** 2018. Host specificity of co-infecting *Botryosphaeriaceae* on ornamental and forest trees in the Western Balkans. *Forest Pathology* 48(2):e12410.

Table 4. Bacterial pathogens of stone and pome fruit trees

Last updated December 2021.

PEST	MAIN HOST(S)	DISEASE	REFERENCES	PRESENCE / ABSENCE		
				CAN	USA	MEX
<i>Candidatus</i> Phytoplasma asteris' Lee <i>et al.</i> (16Srl-B)	<i>P. avium</i> , <i>P. persica</i> , <i>Pyrus communis</i> , <i>Malus domestica</i>	Aster yellows, apple sessile leaf, sweet/sour cherry decline	Caglayan <i>et al.</i> , 2013; Davis, 2020; Jomantiene and Davis, 2005; Lee <i>et al.</i> , 2004; Zunnoon-Khan <i>et al.</i> , 2010b	Ab1	P2	P2
<i>Candidatus</i> Phytoplasma mali Seemüller and Schneider (16SrX)	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>Malus domestica</i> , <i>Pyrus</i> spp.	Apple Proliferation, Sweet/sour cherry decline	Bulgariet <i>et al.</i> , 2012; Mehle <i>et al.</i> , 2006	Ab1	Ab1	A1
<i>Candidatus</i> Phytoplasma pear decline Taiwan	<i>Pyrus</i> spp.	Pear decline Taiwan	Liu <i>et al.</i> , 2007	Ab1	Ab1	Ab1
<i>Candidatus</i> Phytoplasma phoenicium' Verdin <i>et al.</i> (16SrlX-B)	<i>Prunus dulcis</i> , <i>Prunus</i> spp.	Almond's witches broom, almond brooming	Salehi <i>et al.</i> , 2006; Zirak <i>et al.</i> , 2019	Ab1	Ab1	Ab1
<i>Candidatus</i> Phytoplasma pruni Davis <i>et al.</i> (16SrlII-A)	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. cerasus</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i> , <i>P. salicina</i> , <i>Prunus</i> spp.	X-disease, Cherry buckskin, Cherry Western X disease, Prunus X disease	Davis <i>et al.</i> , 2013; Davis <i>et al.</i> , 2019	P5	P2	P2
<i>Candidatus</i> Phytoplasma prunorum' Seemüller and Schneider (16SrX-B)	<i>Prunus armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. salicina</i> , <i>Prunus</i> spp.	European stone fruit yellows, apricot chlorotic leafroll, plum leptonecrosis, plum decline, peach yellows, peach decline, European peach yellows	Krizanac <i>et al.</i> , 2010	Ab1	Ab1	Ab1
<i>Candidatus</i> Phytoplasma pyri' (16SrX-C) Seemüller and Schneider., Syn.: Phytoplasma pyri Seemüller and Schneider	<i>P. persica</i> , <i>Pyrus</i> spp.	Pear decline, Peach yellow leaf roll, peach decline	EPPO, 2017; Hunter <i>et al.</i> , 2010; Sabate <i>et al.</i> , 2014	P2	P2	Ab1
<i>Candidatus</i> Phytoplasma solani'	<i>Malus domestica</i> , <i>P. armeniaca</i> , <i>P. persica</i>	none	Quaglino <i>et al.</i> , 2013	Ab1	Ab1	Ab1

PEST	MAIN HOST(S)	DISEASE	REFERENCES	PRESENCE / ABSENCE		
				CAN	USA	MEX
Quaglino <i>et al.</i> (16SrXII-A)						
<i>Candidatus Phytoplasma ziziphi</i> Jung <i>et al.</i> (16SrV-B)	<i>Malus</i> spp., <i>P. salicina</i>	none	Li <i>et al.</i> , 2014; Gao <i>et al.</i> , 2019	Ab1	Ab1	Ab1
<i>Dickeya fangzhongdai</i> Tian, Zhao, Yuan, Yi, Fan, Xu, <i>et al.</i>	<i>Pyrus</i> spp.	Bleeding canker	Zhao <i>et al.</i> , 2018	Ab1	Ab3	Ab1
<i>Erwinia amylovora</i> (Burrill) Winslow <i>et al.</i>	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus domestica</i> , <i>Chaenomeles</i> spp.	Fire blight	Emeriewen <i>et al.</i> , 2019; Norin and Rumpunen, 2003; Sutton <i>et al.</i> , 2014; Vegh <i>et al.</i> , 2012	P1	P2	P2
<i>Erwinia pyrifoliae</i> Kim <i>et al.</i>	<i>Pyrus pyrifolia</i>	Asian pear blight	Sutton <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Erwinia uzenensis</i> Matsuura <i>et al.</i>	<i>Pyrus</i> spp.	Bacterial black shoot disease of European pear	Matsura <i>et al.</i> , 2012	Ab1	Ab1	Ab1
<i>Pseudomonas amygdale</i> Psallidas and Panagopoulos	<i>Prunus</i> spp.	Almond bacteriosis	Sutton <i>et al.</i> , 2014; Ruinelli <i>et al.</i> , 2019	Ab1	Ab1	Ab1
<i>Pseudomonas syringae</i> pv. <i>morsprunorum</i> (Wormald) Young <i>et al.</i>	<i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i>	Bacterial canker of stone fruits; dieback	Ahmed <i>et al.</i> , 2018; Hulin <i>et al.</i> , 2018; Latorre and Jones 1979; Sutton <i>et al.</i> , 2014	P1	P2	Ab1
<i>Pseudomonas syringae</i> pv. <i>papulans</i> (Rose) Dhanvantari	<i>Malus</i> spp.	Blister spot	Kerkoud <i>et al.</i> , 2000	P2	P2	Ab1
<i>Pseudomonas syringae</i> pv. <i>persicae</i> Prunier, Luisetti and Gardan	<i>Prunus</i> spp.	Bacterial decline of stone fruits, Bacterial shoot dieback	Hulin <i>et al.</i> , 2018; Sutton <i>et al.</i> , 2014; Zhao <i>et al.</i> , 2015	Ab1	Ab1	Ab1
<i>Pseudomonas syringae</i> pv. <i>syringae</i> van Hall	<i>Malus</i> spp., <i>Pyrus</i> spp., <i>P. avium</i> , <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. salicina</i>	Bacterial canker and fruit spot; blossom blast of pear; blister bark of apple	Gasic <i>et al.</i> , 2018; Hulin <i>et al.</i> , 2018; Little <i>et al.</i> , 1998; Ogawa <i>et al.</i> , 1995; Sutton <i>et al.</i> , 2014; Xu <i>et al.</i> , 2008	P1	P2	Ab3
<i>Rhizobium radiobacter</i> (Smith and Townsend) Young <i>et al.</i> , syn.: <i>Agrobacterium tumefaciens</i> (Smith and Townsend) Conn	<i>Malus</i> spp., <i>P. armeniaca</i> , <i>Prunus</i> spp., <i>Pyrus</i> spp.	Crown gall	Aysan <i>et al.</i> , 2003; CABI 2021; Moore and Putnam, 2020	P1	P2	P2

PEST	MAIN HOST(S)	DISEASE	REFERENCES	PRESENCE / ABSENCE		
				CAN	USA	MEX
<i>Rhizobium rhizogenes</i> (Riker <i>et al.</i> 1930) Young <i>et al.</i> , syn.: <i>Agrobacterium rhizogenes</i> (Riker <i>et al.</i> , 1930) Conn	<i>Prunus avium</i> , <i>Prunus</i> spp.	Crown gall, Hairy root of apple	Pulawska <i>et al.</i> , 2016	P2	P2	P2
<i>Xanthomonas arboricola</i> pv. <i>pruni</i> (Smith) Vauterin Hoste, Kersters and Swings	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i>	Leaf and fruit spot and stem canker or bacterial canker of stone fruit	Garita-Cambronero, <i>et al.</i> , 2018	P2	P2	Ab1
<i>Xanthomonas prunicola</i> . The type strain is CFBP 8353, syn. CECT 9404=IVIA 3287.1)	<i>Prunus persica</i> , <i>P. persica</i> var. <i>nectarina</i>	Necrotic lesions on leaves	López, <i>et al.</i> , 2018	Ab1	Ab1	Ab1
<i>Xylella fastidiosa</i> subsp. <i>pauca</i> Schaad <i>et al.</i>	<i>Prunus avium</i>	Citrus variegated chlorosis	Saponari <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Xylella fastidiosa</i> Wells <i>et al.</i>	<i>P. domestica</i> , <i>P. dulcis</i>	Phony peach disease; plum leaf scald; almond scorch; Pierce's disease	Alves <i>et al.</i> , 2004; Chen <i>et al.</i> , 2008; Janse and Obradovic, 2010; Sutton <i>et al.</i> , 2014	P7	P2	P3

References

- Ahmed, R., Inam-ul-Haq, M., Shahzad, U., Hyder, S., Shahzaman, S., Khan, A. U. R., Aatif, H. M., Ahmad, A. & Gondal A. S. 2018. First report of bacterial canker caused by *Pseudomonas syringae* pv. *morsporunorum* race 1 on peach from Khyber Pakhtunkhwa province of Pakistan. *Plant Disease* 102(10):2027.
- Alves, E., Marruci, C. R., Lopes, J. R. S. & Leite, B. 2004. Leaf symptoms on plum, coffee and citrus and the relationship with the extent of xylem vessels colonized by *Xylella fastidiosa*. *Journal of Phytopathology* 152:291-297.
- Aysan, Y., Sahin, F., Mirik, M., Donmez, M. F. & Tekman, H. 2003. First report of crown gall of apricot (*Prunus armeniaca*) caused by *Agrobacterium tumefaciens* in Turkey. *Plant Pathology* 52:793.
- Bulgari, D., Bozkurt, A. I., Casati, P., Caglayan, K., Quaglino, F. & Bianco, P. A. 2012. Endophytic bacterial community living in roots of healthy and '*Candidatus Phytoplasma mali*'-infected apple (*Malus domestica*, Borkh.) trees. *Antonie van Leeuwenhoek* 102:677-687.
- CABI. 2021. Crop Protection Compendium. Commonwealth Agricultural Bureau International (CABI). <https://www.cabi.org/cpc/>.
- Caglayan, K., Gazel, M., Kucukgol, C., Paltrineri, S., Contaldo, N. & Bertaccini, A. 2013. First report of '*Candidatus Phytoplasma asteris*' (group 16SrI-B) infecting sweet cherries in Turkey. *Journal of Plant Pathology* 54: S4.69-S64.77.
- Chen, J., Livingston, S., Grovers, R., & Civerolo, E. L. 2008. High throughput PCR detection of *Xylella fastidiosa* directly from almond tissues. *Journal of Microbiology Methods* 73:57-61.
- Davis, R. E., Dally, E. L., Webb, C. & Appel, J. A. 2019. First report of North American Grapevine Yellows (NAGY) in Kansas: Detection of '*Candidatus Phytoplasma pruni*'- and '*Ca. Phytoplasma asteris*'-related strains in diseased plants. *Plant Disease* 103(4):759.
- Davis, R. E. 2020. MPPL Phytoplasma. Agricultural Research Station (ARS). <https://plantpathology.ba.ars.usda.gov/cgi-bin/resource/phytclass.cgi>.
- Davis, R. E., Zhao, Y., Dally, E. L., Lee, I. M., Jomantiene, R. & Douglas, S. M. 2013. '*Candidatus Phytoplasma pruni*', a novel taxon associated with X-disease of stone fruits, *Prunus* spp.: multilocus characterization based on 16S rRNA, secY, and ribosomal protein genes. *International Journal of Systemic and Evolutionary Microbiology* 66:766-776.
- Emeriewen, O. F., T. Wohner, H. Flachowsky, & A. Peil. 2019. *Malus* host-*Erwinia amylovora* interactions: strains pathogenicity and resistance mechanisms. *frontiers in Plant Science* 10(551):1-.
- EPPO. 2017. PM 7/62 (2) '*Candidatus Phytoplasma mali*, *Ca. P. pyri* and *Ca. P. prunorum*'. *Bulletin OEPP/EPPO Bulletin* 47(2):146-163.
- Garita-Cambronero, J., Palacio-Bielsa, A. & J. Cubero, J. 2018. *Xanthomonas arboricola* pv. *pruni*, causal agent bacterial spot of stone fruits and almond: its genomic and phenotypic characteristics in the *X. arboricola* species context. *Molecular Plant Pathology* 19:2053-2065.
- Gao, R., Yang, S. K., Yan, H. H., Wang, & J. & Lu, X. B. 2019. First report of '*Candidatus Phytoplasma ziziphi*' subgroup 16SrV-B associated with *Prunus salicina* witches'-broom in China. *Plant Disease* 104(2):564.
- Gasic, K., Pavlovic, Z., Santander, R. D., Meredith, & C. & Acimovic, S. G. 2018. First report of *Pseudomonas syringae* pv. *syringae* associated with Bacterial Blossom Blast on apple (*Malus pumila*) in the United States. *Plant Disease* 102(9):1848.
- Hulin, M. T., Mansfield, J. W., Brain, P., Xu, X., Jackson, R. W. & R. J. Harrison, R. J. 2018. Characterization of the pathogenicity of strains of *Pseudomonas syringae* towards cherry and plum. *Plant Pathology* 67:1177-1193.
- Hunter, D. M., Svircev, M., Kaviani, M., Michelutti, R., Wang, L. & Thompson, D. 2010. First report of Pear Decline caused by '*Candidatus Phytoplasma pyri*' in Ontario, Canada. *Plant Disease* 94(5):634.
- Janse, J. D. & Obradovi, Ac. 2010. *Xylella fastidiosa*: its biology, diagnosis, control and risks. *Journal of Plant Pathology* 91: S1.35-S31.48.

- Jomantiene, R. & Davis, R. E. 2005.** Apple sessile leaf: a new disease associated with a '*Candidatus Phytoplasma asteris*' subgroup 16Srl-B phytoplasma in Lithuania. *Plant Pathology* 54:237.
- Kerkoud, M., C. Manceau, L. Gardan, R. Samson, & J.-P. Paulin. 2000.** Epiphytic occurrence of *Pseudomonas syringae* pv. *papulans* (Rose) in France, where blister spot has never been seen *European Journal of Plant Pathology* 106:481-485.
- Krizanac, I., Mikec, I., Budinscak, Z., Seruga Music, M., & D. Skoric, D. 2010.** Diversity of phytoplasma infecting fruit trees and their vector in Croatia. *Journal of Plant Diseases & Protection* 117(5):206-213.
- Latorre, B. A. & A. L. Jones, J.A. 1979.** *Pseudomonas morsprunorum*, the cause of bacterial canker of sour cherry in Michigan, and its epiphytic association with *P. syringae*. *Phytopathology* 69(4):335-339.
- Lee, I. M., Gundersen-Rindal, D. E., Davis, R. E., Bottner, K. D., Marcone, C. & E. Seemüller, E. 2004.** '*Candidatus Phytoplasma asteris*', a novel phytoplasma taxon associated with aster yellows and related diseases. *International Journal of Systemic and Evolutionary Microbiology* 54:1037-1048.
- Li, Z. N., Bai, Y. B., Liu, P., Zhang, L. & We, Y. F. 2014.** Occurrence of '*Candidatus Phytoplasma ziziphi*' in apple trees in China. *Forest Pathology* 44:417-419.
- Little, E. L., Bostock, R. M. & Kirkpatrick, B. C. 1998.** Genetic characterization of *Pseudomonas syringae* pv. *syringae* strains from stone fruits in California. *Applied and Environmental Microbiology* 64(10):3818-3823.
- Liu, H.-L., Chen, C.-C. & Lin, C.-P. 2007.** Detection and identification of the phytoplasma associated with Pear decline in Taiwan. *European Journal of Plant Pathology* 117:281-291.
- López, M. M., Lójepez-Soriano, P., Garita-Cambronero, J., Beltrán, C., Taghoti, G., Portier, P., Cubero, J., Fisher-Le Saux, M & Marco-Noales, E. 2018.** *Xanthomonas prunicola* sp. nov., a novel pathogen that affects nectarine (*Prunus persica* var. *nectarina*) trees. *International Journal of Systematic and Evolutionary Biology* 68(6): 1857-1866.
- Matsura, T., Mizuno, A., Tsukamoto, T., Shimizu, Y., Saito, N., Sato, S., Kikuchi, S., Uzuki, T., Azegami, K. & Sawada, H. 2012.** *Erwinia uzensis* sp. nov., a novel pathogen that affects European pear trees (*Pyrus communis* L.). *International Journal of Systemic and Evolutionary Microbiology* 62:1799-1803.
- Mehle, N., Brzin, J., Boben, J., Hren, M., Frank, J., Petrociv, N., Gruden, K., Dreo, T., Zezlina, I., Seljak, G. & Ravnkar, M. 2006.** First report of '*Candidatus Phytoplasma mali*' in *Prunus avium*, *P. armeniaca* and *P. domestica*. *New Disease Reports* 14:42.
- Moore, L. W. & Putnam, M. L. 2020.** Crown Gall Disease of Nursery Crops. *Pacific Northwest Pest Management Handbooks*. Last accessed 5/21/2020, <https://pnwhandbooks.org/plantdisease/pathogen-articles/pathogens-common-many-plants/bacteria-other-prokaryotes/crown-gall>.
- Norin, I., & K. Rumpunen. 2003.** Pathogens on Japanese quince (*Chaenomeles japonica*) plants in Japanese quince- Potential fruit crop for Northern Europe. 37-58 pp.
- Ogawa, J. M., Zehr, E.I., Bird, G. W., Ritchie, D. F., Uriu, K., & Uyemoto, J. K. 1995.** Compendium of Stone Fruit Diseases. American Phytopathological Society, St. Paul, MN, USA.
- Pulawska, J., Warabieda, W. & Ismail, E. 2016.** Identification and characterization of bacteria isolated from crown galls on stone fruits in Poland. *Plant Pathology* 65:1034-1043.
- Quaglino, F., Zhao, Y., Casati, P., Bulgari, D., Bianco, P. A., Wei, W. & Davis, R. E. 2013.** '*Candidatus Phytoplasma solani*', a novel taxon associated with stolbur- and bois noir-related diseases of plants. *International Journal of Systemic and Evolutionary Microbiology* 63:2879-2894.
- Ruinelli, M., Blom, J., Smits, T. H. M. & Pothier, J. F. 2019.** Comparative genomics and pathogenicity potential of members of the *Pseudomonas syringae* species complex on *Prunus* spp. *BMC Genomics* 20(172):1-16.
- Sabaté, J., Laviña, A. & Batlle, A. 2014.** First report of '*Candidatus Phytoplasma pyri*' causing Peach Yellow Leaf Roll (PYLR) in Spain *Plant Disease* 98(7):989.
- Saponari, M., Boscia, D., Loconsole, G., Palmisano, F., Potere, O., Savino, V. & Martelli, G. P.**

2014. New hosts of *Xylella fastidiosa* strain CoDiRo in Apulia. *Journal of Plant Pathology* 96(3):603-611.
- Salehi, M., Izadpanah, K. & Heydarnejad, J.** 2006. Characterization of a new almond witches' broom phytoplasma in Iran. *Journal of Phytopathology* 154:386-391.
- Sutton, T. B., Aldwinckle, H. S., Agnello, A. M., Walgenbach, J. F.** 2014. *Compendium of Apple and Pear Diseases and Pests*. American Phytopathological Society. St. Paul, MN, USA. 218pp.
- Végh, A., Z. Némethy, L. Hajagos, & L. Palkovics.** 2012. First report of *Erwinia amylovora* causing fire blight on plum (*Prunus domestica*) in Hungary. *Plant Disease* 96(5):759.
- WPBUS.** 2020. Distribution by State. Widely Prevalent Bacteria of the United States (WPBUS). <https://www.prevalentbacteria.org/bacteria.cfm>.
- Xu, L. H., Xie, G. L., Li, B., Zhu, B., Xu, S. & Qian, J.** 2008. First report of Pear Blossom Blast caused by *Pseudomonas syringae* pv. *syringae* in China. *Plant Disease* 92(5):832.
- Zhao, W., Jiang, H., Tian, Q. & Hu, J.** 2015. Draft genome sequence of *Pseudomonas syringae* pv. *persicae* NCPPB 2254. *Genome Announcements* 3(3):1.
- Zhao, Y., Tian, Y., Li, X. & Hu, B.** 2018. Complete genome sequence of a *Dickeya fangzhongdai* type strain causing bleeding canker of pear tree trunks. *Genome Announcements* 6(20):1.
- Zirak, L., Bahar, M. & Ahoonmanesh, A.** 2009. Characterization of phytoplasmas associated with almond diseases in Iran *Journal of Phytopathology* 157:736-741.
- Zunnoon-Khan, S., Michelutti, R., Arocha-Rosete, Y., Scott, J., Crosby, W. & Bertaccini, A.** 2010. First report of '*Candidatus Phytoplasma asteris*'-related strain associated with Peach Rosette in Canada. *Plant Disease* 94(7):916.

Table 5: Nematode pests of stone and pome fruit trees

Last updated December 2021.

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Criconema mutabile</i> Taylor	Ectoparasite	<i>Cydonia oblonga</i> , <i>Prunus persica</i> , <i>Prunus domestica</i> , <i>P armeniaca</i>	Hugo and Storey, 2017; Zasada <i>et al.</i> , 2019	Ab1	P2	P2
<i>Criconemoides curvatum</i>	Ectoparasite	<i>Prunus persica</i> , <i>Prunus avium</i>	Bridge and Starr, 2007	P2	P7	Ab3
<i>Criconemoides ornatus</i>	Ectoparasite	<i>Prunus persica</i>	Ratanaworabhan and Smart, 1970	Ab1	Ab3	Ab1
<i>Helicotylenchus digonicus</i> Perry	Migratory ecto and semi endoparasite	<i>Malus</i> spp., <i>Pyrus</i> spp.	Khan <i>et al.</i> , 2013; Siddiqui <i>et al.</i> , 1973	P2	P2	P2
<i>Helicotylenchus dihystera</i> (Cobb) Sher	Migratory ecto and semi endoparasite	<i>Cydonia oblonga</i> , <i>Malus</i> spp., <i>Prunus avium</i> , <i>Prunus salicina</i> , <i>Prunus persica</i>	Marais and Swart, 1998; Ogawa <i>et al.</i> , 1995; Poiras <i>et al.</i> , 2013; Sharma <i>et al.</i> , 2005	P7	P2	P2
<i>Helicotylenchus erythrinae</i> (Zimmermann) Golden	Migratory ecto and semi endoparasite	<i>Malus</i> spp., <i>Prunus persica</i>	Siddiqui <i>et al.</i> , 1973; Zavaleta-Mejia and Sosa-Moss, 1979	P7	P2	P2
<i>Helicotylenchus platyurus</i>	Migratory ecto and semi endoparasite	<i>Prunus</i> spp.	Hafez <i>et al.</i> , 2010	P2	P2	Ab1
<i>Helicotylenchus pseudorobustus</i> (Steiner) Golden	Migratory ecto and semi endoparasite	<i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus domestica</i> , <i>Prunus persica</i> var <i>nucipersica</i> , <i>Prunus persica</i>	Subbotin <i>et al.</i> , 2015	P2	P2	Ab1
<i>Longidorus africanus</i> Merny	Root tip ectoparasite	<i>Prunus</i> spp.	Xu and Zhao, 2019	Ab1	P2	Ab1
<i>Longidorus attenuatus</i> Hooper	Root tip ectoparasite	<i>Pyrus communis</i>	Arias and Andres, 1989; Andres <i>et al.</i> , 1991; CABI 2021 edition; Griffiths and Robertson, 1984; Raski 1988	Ab1	Ab1	Ab1
<i>Longidorus breviannulatus</i> Norto and Hoffmann	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus avium</i> , <i>Prunus persica</i>	Simard <i>et al.</i> , 2009; Van Driell <i>et al.</i> , 1990; Xu and Zhao, 2019	P2	P2	Ab1
<i>Longidorus brevicaudatus</i> Norton and Hoffmann	Root tip ectoparasite	<i>Malus</i> spp.	Lone <i>et al.</i> , 2018	Ab1	Ab1	Ab1
<i>Longidorus caespiticola</i> Hopper, [<i>Longidorus</i>	Root tip ectoparasite	<i>Malus domestica</i> , <i>Prunus avium</i> , <i>Prunus</i>	Arias and Andres, 1989; Ravichandra, 2008; Xu and Zhao,	Ab1	Ab1	Ab1

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>coespiticola</i> maybe a variant spelling of species name]		<i>domestica</i>	2019			
<i>Longidorus diadecturus</i> Eveleigh and Allen	Root tip ectoparasite	<i>Prunus persica</i>	Eveleigh and Allen, 1982; Allen <i>et al.</i> , 1984	P2	P2	Ab1
<i>Longidorus elongatus</i> (de Man) Micoletzky	Root tip ectoparasite	<i>Malus domestica</i> , <i>Malus sylvestris</i> , <i>Prunus persica</i> , <i>Prunus communis</i>	Xu and Zhao, 2019	P7	P2	Ab1
<i>Longidorus euonymus</i> Mali and Hooper	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus</i> spp.	Liskova <i>et al.</i> , 2007; Lone <i>et al.</i> , 2018; Xu and Zhao, 2019	Ab1	Ab1	Ab1
<i>Longidorus macrosoma</i> Hopper	Root tip ectoparasite	<i>Prunus avium</i> , <i>Pyrus communis</i>	Andres <i>et al.</i> , 1991; Arias and Andres, 1989; Raski, 1988; Xu and Zhao, 2019	Ab1	Ab1	Ab1
<i>Longidorus mirus</i> Khan, Chawla and Seshadri	Root tip ectoparasite	<i>Malus domestica</i>	Lone <i>et al.</i> , 2018	Ab1	Ab1	Ab1
<i>Meloidogyne arenaria</i> (Neal)	Sedentary endoparasite	<i>Prunus persica</i> , <i>Malus</i> spp.	Cid del Prado <i>et al.</i> , 2001; Hugo and Storey, 2017; Power <i>et al.</i> , 2005	P7	P2	P2
<i>Meloidogyne floridensis</i> Handoo, Nyczepir, Esmenjaud, van der Beek, Castognone-Sereno, Carta, Skantar and Higgins	Sedentary endoparasite	<i>Prunus persica</i>	Chitambar, 2018; Shirley, 2013	Ab1	P2	Ab1
<i>Meloidogyne hapla</i> Chitwood	Sedentary endoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus domestica</i> , <i>Prunus persica</i>	Liskova <i>et al.</i> , 2007; Ogawa <i>et al.</i> , 1995; Power <i>et al.</i> , 2005	P2	P2	P2
<i>Meloidogyne incognita</i> (Kofoid and White) Chitwood	Sedentary endoparasite	<i>Chaenomeles</i> spp., <i>Malus pumila</i> , <i>Prunus persica</i>	Maquilan <i>et al.</i> , 2018; Ogawa <i>et al.</i> , 1995; Powers <i>et al.</i> , 2005; Rumpunen, 2002; Khan <i>et al.</i> , 2013	P2	P2	P2
<i>Meloidogyne javanica</i> (Treub) Chitwood	Sedentary endoparasite	<i>Cydonia oblonga</i> , <i>Prunus persica</i>	Cid del Prado <i>et al.</i> , 2001; Maquilan <i>et al.</i> , 2018; Marais and Stewart, 1998; Ogawa <i>et al.</i> , 1995; Powers <i>et al.</i> , 2005	P7	P2	P2
<i>Meloidogyne mali</i> Itoh, Ohshima and Ichinoche	Sedentary endoparasite	<i>Malus</i> spp.	Bridge and Starr, 2007; Itoh <i>et al.</i> , 1969	Ab1	P2	Ab1
<i>Merlinius brevidens</i> (Allen)	Root tip ectoparasite	<i>Prunus</i> spp.	Dong <i>et al.</i> , 2007; MKenry and	P7	P2	Ab1

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
Siddiqi, syn. <i>Geocenamus brevidens</i> (Allen) Siddiqi			Roberts, 1985			
<i>Mesocriconema curvatum</i> (Raski) Loof and de Grisse (accepted name) syn. <i>Criconemella curvatum</i>	Ectoparasite	<i>Malus</i> spp., <i>Prunus persica</i>	Bridge and Starr, 2007; Hafez <i>et al.</i> , 1992	Ab1	P2	Ab1
<i>Mesocriconema ornatum</i> syn. <i>Criconemella ornata</i> (Raski) Luc and Raski, <i>Criconemella ornatum</i>	Ectoparasite	<i>Malus</i> spp.	Hafez <i>et al.</i> , 2010	Ab1	P2	Ab1
<i>Mesocriconema xenoplax</i> (Raski) Luc and Raski, syn. <i>Macroposthonia xenoplax</i> (Raski) De Grisse and Loof, <i>Criconemella xenoplax</i> (Raski) Luc and Raski	Ectoparasite	<i>Prunus persica</i> , <i>Prunus</i> spp.	Dong <i>et al.</i> , 2007; Ferris <i>et al.</i> , 2004; Ogawa <i>et al.</i> , 1995	P2	P2	P2
<i>Nanidorus minor</i> (Colbran) Siddiqi, syn. <i>Paratrichodorus minor</i> , <i>P. christiei</i> , <i>Trichodorus minor</i> , <i>Trichodorus christiei</i>	Root tip ectoparasite	<i>Cydonia oblonga</i> , <i>Malus domestica</i> , <i>Prunus persica</i> , <i>Pyrus</i> spp., <i>Pyrus communis</i>	Dong <i>et al.</i> , 2007; Hafez <i>et al.</i> , 1992; Hugo and Storey, 2017; Kumari and Subbotin, 2012; Xu and Zhao, 2019	P7	P2	Ab1
<i>Paratrichodorus lobatus</i> (Colbran) Siddiqi, syn. <i>Trichodorus labatus</i> Colbran, <i>Trichodorus clarki</i> Yeates	Root tip ectoparasite	<i>Pyrus communis</i> , <i>Prunus persica</i> , <i>Prunus armeniaca</i>	Hugo and Storey, 2017	Ab1	Ab1	Ab1
<i>Paratrichodorus pachydermus</i> (Seinhorst) Siddiqi	Ectoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus persica</i> , <i>Prunus cerasus</i> , <i>Prunus avium</i>	Kumari, 2010; Ravichandra, 2008	P7	P2	Ab1
<i>Paratrichodorus porosus</i> (Allen) Siddiqi, syn. <i>Trichodorus porosus</i> Allen	Root tip ectoparasite	<i>Malus domestica</i> , <i>Prunus persica</i> , <i>Prunus serrula</i> , <i>Pyrus</i> spp.	Hugo and Storey, 2017; Li <i>et al.</i> , 2020; Siddiqui <i>et al.</i> , 1973; Xu and Zhao, 2019	Ab1	P2	Ab1
<i>Paratylenchus hamatus</i> Thorne and Allen	Migratory endo and ectoparasite	<i>Malus</i> spp., <i>Prunus avium</i> , <i>Prunus persica</i> , <i>Prunus</i> spp., <i>Pyrus</i> sp..	Dong <i>et al.</i> , 2007; Raski, 1975; Ravichandra, 2008; Siddiqui <i>et al.</i> , 1973; Van den Berg <i>et al.</i> , 2014	P2	P2	Ab1
<i>Paratylenchus neglectus</i> Filipjev and Schuurmans-Stekhoven	Migratory endo and ectoparasite	<i>Malus</i> spp., <i>Prunus avium</i> , <i>Prunus persica</i> , <i>Prunus</i> spp.	Carta <i>et al.</i> , 2001; Hafez <i>et al.</i> , 2010; Siddiqui <i>et al.</i> , 1973; Subbotin <i>et al.</i> , 2008; Wang <i>et al.</i> , 2016	P2	P2	Ab1

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
<i>Paratylenchus neoamblycephalus</i> Geraert	Migratory endo and ectoparasite	<i>Prunus persica</i>	Dong <i>et al.</i> , 2001; McKenry and Roberts, 1985; Ravichandra, 2008	Ab1	P2	Ab1
<i>Paratylenchus pratensis</i> (de Man) Filipjev	Migratory endo and ectoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus persica</i> , <i>Prunus</i> spp.	Handoo and Morgan, 1989; Hugo and Storey, 2017; Liskova <i>et al.</i> , 2007	P7	P2	P2
<i>Paratylenchus projectus</i> Jenkins, [<i>Pratylenchus projectus</i> (Incorrect genus name)]	Migratory endo and ectoparasite	<i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus persica</i> , <i>Prunus</i> spp.	Liskova <i>et al.</i> , 2007; Siddiqui <i>et al.</i> , 1973 as cited in Chitambar <i>et al.</i> , 2012	P2	P2	Ab1
<i>Pratylenchus brachyurus</i> (Godfrey) Filipjev and Schuurmans Stekhoven	Migratory endo and ectoparasite	<i>Pyrus malus</i> , <i>Malus</i> spp., <i>Pyrus</i> spp., <i>Prunus</i> spp.	Cepeda and Hernández, 1991; Dong <i>et al.</i> , 2007; Jones and Aldwinckle, 1990; McKenry and Roberts, 1985; Oliveira <i>et al.</i> , 1999; Siddiqui <i>et al.</i> , 1973	Ab3	P2	P2
<i>Pratylenchus coffeae</i> (Zimmermann) Filipjev and Schuurmans Stekhoven	Migratory endo and ectoparasite	<i>Malus domestica</i> , <i>Malus</i> spp., <i>Pyrus</i> spp.	Hafez <i>et al.</i> , 1992; Hafez <i>et al.</i> , 2010; Silva and Inomoto, 2002; Sutton <i>et al.</i> , 2014; Wang <i>et al.</i> , 2016	Ab1	P2	Ab3
<i>Pratylenchus crenatus</i> Loof	Migratory endo and ectoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus persica</i>	Brown <i>et al.</i> , 1980; Hafez <i>et al.</i> , 2010; Liskova <i>et al.</i> , 2007; Siddiqui <i>et al.</i> , 1973; Sutton <i>et al.</i> , 2013	P2	P2	Ab1
<i>Pratylenchus penetrans</i> (Cobb) Filipjev and Schuurmans-Stekhoven	Migratory endo and ectoparasite	<i>Cydonia oblongta</i> , <i>Malus domestica</i> , <i>Prunus armeniaca</i> , <i>Prunus persica</i> , <i>Prunus avium</i>	Carta <i>et al.</i> , 2001; Khan <i>et al.</i> , 2013; Liskova <i>et al.</i> , 2007; Ogawa <i>et al.</i> , 1995; Potter <i>et al.</i> , 1984; Subbotin <i>et al.</i> , 2008; Villalobos <i>et al.</i> , 1980; Wang <i>et al.</i> , 2016;	P1	P2	Ab3
<i>Pratylenchus scribneri</i> Steiner	Migratory endo and ectoparasite	<i>Malus</i> spp.	Handoo and Morgan, 1989; Hugo and Storey, 2017	Ab1	P2	Ab3
<i>Pratylenchus thornei</i> Sher and Allen	Migratory endo and ectoparasite	<i>Malus domestica</i> , <i>Prunus armeniaca</i> , <i>Prunus persica</i>	Handoo and Morgan, 1989; Liskova <i>et al.</i> , 2007; Subbotin <i>et al.</i> , 2008	P7	P2	P2
<i>Pratylenchus vulnus</i> Allen and Jensen	Migratory endo and ectoparasite	<i>Malus domestica</i> , <i>Prunus</i> spp., <i>Pyrus communis</i>	Chitambar and Raski, 1984; Handoo and Morgan, 1989; Hugo and Storey, 2017; Ogawa <i>et al.</i> , 1995	Ab4	P2	Ab3
<i>Pratylenchus zaeae</i> Graham	Migratory endo and	<i>Cydonia oblonga</i> ,	Hugo and Storey, 2017; Marais	P7	P2	P2

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
	ectoparasite	<i>Pyrus communis</i>	and Swart, 1998			
<i>Quinisulcius acutus</i> (Allen) Siddiqi, syn. <i>Tylenchorhynchus acutus</i> Allen	Ectoparasite	<i>Malus</i> spp., <i>Prunus persica</i>	Siddiqui <i>et al.</i> , 1973; Hafez <i>et al.</i> , 2010	P7	P2	Ab1
<i>Quinisulcius capitatus</i> (Allen) Siddiqi, syn. <i>Tylenchorhynchus capitatus</i>	Ectoparasite	<i>Malus</i> spp., <i>Prunus</i> spp.	Siddiqui <i>et al.</i> , 1973	P7	P2	Ab3
<i>Rotylenchus robustus</i> (de Man) Filipev syn. <i>Rotylenchus fallorobustus</i> Sher	Semiendoparasite	<i>Malus</i> spp., <i>Prunus persica</i> , <i>Prunus</i> spp.	Cantalapiedra-Navarrete <i>et al.</i> , 2013; Dong <i>et al.</i> , 2007; Liskova <i>et al.</i> , 2007; Siddiqui <i>et al.</i> , 1973	P7	P2	Ab1
<i>Scutellonema brachyurus</i> (Steiner) Andrassy	Semiendoparasite	<i>Prunus persica</i> , <i>Prunus</i> spp.	Dong <i>et al.</i> , 2007	Ab1	P2	Ab1
<i>Tylenchorhynchus annulatus</i> (Cassidy) Golden	Ectoparasite	<i>Prunus</i> spp.	Dong <i>et al.</i> , 2007; Handoo and Morgan, 1989	Ab1	P2	Ab1
<i>Tylenchorhynchus capitatus</i>	Ectoparasite	<i>Malus domestica</i> , <i>Prunus</i> spp.	Allen, 1955; Siddiqui <i>et al.</i> , 1973	P7	P3	Ab3
<i>Tylenchorhynchus clarus</i> Allen	Ectoparasite	<i>Prunus persica</i> , <i>Prunus</i> spp.	Handoo and Morgan, 1989; McKenry and Roberts, 1985; Siddiqui <i>et al.</i> , 1973	Ab1	P2	Ab3
<i>Tylenchorhynchus claytoni</i> Steiner	Ectoparasite	<i>Malus</i> spp., <i>Prunus persica</i>	Siddiqui <i>et al.</i> , 1973	P2	P1	Ab3
<i>Tylenchorhynchus cylindricus</i> Cobb	Ectoparasite	<i>Malus domestica</i> , <i>Prunus armeniaca</i> , <i>Prunus persica</i> , <i>Prunus</i> spp.	Liskova <i>et al.</i> , 2007; Siddiqui <i>et al.</i> , 1973	Ab1	P2	Ab3
<i>Tylenchorhynchus dubius</i> Thorne	Ectoparasite	<i>Prunus</i> spp.	Siddiqui <i>et al.</i> , 1973	P2	P2	Ab1
<i>Tylenchorhynchus ebriensis</i> Seinhorst	Ectoparasite	<i>Prunus</i> spp.	Dong <i>et al.</i> , 2007	Ab1	P2	Ab1
<i>Tylenchorhynchus maximus</i> Allen	Ectoparasite	<i>Malus domestica</i> , <i>Prunus</i> spp.	Chitwood, 1953	P2	P2	Ab1
<i>Tylenchorhynchus nudus</i>	Ectoparasite	<i>Prunus</i> spp.	Dong <i>et al.</i> , 2007	P2	P2	Ab1
<i>Xiphinema americanum</i> Cobb	Root tip ectoparasite	<i>Prunus persica</i> , <i>Malus</i> spp.	Barsi, 1994; Lone <i>et al.</i> , 2018; Martínez, 1980; Ogawa <i>et al.</i> , 1995; Ramírez and Jiménez, 1987; Vrain and Rouselle, 1980; Weimin <i>et al.</i> , 2004; Xu and Zhao, 2019; Ye <i>et al.</i> , 2004;	P2	P1	Ab3
<i>Xiphinema basiri</i> Siddiqui	Root tip ectoparasite	<i>Malus domestica</i>	Lone <i>et al.</i> , 2018; Xu and Zhao,	Ab1	P2	Ab3

PEST	MODE OF PARASITISM	MAIN HOST(S)	REFERENCES	PRESENCE/ABSENCE		
				CAN	USA	MEX
			2019			
<i>Xiphinema brevicolle</i> Bordello and DaCosta	Root tip ectoparasite	<i>Malus pumila</i> , <i>Prunus armeniaca</i> , <i>Prunus persica</i> , <i>Prunus</i> spp., <i>Pyrus</i> spp.	Bridge and Starr, 2007; Oliveira <i>et al.</i> , 2004; Xu and Zhao, 2019	Ab1	P2	Ab1
<i>Xiphinema bricolensis</i> Ebsary, Vrain and Graham	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus</i> spp.	Ebsary <i>et al.</i> , 1989; Singh <i>et al.</i> , 2013; Robbins and Brown, 1991; Vrain and Rouselle, 1980	P2	P2	Ab1
<i>Xiphinema californicum</i> Lamberti and Bleve-Zacheo	Root tip ectoparasite	<i>Malus domestica</i> , <i>Prunus</i> spp.	Georgi, 1988; Bridge and Starr, 2007	Ab1	P2	Ab1
<i>Xiphinema diffusum</i> Lamberti and Bleve-Zecheo	Root tip ectoparasite	<i>Prunus persica</i>	Hugo and Storey, 2017; Oliveira <i>et al.</i> , 2004	Ab1	P2	Ab1
<i>Xiphinema diversicaudatum</i> (Micoletzky) Thorne	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus armeniaca</i>	Hugo and Storey, 2017; Lone <i>et al.</i> , 2018; Weimin <i>et al.</i> , 2004; Xu and Zhao, 2019; Ye <i>et al.</i> , 2004	P7	P2	Ab1
<i>Xiphinema elongatum</i> Schuurmans Stekhoven and Teunisse	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus avium</i> , <i>Pyrus communis</i>	Hugo and Storey, 2017; Lone <i>et al.</i> , 2018; Xu and Zhao, 2019	Ab1	P2	Ab1
<i>Xiphinema index</i> Thorne and Allen	Root tip ectoparasite	<i>Malus domestica</i> , <i>Prunus dulcis</i> , <i>Prunus persica</i> , <i>Pyrus malus</i>	Lone <i>et al.</i> , 2018; Téliz and Goheen, 1968; Weimin <i>et al.</i> , 2004; Xu and Zhao, 2019; Ye <i>et al.</i> , 2004	Ab1	P2	Ab3
<i>Xiphinema insigne</i> Loos	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus</i> spp.	Lambert <i>et al.</i> , 1997; Lone <i>et al.</i> , 2018; Luc and Southey, 1980	Ab1	P2	Ab1
<i>Xiphinema mirus</i>	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus</i> spp.	Lone <i>et al.</i> , 2018	Ab1	Ab1	Ab1
<i>Xiphinema mluci</i> Heyns	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus persica</i> , <i>Pyrus communis</i>	Hugo and Storey, 2017; Xu and Zhao, 2019	Ab1	Ab1	Ab1
<i>Xiphinema occiduum</i>	Root tip ectoparasite	<i>Malus domestica</i>	Ebsary <i>et al.</i> , 1984	P2	Ab1	Ab1
<i>Xiphinema parvistillus</i> Heyns	Root tip ectoparasite	<i>Malus domestica</i> , <i>Prunus persica</i>	Hugo and Storey, 2017; Xu and Zhao, 2019	Ab1	Ab1	Ab1
<i>Xiphinema rivesi</i> Dalmasso	Root tip ectoparasite	<i>Malus</i> spp., <i>Prunus persica</i>	Akinbade <i>et al.</i> , 2014; Ebsary <i>et al.</i> , 1984; Georgi, 1988	P2	P2	Ab1
<i>Xiphinema vuittenezi</i> Luc, Lima, Weischer and Flegg	Root tip ectoparasite	<i>Malus domestica</i> , <i>Malus</i> spp., <i>Prunus armeniaca</i> , <i>Prunus avium</i> , <i>Prunus</i> spp., <i>Pyrus communis</i>	Bride and Starr, 2007; Weimin <i>et al.</i> , 2004; Xu and Zhao, 2019; Ye <i>et al.</i> , 2004	Ab1	P2	Ab1

References

- Akinbade, S. A., Mojtahedi, H., Guerra, L. Eastwell, K., Villamor, D. E. V. & Handoo, Z. A. & Skantar, A. M.** 2014. First Report of *Xiphinema rivesi* (Nematoda, Longidoridae) in Washington State. *Plant Disease* 98:1018.
- Allen, M. W.** 1955. A review of the nematode genus *Tylenchorhynchus*. University of California Publications in Zoology, 61, 129–166.
- Allen, W. R., Van Schagen, J. G., & Ebsary, B. A.** (1984). Comparative transmission of the peach rosette mosaic virus by Ontario populations of *Longidorus diadecturus* and *Xiphinema americanum* (Nematoda: Longidoridae). *Canadian Journal of Plant Pathology*, 6, 29–32.
- Arias, M. & Andrés, M. F.** 1989. Virus-vector nematodes in cereals and fruit crops in Spain. *EPPO Bulletin* 19,625-632.
- Andrés, M., Arias, M. & Bello, A.** 1991. Distribución ecológica del género *Longidorus* (Micoletzky) Filipjev en la Región Central de España. *Nematropica* 21: 79-87.
- Barsi, L.** 1994b. Specimens of the *Xiphinema americanum*-Group (Nematoda: Dorylaimida) on the territory of the former Yugoslavia. *Nematologia Mediterranea*. 22: 25-34.
- Bridge, J. & Starr, J. L.** 2007. *Plant Nematodes of Agricultural Importance: A Color Handbook*. Academic Press Burlington, MA 152pp.
- Brown, M. J., Riedel, R. M. & Rowe R. C.** 1980. Species of *Pratylenchus* Associated with *Solanum tuberosum* cv Superior in Ohio. *Journal of Nematology*. 12: 189-192.
- Cantalapiedra-Navarrete, C., Navas-Cortés, J. A., Liébanas, G., Vovlas, N., Subbotin, S. A., Palomares-Rius, J. E., & Castillo, P.** 2013. Comparative molecular and morphological characterisations in the nematode genus *Rotylenchus*: *Rotylenchus paravitis* n. sp., an example of cryptic speciation. *Zoologischer Anzeiger*, 252, 246–268.
- Carta, K. L., Skantar, M. A. & Handoo, A. Z.** 2001. Molecular, morphological, and thermal characters of 19 *Pratylenchus* spp. and relatives using the D3 segments of the nuclear LSU rDNA gene. *Nematropica*. 31: 193-208.
- Cepeda, S. M., & Hernández, B. J. R.** 1991. Control químico del nematodo de la lesión *Pratylenchus brachyurus* en el cultivo del manzano (*Pyrus malus* L.) en Arteaga, Coahuila. *Memorias XVIII Congreso Nacional de Fitopatología*. Puebla, Pue. Resumen 214.
- Chitambar J. J. & Raski D. J.** 1984. Reactions of grape rootstocks to *Pratylenchus vulnus* and *Meloidogyne* spp. *Journal of Nematology* 16: 166-170.
- Chitambar, J. J.** 2018. California pest rating for *Meloidogyne floridensis* Handoo *et al.*, 2004. California Department Food and Agriculture. <http://blogs.cdfa.ca.gov/Section3162/?p=5666>
- Chitambar, J. J., Westerdahl, B. B. & Subbotin, S. A.** 2012. Plant parasitic nematodes in California agriculture. Pages 92 *in* S. A. Subbotin and J. J. Chitambar, (eds.). *Plant Parasitic Nematodes in Sustainable Agriculture of North America Volume 2: western USA and Mexico*. Springer, Cham.
- Chitwood, B. G.** 1953. Plant parasitic nematode problems in Michigan. Pages 35 *in* N. A. Knobloch, G. W. Bird, eds. *Plant parasitic nematodes of Michigan: with special reference to the genera of the Tylenchorhynchinae* (Nematoda). Michigan Agricultural Research Station.
- Cid del Prado, V. I., Tovar, S. A. & Hernández, J. A.** 2001. Distribución de especies y razas de *Meloidogyne* en México. *Revista Mexicana de Fitopatología*, 19, 32–39.
- Dong, K., Chitambar, J., Subbotin, S., Alzubaidy, M., Luque-Williams, M., Romero, J., Kosta, K. & Luna, R.** 2007. Significant records in Nematology: California statewide nematode survey project for 2006. *California Plant Pest and Damage Report.*, (July 2005 through December 2006), 23, 45-71.
- Ebsary, B. A., Potter, J. W. & Allen, W. R.** (1984). Redescription and distribution of *Xiphinema rivesi* Dalmasso, 1969 and *Xiphinema americanum* Cobb, 1913 in Canada with a description of

Xiphinema occiduum n.sp. (Nematoda: Longidoridae). Canadian Journal of Zoology, 62, 1696–1702.

- Eveleigh, E. S. & Allen, W. R.** 1982. Description of *Longidorus diadecturus n. sp.* (Nematoda: Longidoridae), a vector of the peach rosette mosaic virus in peach orchards in Southwestern Ontario, Canada. Canadian Journal of Zoology, 60, 112–115.
- Ferris, H.; McKenry, M. V.; Jaffee, B. A., Anderson, C. E. & Juurma, A.** 2004. Population Characteristics and Dosage Trajectory Analysis for *Mesocriconema xenoplax* in California *Prunus* Orchards. Journal of nematology. 36: 505-516.
- Hugo, H. J. & Storey, S. G.** 2017. Nematode Pests in Deciduous Fruit. Chapter 15 in: Nematology in South Africa: A View from the 21st Century. edited by Hendrika Fourie, Vaughan W. Spaull, Robin K. Jones, Mieke S. Daneel, Dirk De Waele. Springer International, Switzerland. DOI 10.1007/978-3-319-44210-5_15.
- Georgi, L. L.** 1988. Morphological Variation in *Xiphinema spp.* from New York Orchards. Journal of Nematology 20: 47-57.
- Griffiths, B. S. & Robertson, W. M.** 1984. Morphological and Histochemical Changes Occurring during the Lifespan of Root-tip Galls on *Lolium perenne* Induced by *Longidorus elongatus*. Journal of Nematology 16: 223-229.
- Hafez, S., Golden, A. M., Rashid, F. & Handoo, Z.** 1992. Plant parasitic nematodes associated with crops in Idaho and Eastern Oregon. Nematropica, 22, 193–204.
- Hafez, S. L., Sundararaj, P., Handoo, Z. A. & Siddiqi, M. R.,** 2010. Occurrence and distribution of nematodes in Idaho crops. International Journal of Nematology, 20(1), pp.91-98.
- Handoo, A. Z. & Morgan Golden, A.** 1989. A Key and Diagnostic Compendium to the Species of the Genus *Pratylenchus* Filipjev, 1936 (Lesion Nematodes). Journal of Nematology. 21: 202-218.
- Itoh, Y., Y. Ohshima, Y. & Ichinohe, M.** 1969. A root-knot nematode, *Meloidogyne mali n. sp.* on apple-tree from Japan (Tylenchida: Heteroderidae). Applied Entomology and Zoology 4:194-202.
- Jones, A. L. & Aldwinckle, H. S.** 1990. Compendium of Apple and Pear Diseases. American Phytopathological Society, St. Paul, MN, USA. 100 pp.
- Khan, A., Khatoon, N., Shaukat, S. S. & Kazmi, A. R.** 2013. Nematodes associated with apple (*Malus pumila* Mill) in Balochistan, Pakistan. International Journal of Biological Research 1: 9-11.
- Kumari, S.** 2010. Research Note: Description of *Paratrichodorus pachydermus* (Nematoda: Trichodoridae) from the Czech Republic. HELMINTHOLOGIA, 47, 3: 196 – 198.
- Kumari, S., & Subbotin, S. A.** 2012. Molecular characterization and diagnostics of stubby root and virus vector nematodes of the family Trichodoridae (Nematoda: Triplochoda) using ribosomal RNA genes. Plant Pathology, 61, 1021–1031.
- Lambert, F., Lovev, T., Choleva, B., Brown, D. G. F., Agostinelli, A. & Radicci, V.** 1997. Morphometric variation and juvenile stages of some Longidorid nematodes from Bulgaria with comments on the number of juveniles stages of *Longidorus africanus*, *L. closelongatus* and *Xiphinema santos*. Nematologia Mediterranea. 26: 213-237. Lamberti, F.; Blevé Zacheo, T.; Arias, M. 1982. 1982 The Longidoridae of the Maltese Islands with the description of *Longidorus magnus sp. n.* and *Xiphinema melitense sp. n.* Nematologia Mediterranea., 10: 183-200.
- Li, X., Maria, M., Cai, R., Barsalote, E. M., Peneva, V. & Zheng, J.** 2020. Distribution of trichodorid species in mainland China with description of *Trichodorus hangzhouensis* sp. nov. (Nematoda, Triplonchida). ZooKeys 945.
- Lišková, M., Sasanelli, N. & D’addabbo, T.** 2007. Some notes on the occurrence of plant parasitic nematodes on fruit trees in Slovakia. Plant Protection Science 43:26-32.

- Lone G. M., Zaki, F. A., Pathania S. S. & Mohi, Sajad.** 2018. Population Abundance and Distribution of Ectoparasitic Plant Nematodes Associated with Apple Trees (*Malus domestica*) in Apple Belts of Kashmir, India. *International Journal of Current Microbiology and Applied Sciences*. 7(06): 3817-3824.
- Luc, M. & Southey, J. F.**, 1980. Study of biometrical variability in *Xiphinema insigne* Loos, 1949, and *X. elongatum* Schuurmans Stekhoven & Teunissen, 1938; description of *X. savanicola* n. sp. (Nematoda: Longidoridae) and comments on thelytokous species. *Revue de Nematologie*, 3(2), pp.243-269.
- Maquilan, M. A. D., Sarkhosh, A. & Dickson, D. A.** 2018. Peach root-knot nematode. Pages 5 in. University of Florida, The Institute of Food and Agricultural Sciences (IFAS), Gainesville, FL.
- Martínez, G. M.** (1980). Observaciones sobre la distribución espacial en el suelo de *Xiphinema americanum* Cobb y *Meloidogyne incognita* Chitwood, en viñedos en la costa de Hermosillo. Tesis. Escuela de Agricultura y Ganadería. Universidad de Sonora.
- Marais, M. & Swart, A.** 1998. Plant nematodes in South Africa. 1. Caledon area, Western Cape Province. *African Plant Protection* 4: 27-33
- McKenry, M. V. & Roberts, P. A.** 1985. Phytonematology study guide. Publications, Division of Agriculture and Natural Resources, University of California.
- Ogawa, J. M., Zehr, E. I., Bird, G. W., Ritchie, D. F., Uriu, K. & Uyemoto, J. K.** 1995. Compendium of Stone Fruit Diseases. American Phytopathological Society, St. Paul, MN, USA. pp 61-64.
- Oliveira, C. M. G., Inomoto, M. M., Vieira, A. M. C. & Monteiro, A. R.** 1999. Efeito de Densidades Populacionais de *Pratylenchus brachyurus* no Crescimento de Plantulas de *Coffea arabica* cv. Mundo Novo e *C. canephora* cv. Apoata. *Nematropica*. 29: 215-221.
- Oliveira, C. M. G., Hübschen, J., Brown, Derek J. F., Ferraz, L. C. C. B., Wright, F. & Neilson, R.** 2004. Phylogenetic Relationships Among *Xiphinema* and *Xiphidorus* Nematode Species from Brazil Inferred from 18S rDNA Sequences. *Journal of Nematology* 36: 153-159.
- Poiras, L., Iurcu-Straistraru, E., Bivol, A., Poiras, N. & Cernet, A.** 2013. Plant parasitic and free-living nematodes of some orchards (peach, apple) in the Republic of Moldova. *Stiintele Naturii - Studii si comunicari* 29(2):166-171.
- Potter, J. W., Dirks, V. A., Johnson, P. W., Olthof, T. H. A., Layne, R. E. C. & McDonnell, M. M.** 1984. Response of Peach Seedlings to Infection by the Root Lesion Nematode *Pratylenchus penetrans* under Controlled Conditions. *J Nematol.* 16: 317–322.
- Powers, T. O., Mullin, P. G., Harris, T. S., Sutton, L. A. & Higgins, R.S.** 2005. Incorporating Molecular Identification of *Meloidogyne spp.* into a Large-scale Regional Nematode Survey. *Journal of Nematology* 37: 226-235.
- Ramírez, A. J. A. & Jiménez, L. M.** 1987. Identificación y cuantificación de nematodos fitoparásitos asociados a la vid en la costa de Hermosillo, Sonora. *Memorias del XIV Congreso Nacional de Fitopatología*. Morelia, Michoacán. Resumen 134.
- Raski, D. J.** 1975. Revision of the genus *Paratylenchus* Micoletzky, 1922, and descriptions of new species. Part II of three parts. *Journal of Nematology* 7: 274-295.
- Raski, D.J.** 1988. Nematode Parasites of Grape. In: *Compendium of Grape Diseases*. Pearson. R.C & A.C. Austin (eds). American Phytopathological Society, St. Paul, MN, USA. pp 55-59.
- Ratanaworabhan, S. & Smart, G. C.** 1970. The Ring Nematode, *Criconemoides ornatus*, on Peach and Centipede Grass. *Journal of Nematol.* 2:204-208.
- Ravichandra, N. G.** 2008. *Plant Nematology*. I.K. International Publishing House Pvt. Ltd., New Delhi, India. 693pp.
- Robbins, R. T. & Brown, D. J. F.** 1991. Comments on the Taxonomy, occurrence and distribution of Longidoridae (Nematoda) in North America. *Nematologica* 37: 395-419.
- Rumpunen, K.** 2002. *Chaenomeles*: Potential New Fruit Crop for Northern Europe. In: *Trends in new crops and new uses*. Janick, J. and Whipkey, A. (eds.). ASHS Press, Alexandria, VA. pp 385-392.

- Sharma, G. C., Thakur, B. S. & Kashyap, A. S.** 2005. Impact of npk on the nematode populations and yield of plum (*Prunus salicina*). Pages 433-436 in. International Society for Horticultural Science (ISHS), Leuven, Belgium.
- Shirley, A. M.** 2013. Management of plant-parasitic nematodes on peach utilizing post-plant nematicides and crop rotations. Doctoral dissertation, University of Georgia.
- Siddiqui, I. A., Sher, S. A., & French, A. M.** 1973. Distribution of plant parasitic nematodes in California. State of California Department of Food and Agriculture, Division of Plant Industry. 324 p.
- Silva, R. A. & Inomoto, N. M.** 2002. Host-range Characterization of Two *Pratylenchus coffeae* Isolates from Brazil. *Journal of Nematology*. 34:135-139.
- Simard, L., Bélair, G., & Miller, S.** 2009. First report of *Longidorus breviannulatus* associated with damage on creeping bentgrass golf greens in Québec, Canada. *Plant Disease*, 93, 846.
- Subbotin, S. A., Ragsdale, E. J., Mullens, T., Roberts, P. A., Mundo-Ocampo, M. & Baldwin, J. G.,** 2008. A phylogenetic framework for root lesion nematodes of the genus *Pratylenchus* (Nematoda): Evidence from 18S and D2–D3 expansion segments of 28S ribosomal RNA genes and morphological characters. *Molecular phylogenetics and evolution*, 48(2), pp.491-505.
- Subbotin, S. A., Vovlas, N., Yeates, G. W., Hallmann, J., Kiewnick, S., Chizhov, V. N., Manzanilla-López, R. H., Inserra, R. N., & Castillo, P.** 2015. Morphological and molecular characterization of *Helicotylenchus pseudorobustus* (Steiner, 1914) Golden, 1956 and related species (Tylenchida: Hoplolaimidae) with phylogeny of the genus. *Nematology*, 17, 27-52.
- Sutton, T. B., Aldwinckle, H. S., Agnello, A. M. & Walgenbach, J. F.** 2014. Compendium of Apple and Pear Diseases and Pests (Second Edition). American Phytopathological Society, St. Paul, MN, USA. 218 pp.
- Téliz, O. D., & Goheen, A. C.** 1968. Diseases of grapevines in Mexico. *Plant Disease Report*, 52, 372–373.
- Van den Berg, E., Tiedt, L. R. & Subbotin, S. A.** 2014. Morphological and molecular characterisation of several *Paratylenchus* Micoletzky, 1922 (Tylenchida: Paratylenchidae) species from South Africa and USA, together with some taxonomic notes. *Nematology*, 16(3), pp.323-358.
- Van den Berg, E., Tiedt, L. R., & Subbotin, S. A.** 2014. Morphological and molecular characterisation of several *Paratylenchus* Micoletzky, 1922 (Tylenchida: Paratylenchidae) species from South Africa and USA, together with some taxonomic notes. *Nematology* 16(3): 323-358.
- Van Driell, L., Potter, J. W., & Ebsary, B. A.** 1990. Distribution of virus vector nematodes associated with peach and other fruit crops in Essex County, Ontario. *Canadian Plant Disease Survey*, 70, 23–26.
- Villalobos, B. O.** 1980. Efecto residual de Nemaicur 10G y Furadan 10G sobre las poblaciones de nematodos y el rendimiento en dos variedades de vid en la P. P. La Enramada, Municipio de Matamoros, Coahuila. Tesis de Licenciatura. Universidad Juárez del Estado de Durango, Gómez Palacio, México, 78 p.
- Vrain, T. C., & Rousselle, G. L.** 1980. Distribution of plant parasitic nematodes in Quebec apple orchards. *Plant Disease*, 64, 582–583.
- Wang, H., Zhuo, K. & Liao, J.** 2016. Morphological and molecular characterization of *Pratylenchus hippeastri*, a new record of root-lesion nematode associated with apple in China. *Pakistan Journal of Zoology* 48: 665-671.
- Weimin, Y., Szalanski, A. & Robbins, R. T.** 2004. Phylogenetic relationships and genetic variation in *Longidorus* and *Xiphinema* species (Nematoda: Longidoridae) using ITS1 sequences of nuclear ribosomal DNA. *Journal of Nematology*. 36:14–19.
- Xu, Y.-M. & Zhao, Z.-Q.** 2019. Longidoridae and Trichodoridae (Nematoda: Dorylaimida and

Triplonchida). Landcare Research, Auckland, New Zealand. 149 pp.

- Ye, W., Szalanski, A. & Robbins, R. T.** 2004. Phylogenetic Relationships and Genetic Variation in *Longidorus* and *Xiphinema* Species (Nematoda: Longidoridae) Using ITS1 Sequences of Nuclear Ribosomal DNA. *Journal of Nematology* 36: 14-19.
- Zasada, I. A., Kitner, M., Wram, C., Wade, N., Ingham, R. E., Hafez, S., Mojtahedi, H., Chavoshi, S. & Hammack, N.** 2019. Trends in occurrence, distribution, and population densities of plant-parasitic nematodes in the Pacific Northwest of the United States from 2012-2016. *Plant Health Progress* 20:20-28.
- Zavaleta-Mejia, E. & Sosa-Moss, C.** 1979. Description of a new species of *Helicotylenchus* Steiner, 1945 (Nematoda: Hoplolaimida) and observations on three other spiral nematodes. *Nematropica* 9: 72-75.

Table 6: Arthropod pests of stone and pome fruit trees

Context: The pests named in this table are considered vectors, which promote the introduction (entry and establishment) and spread of the stone and pome fruit diseases listed in this standard.

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
Acari					
Trombidiformes: Eriophyidae					
<i>Aculus fockeui</i> (Nalepa & Trouessart)	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i>	Abou-Awad <i>et al.</i> , 2010; Alford, 2014; Isac <i>et al.</i> , 1998; Keifer, 1946	P2	P2	P2
<i>Eriophyes inaequalis</i> Wilson & Oldfield	<i>P. avium</i>	Beers <i>et al.</i> , 1993; Caprile <i>et al.</i> , 2016; Lindquist <i>et al.</i> , 1996; Vacante, 2016	P2	P2	Ab1
<i>Eriophyes insidiosus</i> Keifer & Wilson	<i>P. persica</i> , <i>P. var. nectarina</i> , <i>Prunus</i> spp.	Acuña-Soto & Estrada-Venegas, 2009; Acuña-Soto <i>et al.</i> , 2014; Gispert <i>et al.</i> , 1998; Lindquist <i>et al.</i> , 1996; Oldfield <i>et al.</i> , 1995	Ab1	P2	Ab3
Insecta					
Coleoptera: Nitidulidae					
<i>Carpophilus freemani</i> Dodson	Indiscriminate pest of <i>Prunus</i>	Connell, 1991 ; Connell <i>et al.</i> , 2005; Ewing & Cline 2005; Moller & DeVay, 1968; Rodríguez-del-bosque <i>et al.</i> , 1998; UF/IFAS 2018	Ab1	P2	Ab3
Diptera: Drosophilidae					
<i>Chymomyza procnemoides</i> Wheeler	<i>P. dulcis</i>	Connell <i>et al.</i> , 2005; Moller & DeVay, 1968; Wheeler, 1952	P2	P2	Ab1
Hemiptera: Aphididae					

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
<i>Aphis craccivora</i> Koch	<i>P. persica</i>	Alford, 2014; Blackman & Eastop, 2000; Labonne <i>et al.</i> , 1994	P2	P2	P2
<i>Aphis fabae</i> (Scopoli)	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i> , <i>Prunus</i> spp.	Blackman & Eastop, 2000; Gildow <i>et al.</i> , 2004; Smith & Parron, 1978; Wallis <i>et al.</i> , 2005	P1	P1	P2
<i>Aphis gossypii</i> Glover	<i>P. domestica</i> , <i>P. dulcis</i>	Blackman & Eastop, 2000; Gildow <i>et al.</i> , 2004; Hill, 1987; Labonne <i>et al.</i> , 1994; Smith & Parron, 1978; Wallis <i>et al.</i> , 2005	P1	P1	P2
<i>Aphis hederae</i> Kaltentbach	<i>P. dulcis</i>	Blackman & Eastop, 2000; Halima, 2012; Labonne <i>et al.</i> , 1994; Smith & Parron, 1978	P2	P2	P2
<i>Aphis spiraecola</i> Patch	<i>Prunus</i> spp., <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. domestica</i> , <i>P. persica</i>	Blackman & Eastop, 2000; Blackman, 2019; CABI, 2019; Gildow <i>et al.</i> , 2004; Labonne <i>et al.</i> , 1994; Wallis <i>et al.</i> , 2005	P2	P2	P2
<i>Brachycaudus cardui</i> (L.)	<i>Prunus</i> spp., <i>P. domestica</i>	Alford, 2014; Isac <i>et al.</i> , 1998; Blackman & Eastop, 2000; Kunze & Krczal, 1971; Smith & Parron, 1978	P2	P2	Ab1
<i>Brachycaudus helichrysi</i> (Kaltentbach)	<i>Prunus</i> spp., <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i>	Blackman & Eastop, 2000; Hadidi <i>et al.</i> , 2011; Isac <i>et al.</i> , 1998; Kunze & Krczal, 1971; Smith & Parron, 1978; Zsuzsa <i>et al.</i> , 1997	P2	P2	P2
<i>Brachycaudus persicae</i> (Passerini)	<i>Prunus</i> spp., <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i>	Blackman & Eastop, 2000; Gildow <i>et al.</i> , 2004; Leclant, 1973; Wallis <i>et al.</i> , 2005	P2	P2	P2
<i>Hyalopterus pruni</i> (Geoffroy)	<i>Prunus</i> spp., <i>P. domestica</i> , <i>P. armeniaca</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i>	Blackman & Eastop, 2000; Hadidi <i>et al.</i> , 2011; Hill, 1987; Isac <i>et al.</i> , 1998; Kunze & Krczal, 1971; Smith & Parron, 1978; Zsuzsa <i>et al.</i> , 1997	P2	P2	P2
<i>Myzus cerasi</i> (Fabricius)	<i>P. avium</i> , <i>P. persica</i>	Blackman & Eastop, 2000;	P1	P2	P2

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
		Smith & Parron, 1978; Zsuzsa <i>et al.</i> , 1997			
<i>Myzus persicae</i> (Sulzer)	<i>Prunus</i> spp., <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. var. nectarina</i>	Alford, 2014; Blackman & Eastop, 2000; Gildow <i>et al.</i> , 2004; Hill, 1987; Isac <i>et al.</i> , 1998; Kunze & Krczal, 1971; Labonne <i>et al.</i> , 1994; Leclant, 1973; Smith & Parron, 1978; Wallis <i>et al.</i> , 2005	P1	P1	P1
<i>Myzus varians</i> (Davidson)	<i>Prunus persica</i>	Manachini <i>et al.</i> , 2004; Németh, 1994	Ab1	P2	P2
<i>Phorodon humuli</i> (Schrank)	<i>P. domestica</i> , <i>Prunus</i> spp.	Alford, 2014; Blackman & Eastop, 2000; Hill, 1987; Isac <i>et al.</i> , 1998; Kunze & Krczal, 1971; Smith & Parron, 1978; Zsuzsa <i>et al.</i> , 1997	P2	P2	Ab1
<i>Rhopalosiphum padi</i> (L.)	<i>Prunus</i> spp.	Blackman & Eastop, 2000; Labonne <i>et al.</i> , 1994; Smith & Parron, 1978; Zsuzsa <i>et al.</i> , 1997	P1	P1	P2
<i>Toxoptera citricida</i> (Kirkaldy)	<i>P. domestica</i>	Blackman & Eastop, 2000; Halbert <i>et al.</i> , 2000; Gildow <i>et al.</i> , 2004	Ab1	P2	P2
Hemiptera: Cercopidae					
<i>Philaenus spumarius</i> L.	<i>P. dulcis</i> , <i>P. persica</i>	CABI, 2019; Cornara <i>et al.</i> , 2017; Ivanauskas <i>et al.</i> , 2014; Redak <i>et al.</i> , 2004; Wiegert, 1964	P1	P2	Ab1
Hemiptera: Cicadellidae					
<i>Colladonus clitellarius</i> (Say)	<i>P. persica</i> , <i>P. avium</i> , <i>Prunus</i> spp.	George & Davidson, 1959; Gilmer <i>et al.</i> , 1966; Metcalf, 1967; Taboada <i>et al.</i> , 1975	P2	P2	Ab1
<i>Colladonus geminatus</i> (Van Duzee)	<i>P. persica</i> , <i>P. avium</i>	Metcalf, 1967; Nielson, 1968; Wolfe <i>et al.</i> , 1950	P2	P2	Ab1

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
<i>Colladonus montanus</i> (Van Duzee)	<i>P. persica</i> in the laboratory	Caprile <i>et al.</i> , 2016; Diekmann & Putter, 1996; Kirkpatrick <i>et al.</i> , 1990; Metcalf, 1967; Nielson, 1968	P2	P2	Ab1
<i>Cuerna costalis</i> (F.)	<i>P. persica</i>	Janse, 2005; Janse & Obradovic, 2010; Metcalf, 1965; Nielson, 1968; Overall & Rebek, 2017; Purcell, 2008; Redak <i>et al.</i> , 2004; Turner & Pollard, 1955; Turner & Pollard, 1959a, b	P2	P2	Ab1
<i>Fieberiella florii</i> (Stal)	<i>P. avium</i> , <i>P. persica</i> , <i>P. var. nectarina</i> , <i>Malus domestica</i>	Alford, 2014; Anthon & Wolfe, 1951; Caprile <i>et al.</i> , 2016; Douglas 1999; Gilmer <i>et al.</i> , 1966; Jensen, 1957; Kirkpatrick <i>et al.</i> , 1990; Krczal <i>et al.</i> , 1989; Metcalf, 1967; Nielson, 1968	P2	P2	Ab1
<i>Graphocephala confluens</i> Uhler) = <i>Keonella confluens</i> (Uhler) – typo: <i>Keonolla confluens</i> , now <i>Neokolla confluens</i> (Uhler)	<i>Prunus</i> spp.	Anthon & Wolfe, 1951; Nielson, 1968	P2 (As <i>Neokolla confluens</i> (Uhler)	P2	Ab1
<i>Graphocephala versuta</i> (Say)	<i>P. persica</i> , <i>P. domestica</i> , <i>Prunus</i> spp.	Janse, 2005; Metcalf, 1965; Nielson, 1968; Purcell, 2008; Turner & Pollard, 1959a; Turner & Pollard, 1955; Wilson <i>et al.</i> , 2009	Ab1	P2	Ab3
<i>Gyponana lamina</i> DeLong = <i>Gyponana expanda</i> DeLong	<i>Prunus</i> spp.	Gilmer <i>et al.</i> , 1966; Hamilton, 1982; Metcalf, 1965; Nielson, 1968	P2 (as <i>G. expanda</i>)	P2	Ab1
<i>Homalodisca vitripennis</i> (Germar) syn: <i>H. coagulata</i> (Say)	<i>P. avium</i> , <i>P. domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>P. salicina</i> , <i>Prunus</i> spp.	CABI 2019; Janse, 2005; Janse & Obradovic, 2010; Martin <i>et al.</i> , 2012; Nielson, 1968; Overall & Rebek, 2017; Purcell, 2008; Redak <i>et al.</i> , 2008; Triapitsyn &	Ab1	P1	P2

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
		Phillips, 2000; Turner & Pollard, 1959a			
<i>Macropsis trimaculata</i> (Fitch)	<i>P. persica</i> , <i>P. armeniaca</i> , <i>P. domestica</i> , <i>Prunus</i> spp.	Layne & Bassi, 2008; Metcalf, 1966; Nielson, 1968	P2	P2	Ab1
<i>Norvellina seminuda</i> (Say)	<i>P. persica</i>	Gilmer <i>et al.</i> , 1966; Metcalf, 1967; Nielson 1968	P2	P2	Ab1
<i>Oncometopia orbona</i> (F.)	<i>P. persica</i> , <i>Prunus</i> spp.	Janse, 2005; Janse & Obradovic, 2010; Metcalf, 1965; Nielson, 1968; Overall & Rebek, 2017	P2	P2	Ab1
<i>Oncometopia nigricans</i> (Walker)	<i>P. persica</i>	Coronado-Blanco <i>et al.</i> , 2000; Layne & Bassi, 2008; Metcalf, 1965 ; Nielson, 1968; Purcell, 2008; Redak <i>et al.</i> , 2004	Ab1	P2	Ab3
<i>Osbornellus borealis</i> DeLong & Mohr.	<i>P. persica</i>	Jensen, 1957; Nielson, 1968	P2	P2	Ab1
<i>Paraphlepsius irroratus</i> (Say)	<i>P. persica</i>	Gilmer <i>et al.</i> , 1966; Layne & Bassi, 2008; López-Rosas <i>et al.</i> , 2009 ; Metcalf, 1967; Nielson, 1968; Rosenberger & Jones, 1978	P1	P2	Ab3
<i>Scaphytopius acutus</i> (Say) syn.: <i>Scaphytopius acutus delongi</i> Young	<i>P. persica</i> , <i>P. avium</i> , <i>Prunus</i> spp.	Anthon & Wolfe, 1951; GBIF, 2022; Gilmer <i>et al.</i> , 1966; Metcalf, 1967; Nielson, 1968; Rosenberger & Jones, 1978; Taboada <i>et al.</i> , 1975	P1	P2	Ab3
Hemiptera: Cixidae					
<i>Hyalesthes obsoletus</i> Signoret	<i>P. domestica</i> , <i>P. avium</i> , <i>Prunus</i> spp.	Alford, 2014; Bressan <i>et al.</i> , 2007; Sforza <i>et al.</i> , 1998	Ab1	Ab1	Ab1
Hemiptera: Pseudococcidae					
<i>Phenacoccus aceris</i> (Signoret)	<i>P. armeniaca</i> , <i>P. avium</i> , <i>P.</i>	García Morales <i>et al.</i> , 2016;	P2	P2	Ab1

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
	<i>domestica</i> , <i>P. dulcis</i> , <i>P. persica</i> , <i>Prunus</i> spp.	Raine <i>et al.</i> , 1986			
<i>Pseudococcus maritimus</i> (Ehrhorn)	<i>P. armeniaca</i> , <i>P. persica</i>	García Morales <i>et al.</i> , 2016; Hill, 1987; Long <i>et al.</i> , 2017; Mekuria <i>et al.</i> , 2013	P2	P2	P2
Hemiptera: Psyllidae					
<i>Cacopsylla melanoneura</i> (Förster)	<i>Malus</i> spp.	Jarausch & Jarausch, 2010	Ab1	Ab1	Ab1
<i>Cacopsylla picta</i> (Förster) syn. <i>C. costalis</i> (Flor)	<i>Malus</i> spp.	Hadidi <i>et al.</i> , 2011; Jarausch & Jarausch, 2010	Ab1	Ab1	Ab1
<i>Cacopsylla pruni</i> (Scopoli)	<i>P. dulcis</i> , <i>P. armeniaca</i> , <i>P. domestica</i> , <i>Prunus</i> spp.	Gallinger & Gross, 2018; Hadidi <i>et al.</i> , 2011; Layne & Bassi, 2008	Ab1	Ab1	Ab1
<i>Cacopsylla pyri</i> (Linnaeus)	<i>Pyrus</i> spp.	Hadidi <i>et al.</i> , 2011; Jarausch & Jarausch, 2010	Ab1	Ab1	Ab1
<i>Cacopsylla pyricola</i> (Förster)	<i>P. persica</i> , <i>Pyrus</i> spp.	Alford, 2014; Blomquist & Kirkpatrick; 2002; Hadidi <i>et al.</i> , 2011; Hodkimson, 1988; Jarausch & Jarausch, 2010; Layne & Bassi, 2008	P2	P2	Ab1
Lepidoptera: Pyralidae					
<i>Amyelois transitella</i> (Walker)	<i>P. domestica</i> , <i>P. armeniaca</i> , <i>P. persica</i> , <i>P. dulcis</i>	Ampt <i>et al.</i> , 2015; Heinrich, 1956 ; Palumbo <i>et al.</i> , 2014; Wenneker <i>et al.</i> , 2015	P6	P2	P2
<i>Euzophera semifuneralis</i> (Walker)	<i>P. avium</i> , <i>P. domestica</i>	Biddinger & Howitt, 1992 ; Connell <i>et al.</i> , 2005; Long <i>et al.</i> , 2017 Robinson <i>et al.</i> , 2002; Van Steenwyk <i>et al.</i> , 1986	P2	P2	Ab1
Thysanoptera: Thripidae					

VECTOR NAME	HOST	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
<i>Frankliniella occidentalis</i> (Pergande)	<i>P. avium</i> , <i>P. domestica</i> , <i>P. persica</i> , <i>P. persica</i> var. <i>nectarina</i>	CABI, 2019; Caprile <i>et al.</i> , 2016; Hadidi <i>et al.</i> , 2011; Hill, 1987; Long <i>et al.</i> , 2017; Milne & Walter, 2003; Saccaggi & Pieterse, 2013	P2	P2	P2
<i>Thrips tabaci</i> Lindeman	<i>P. persica</i>	Greber <i>et al.</i> , 1991; Hill, 1987; Hodges <i>et al.</i> , 2009; Milne & Walter, 2003; Pinent <i>et al.</i> , 2008	P1	P1	P2

References

- Abou-Awad, B. A., AL-Azzazy, M. M. and El-Sawi, S. A.** 2010. The life–history of the peach silver mite, *Aculus fockeui* (Acari: Eriophyidae) in Egypt. Archives of Phytopathology and Plant Protection 43(4):384-389.
- Acuña-Soto, J. A. and Estrada-Venegas, E. G.** 2009. Actualización al conocimiento de la superfamilia Eriophyoidae (Acari: Prostigmata) en México. Entomología Mexicana: 119-124.
- Acuña-Soto, J., Estrada-Venegas, E. G. and Equihua-Martínez, A.** 2014. Ácaros eriófidos (Acari: Prostigmata), en plantas no nativas de México. Entomología Mexicana, 1: 90-95.
- Alford, D. V.** 2014. Pests of fruit crops: a colour handbook, Second Edition. CRC Press, Boca Raton, FL. 461 pp.
- Ampt, E. A., Bush, D. S., Siegel, J. P. and Berenbaum, M. R.** 2015. Larval Preference and Performance of *Amyelois transitella* (Navel Orangeworm, Lepidoptera: Pyralidae) in Relation to the Fungus *Aspergillus flavus*. Environmental Entomology 45(1):155-162.
- Anthon, E. and Wolfe, H.** 1951. Additional insect vectors of western X-disease. Plant Dis Rep 35:245-46.
- Beers, E. H., Brunner, J., Willett, M. and Warner, G.** 1993. Orchard Pest Management: A Resource Book for the Pacific Northwest.
- Biddinger, D. J. & Howitt, A. J.** 1992. The food plants and distribution of the American plum borer (Lepidoptera: Pyralidae). The Great Lakes Entomologist 25(3): 149-159.
- Blackman, R. L.** n.d. Aphids on the world's plants, an online identification and information guide. [Online] Available: <http://www.aphidsonworldsplants.info/> (Accessed 5/3/2019, 2019).
- Blackman, R. L. and Eastop, V. F.** 2000. Aphids on the World's Crops: An Identification and Information Guide. 2nd ed. John Wiley & Sons, Ltd, Chichester, UK. 466 pp.
- Blomquist, C. and Kirkpatrick, B.** 2002. Identification of phytoplasma taxa and insect vectors of peach yellow leaf roll disease in California. Plant disease 86(7):759-763.
- Bressan, A., Turata, R., Maixner, M., Spiazzi, S., Boudon-Padieu, E. & Girolami, V.** 2007. Vector activity of *Hyalesthes obsoletus* living on nettles and transmitting a stolbur phytoplasma to grapevines: a case study. Annals of Applied Biology:331-339.
- CABI.** 2019. Crop Protection Compendium [CD]. Commonwealth Agricultural Bureau International, Wallingford, UK.
- Caprile, J. A., Symmes, E. J., Adaskaveg, J. E., Baldwin, R. A., Roncoroni, J. A., Westerdahl, B. B., Coates, W. W., Daane, K. M., Grant, J. A. and Van Steenwyk, R. A.** 2016. Cherry. Pest Management Guidelines for Agriculture. University of California.
- Connell, W. A.** 1991. Sap beetles (Nitidulidae: Coleoptera). Pages 151-174 *In* Insect and Mite Pests in Food: and Illustrated Key Vol. 1. United States Department of Agriculture, Agricultural Research Service, Washington D.C.
- Connell, J. H., Gubler, W. D. and Steenwyk, R. A.** 2005. Almond trunk injury treatment following bark damage during shaker harvest. Proc. XIII GREMPA Meeting on Almonds and Pistachios, Zaragoza, Spain.
- Cornara, D., Cavalieri, V., Dongiovanni, C., Altamura, G., Palmisano, F., Bosco, D., Porcelli, F., Almeida, R. and Saponari, M.** 2017. Transmission of *Xylella fastidiosa* by naturally infected *Philaenus spumarius* (Hemiptera, Aphrophoridae) to different host plants. Journal of Applied Entomology 141(1-2):80-87.
- Coronado-Blanco, J.M., Ruíz-Cancino, E., y Triapitsyn, S.V.** 2000. Chicharritas de la tribu Proconiini (Homoptera: Cicadellidae) asociadas a cítricos en Tamaulipas México. Acta Zoológica Mexicana, 81: 133-134.
- Diekmann, M. and Putter, C. A. J.** 1996. Stone fruits. Biodiversity International.

- Douglas, S.** 1999. Management of X-Disease in Stone Fruits. The Connecticut Agricultural Experiment Station Extension Publ PP062.
- Ewing, C. P. and Cline, A. R.** 2005. Key to Adventive Sap Beetles (Coleoptera: Nitidulidae) in Hawaii, with Notes on Records and Habits. The Coleopterists Bulletin 59(2):167-183.
- Gallinger, J. & Gross, J.** 2018. Unraveling the host plant alternation of *Cacopsylla pruni*—adults but not nymphs can survive on conifers due to phloem/xylem composition. *Frontiers in Plant Science* 9:484.
- García Morales, M., Denno, B., Miller, D. R., Miller, G. L., Ben-Dov, Y. and Hardy, N. B.** 2016. ScaleNet: A literature-based Model of Scale Insect Biology and Systematics.
- George, J. and Davidson, T.** 1959. Notes on life-history and rearing of *Colladonus clitellarius* (Say) (Homoptera: Cicadellidae). *The Canadian Entomologist* 91(6):376-379.
- Gildow, F., Damsteegt, V., Stone, A., Schneider, W., Luster, D. & Levy, L.** 2004. Plum pox in North America: identification of aphid vectors and a potential role for fruit in virus spread. *Phytopathology* 94(8):868-874.
- Gilmer, R.M., Palmiter, D.H., Schaefers, G.A., and McEwen, F.L.** 1966. Insect transmission of X-disease virus of stone fruits in New York. New York State Agricultural Experiment Station. Bulletin 813 (Geneva). 22pp.
- Gispert, C., Oldfield, G. N., Perring, T. M. & Creamer, R.** 1998. Biology of the transmission of peach mosaic virus by *Eriophyes insidiosus* (Acari: Eriophyidae). *Plant disease* 82(12):1371-1374.
- Greber, R., Klose, M., Milne, J. & Teakle, D.** 1991. Transmission of Prunus necrotic ringspot virus using plum pollen and thrips. *Annals of Applied Biology* 118(3):589-593.
- GBIF.** 2022. *Scaphytopius acutus* (Say, 1830). Global Biodiversity Information Facility (GBIF). Online at https://www.gbif.org/es/occurrence/search?taxon_key=2030616.
- Hadidi, A., Barba, M., Candresse, T. & Jelkmann, W.** 2011. Virus and virus-like diseases of pome and stone fruits. *Am. Phytopath. Society*.
- Halbert, S. E., Remaudière, G., & Webb, S. E.** 2000. Newly established and rarely collected aphids (Homoptera: Aphididae) in Florida and the southern United States. *Florida Entomologist* 83(1):79-91.
- Halima, M. K. B.** 2012. Aphid fauna (Hemiptera Aphididae) and their host association of chott mariem, coastal area of Tunisia. *Annals of Biological Research* 3(2):746-754.
- Hamilton, K. G. A.** 1982. Review of the Nearctic Species of the nominate subgenus of *Gyponana* Ball (Rhynchota: Homoptera: Cicadellidae). *Journal of the Kansas Entomological Society* 55(3):547-562.
- Heinrich, C.** 1956. American moths of the subfamily Phycitinae. *Bulletin of the United States National Museum*. 581 pp.
- Hill, D. S.** 1987. *Agricultural insect pests of temperate regions and their control*. Cambridge University Press, Cambridge. 659 pp.
- Hodges, A., Ludwig, S., Osborne, L., & Edwards, G. B.** 2009. *Pest Thrips of the United States: Field Identification Guide*. USDA-CSREES. 143 pp.
- Hodkinson, I. D.** 1988. The Nearctic Psylloidea (Insecta: Homoptera): an annotated check list. *Journal of Natural History* 22:1243
- Isac, M., Preda, S. & Marcu, M.** 1998. Aphid species--vectors of plum pox virus. *Acta virologica* 42(4):233-234.
- Ivanauskas, A., Valunas, D., Jomantiene, R., Picciau, L. & Davis, R. E.** 2014. Possible insect vectors of *Candidatus Phytoplasma asteris* and *Ca. Phytoplasma pruni*-related strains in Lithuania. *Zemdirbyste-Agriculture* 101(3).

- Jarausch, B. & Jarausch, W.** 2010. Psyllid Vectors and their Control. In P.G. Weintraub & P. Jones, eds. *Phytoplasmas - Genomes, Plant Hosts and Vectors*, pp 250 - 271. Cambridge, Massachusetts, CAB International.
- Janse, J. and Obradovic, A.** 2010. *Xylella fastidiosa*: its biology, diagnosis, control, and risks. *Journal of Plant Pathology*: S35-S48.
- Janse, J. D.** 2005. *Phytobacteriology Principles and Practice*. CABI International, Wallingford, UK. 360 pp.
- Jensen, D. D.** 1957. Transmission of Peach Yellow Leaf Roll Virus by *Fieberiella florii* (Stål) and a New Vector, *Osbornellus borealis* DeL. & M. *Journal of Economic Entomology* 50(5):668-672.
- Keifer, H. H.** 1946. A review of North American economic Eriophyid mites. *Journal of Economic Entomology* 39(5): 563-570.
- Kirkpatrick, B., Fisher, G., Fraser, J. & Purcell, A.** 1990. Epidemiological and phylogenetic studies on western X-disease mycoplasma-like organisms. Proc. Recent advances in mycoplasmaology. Proceedings of the 7th congress of the International Organization for Mycoplasmaology, Baden near Vienna, 1988.
- Krczal, G., Krczal, H., and Kunze L.** 1989. *Fieberiella florii* (Stäl), a vector of apple proliferation agent. *Acta Horticulturae* 235, 99-106.
- Kunze, L. & Krczal, H.** 1971. Transmission of sharka virus by aphids. Proc. Eur Symp Fruit Tree Virus Dis Proc. 255-260.
- Labonne, G., Yvon, M., Quiot, J., Avinent, L. & Llacer, G.** 1994. Aphids as potential vectors of plum pox virus: comparison of methods of testing and epidemiological consequences. *Acta Horticulturae* 386:207-218.
- Layne, D. R. & Bassi, D.,** (eds.) 2008. *The Peach: Botany, Production and Uses*. CAB International, Wallingford, UK.
- Leclant, F.** 1973. Ecological aspects of the transmission of Sharka disease (Plum pox) in south-eastern France. Discovery of new species of aphid vectors. Proc. Annales de Phytopathologie.
- Lindquist, E. E., Sabelis, M. W. and Bruin, J.,** (eds.) 1996. *Eriophyoid Mites: Their Biology, Natural Enemies, and Control*. Elsevier, Amsterdam.
- Long, L., Stoven, H. & Bush, M.** 2017. Tree Fruit Crops: Cherry (Sweet and Sour) Pests Pages 32-46 in Hollingsworth, C. S., ed. *Pacific Northwest Insect Management Handbook*. Oregon State University.
- López-Rosas, F. J., Rodríguez-Navarro, S. and Torreón-Sierra, R. A.** 2009. Artrópodos asociados al cultivo de hierbabuena (*Mentha spicata* L. var. tashkent) en el predio "Las animas", Tulyehualco, D.F. *Entomología Mexicana*, 1039-1046.
- Manachini, B., Lozzia, G. C., Casati, P., Bianco, P. A. & Longoni, C.** 2004. On the transmissibility of PPV by *Myzus varians* Davidson (Homoptera: Aphididae) to *Clematis vitalba* L. (Ranunculaceae). *IOBC/wprs Bulletin* 27(5):25-28.
- Martin, K. W., Hodges, A. C., Leppla, N. C.** 2012. ID tools: Citrus pests, glassy-winged sharpshooter. Idtools.org
- Mekuria, T., Smith, T., Beers, E., Watson, G. & Eastwell, K.** 2013. First report of transmission of Little cherry virus 2 to sweet cherry by *Pseudococcus maritimus* (Ehrhorn)(Hemiptera: Pseudococcidae). *Plant disease* 97(6):851-851.
- Milne, J. & Walter, G.** 2003. The coincidence of thrips and dispersed pollen in PNRSV-infected stonefruit orchards—a precondition for thrips-mediated transmission via infected pollen. *Annals of applied biology* 142(3):291-298.
- Metcalf, Z.** 1965. *General Catalogue of Homoptera*. Fasc6. Cicadelloidea. Part 1-3. Tettigellidae. USDA, ARS. 975 pp.

- Metcalf, Z. 1966.** General Catalogue of Homoptera. Fasc6. Cicadelloidea. Part 13. Macropsidae. USDA, ARS. 261 pp.
- Metcalf, Z. 1967.** General Catalogue of Homoptera. Fasc6. Cicadelloidea. Part 10. Euscelidae. USDA, ARS. 2695 pp.
- Moller, W. & DeVay, J. 1968.** Insect transmission of *Ceratocystis fimbriata* in deciduous fruit orchards. *Phytopathology* 58(11).
- Németh, M. 1994.** History and importance of plum pox in stone-fruit production. *EPPO Bulletin* 24(3):525-536.
- Nielson, M. W. 1968.** The leafhopper vectors of phytopathogenic viruses (Homoptera, Cicadellidae): taxonomy, biology, and virus transmission. United States Department of Agriculture, Agricultural Research Service.
- Oldfield, G. N., Creamer, R., Guispert, C., Osorio, F., Rodriguez, R. and Perring, T. M. 1995.** Incidence and distribution of peach mosaic and its vector, *Eriophyes insidiosus* (Acari: Eriophyidae) in Mexico. *Plant Disease*, 79(2): 186-189.
- Overall, L. M. & Rebek, E. J. 2017.** Insect vectors and current management strategies for diseases caused by *Xylella fastidiosa* in the southern United States. *Journal of Integrated Pest Management* 8(1):12.
- Palumbo, J. D., Mahoney, N. E., Light, D. M., Siegel, J., Puckett, R. D. & Michailides, T. J. 2014.** Spread of *Aspergillus flavus* by navel orangeworm (*Amyelois transitella*) on almond. *Plant disease* 98(9):1194-1199.
- Pinent, S. M., Mascaro, F., Botton, M. & Redaelli, L. R. 2008.** Thrips (Thysanoptera: Thripidae, Phlaeothripidae) damaging peach in Paranapanema, São Paulo State, Brazil. *Neotropical Entomology* 37(4):486-488.
- Purcell, A. H. 2008.** Transmission of *Xylella fastidiosa* bacteria by xylem-feeding insects. *Encyclopedia of Entomology*:3885-3886.
- Raine, J., McMullen, R. & Forbes, A. 1986.** Transmission of the agent causing little cherry disease by the apple mealybug *Phenacoccus aceris* and the dodder *Cuscuta lupuliformis*. *Canadian Journal of Plant Pathology* 8(1):6-11.
- Redak, R. A., Purcell, A. H., Lopes, J. R., Blua, M. J., Mizell Iii, R. F. & Andersen, P. C. 2004.** The biology of xylem fluid-feeding insect vectors of *Xylella fastidiosa* and their relation to disease epidemiology. *Annual Reviews in Entomology* 49(1):243-270.
- Robinson, G. S., Ackery, P. R., Kitching, I. J., Beccaloni, G. W. & Hernández, L. M. 2002.** Hostplants of the Moth and Butterfly Caterpillars of America North of Mexico. American Entomological Institute, Gainesville, FL. 824 pp.
- Rodríguez-del-bosque, L.A., Leos-Martínez, J. & Dowd, P. F. 1998.** Effect of ear wounding and cultural practices on abundance of *Carpophilus freemani* (Coleoptera: Nitidulidae) and other micro coleopterans in maize in Northeastern Mexico. *Journal of Economic Entomology*, 91(4): 796-801.
- Rosenberger, D. A. & Jones, A. L. 1978.** Leafhopper vectors of the peach X disease pathogen and its seasonal transmission from chokeberry. *Phytopathology* 68:782-790.
- Saccaggi, D. L. & Pieterse, W. 2013.** Intercepting Aliens: Insects and Mites on Budwood Imported to South Africa. *Journal of Economic Entomology* 106(3):1179-1189.
- Sforza, R., Clair, D., Daire, X., Larrue, J. & Boudon-Padieu, E. 1998.** The role of *Hyalesthes obsoletus* (Hemiptera: Cixiidae) in the occurrence of bois noir of grapevines in France. *Journal of Phytopathology* 146(11-12):549-556.
- Simith, C. F., & Parron, C. S. 1978.** An annotated list of Aphididae (Homoptera) of North America. Technical Bulletin 255. North Carolina Agricultural Experimental Station, North Carolina State University, Raleigh NC. 428 pp.

- Taboada, O., Rosenberger, D. A. & Jones, A. L.** 1975. Leafhopper fauna of X-diseased peach and cherry orchards in southwest Michigan. *Journal of Economic Entomology* 68(2):255-257.
- Triapitsyn, S. V. & Phillips, P. A.** 2000. First record of *Gonatocerus triguttatus* (Hymenoptera: Mymaridae) from eggs of *Homalodisca coagulata* (Homoptera: Cicadellidae) with notes on the distribution of the host. *Florida Entomologist* 83(2):200.
- Turner, W. F. & Pollard, H. N.** 1955. Additional leafhopper vectors of phony peach. *Journal of Economic Entomology* 48(6):771-72.
- Turner, W. F. & Pollard, H. N.** 1959a. Insect transmission of phony peach disease. US Dept. of Agriculture.
- Turner, W. F. & Pollard, H. N.** 1959b. Life histories and behavior of five insect vectors of phony peach disease. US Dept. of Agriculture.
- UF/IFAS.** 2018. Featured Creatures. University of Florida Institute of Food and Agricultural Science (UF/IFAS), Gainesville, FL.
- Vacante, V.** 2016. The Family Eriophyidae Nalepa (Chapter 16). Pages 214-390 *In* The handbook of Mites of Economic Plants: Identification, Bio-ecology and Control. CAB International, Wallingford U.K.
- Van Steenwyk, R., Hendricks, L., Barclay, L. & Younce, E.** 1986. Borer control in young almond trees. *California Agriculture* 40(3):10-11.
- Wallis, C. M., Fleischer, S. J., Luster, D. & Gildow, F. E.** 2005. Aphid (Hemiptera: Aphididae) species composition and potential aphid vectors of plum pox virus in Pennsylvania peach orchards. *Journal of Economic Entomology* 98(5):1441-1450.
- Wheeler, M. R.** 1952. The Drosophilidae of the Nearctic region, exclusive of the genus *Drosophila*. University of Texas Publication 5204: 162-218.
- Wiegert, R. G.** 1964. Population energetics of meadow spittlebugs (*Philaenus spumarius* L.) as affected by migration and habitat. *Ecological Monographs* 34(3):217-241.
- Wilson M. R., Turner, J. A. and Mckamey, S. H.** 2009. Sharpshooter leafhoppers of the word (Hemiptera: Cicadellidae subfamily Cicadellinae). Amguedfa Cymru-National Museum Wales. Online: <https://naturalhistory.museumwales.ac.uk/Sharpshooters/>
- Wenneker, M., Duffy, B. & Montesinos, E.** 2015. DROPSA: Strategies to develop effective, innovative, and practical approaches to protect major European fruit crops from pests and pathogens. Proc. 2nd International Workshop on bacterial diseases of Stone fruits and nuts.
- Wolfe, H., Anthon, E. & Jones, L.** 1950. Transmission of western X-disease of Peaches by the leafhopper, *Colladonus geminatus* (Van D.). *Phytopathology* 40(10): 971.
- Zsuzsa, B., Dalma, P. & Richard, G.** 1997. Flight and transmission activity of PPV vector aphids. *Journal of Aphidology* 11:21-27.

Annex 2: Grapevine Pests

LEGEND FOR SYMBOLS USED IN TABLES

Presence or absence unless otherwise noted conform to the categories listed in ISPM 8: 2021 (*Determination of pest status in an area*). For ease of reference alphanumeric designations have been added here.

Ab1: Absent: pest not recorded

Ab2: Absent: the entire country is pest free

Ab3: Absent: pest records invalid

Ab4: Absent: pest no longer present

Ab5: Absent: pest eradicated

P1: Present: widely distributed

P2: Present: not widely distributed and not under official control

P3: Present: not widely distributed and under official control

P4: Present: at low prevalence

P5: Present: except in specified pest free areas

P6: Present: transient

P7: Present: not associated with host crop (NAPPO category)

Table 1: Virus pests of grapevine

Table 2: Fungal pathogens (incl. Chromista) of grapevine

Table 3. Bacterial pathogens of grapevine

Table 4: Nematode pests of grapevine

Table 5: Arthropod pests of grapevine

Table 1: Virus pests of grapevine

Viroid infections are ubiquitous throughout all grapevine growing regions of the world. All known viroids in grapevines are not known to cause any agronomic significant effects. Therefore, the viroids that are known to occur in the NAPPO region will not be included in this standard. Last updated December 2021.

PEST	ABBREVIATION	FAMILY	GENUS	REFERENCES	PRESENCE/ABSENCE		
					CAN	USA	MEX
Alfalfa mosaic virus	AMV	<i>Bromoviridae</i>	<i>Alfavirus</i>	Meng <i>et al.</i> , 2017	P7	P2	P2
Arabis mosaic virus	ArMV	<i>Secoviridae</i>	<i>Nepovirus</i>	MacKenzie <i>et al.</i> , 1996; Meng <i>et al.</i> , 2017	P2	P2	Ab3
Artichoke Italian latent virus	AILV	<i>Secoviridae</i>	<i>Nepovirus</i>	Jankulova <i>et al.</i> , 1976; Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Bean common mosaic virus	BCMV-PSt; peanut strain	<i>Potyviridae</i>	<i>Potyvirus</i>	Meng <i>et al.</i> , 2017	Ab1	P2	Ab3
Blackberry virus S	BVS	<i>Tymoviridae</i>	<i>Marafivirus</i>	Meng <i>et al.</i> , 2017	Ab1	P2	Ab1
Blueberry leaf mottle virus	BBLMV	<i>Secoviridae</i>	<i>Nepovirus</i>	Meng <i>et al.</i> , 2017	P7	P2	Ab1
Broad bean wilt virus	BBWV	<i>Secoviridae</i>	<i>Fabavirus</i>	Basso <i>et al.</i> , 2017; Castrovilli <i>et al.</i> , 1985; Pearson, 1988	P7	P2	Ab1
Carnation mottle virus	CarMV	<i>Tombusviridae</i>	<i>Alphacarmovirus</i>	Basso <i>et al.</i> , 2017; Wilcox, 2015	P7	P2	P2
Cherry leaf roll virus	CLRV	<i>Secoviridae</i>	<i>Nepovirus</i>	Basso <i>et al.</i> , 2017	P7	P2	Ab1
Cucumber mosaic virus	CMV	<i>Bromoviridae</i>	<i>Cucumovirus</i>	Basso <i>et al.</i> , 2017; Bovey, 1985	P7	P2	P2
Grapevine Ajinashika virus	GAgV	Unclassified	Unassigned	Meng <i>et al.</i> , 2017; Wilcox, 2015	Ab1	Ab1	Ab1
Grapevine Algerian latent virus	GALV	<i>Tombusviridae</i>	<i>Tombusvirus</i>	Basso <i>et al.</i> , 2017; Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine Anatolian ringspot virus	GARSV	<i>Secoviridae</i>	<i>Nepovirus</i>	Basso <i>et al.</i> , 2017; Hajizadeh <i>et al.</i> , 2012; Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine angular mosaic virus	GAMoV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	Basso <i>et al.</i> , 2017; Girgis, <i>et al.</i> , 2009; Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine asteroid mosaic-associated virus	GAMaV	<i>Tymoviridae</i>	<i>Marafivirus</i>	Martelli, 2014; Xiao and Meng, 2016	P2	P2	Ab1
Grapevine berry	GINV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	Giampetruzzi <i>et al.</i> , 2012;	Ab1	Ab1	Ab1

PEST	ABBREVIATION	FAMILY	GENUS	REFERENCES	PRESENCE/ABSENCE		
					CAN	USA	MEX
Inner necrosis virus				Martelli, 2014; Meng <i>et al.</i> , 2017			
Grapevine Bulgarian latent virus	GBLV	<i>Secoviridae</i>	<i>Nepovirus</i>	Meng <i>et al.</i> , 2017; Uyemoto <i>et al.</i> , 1977	Ab1	Ab1	Ab1
Grapevine chrome mosaic virus	GCMV	<i>Secoviridae</i>	<i>Nepovirus</i>	Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine deformation virus	GDefV	<i>Secoviridae</i>	<i>Nepovirus</i>	Hajizadeh <i>et al.</i> , 2012; Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine enamovirus 1	GEV-1	<i>Luteoviridae</i>	<i>Enamovirus</i>	Silva <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine fabavirus		<i>Secoviridae</i>	Fabavirus	Al Rwahnih <i>et al.</i> , 2016b	Ab1	Ab1	Ab1
Grapevine fanleaf virus	GFLV	<i>Secoviridae</i>	<i>Nepovirus</i>	CISEH, 2018a; MacKenzie <i>et al.</i> , 1996; Meng <i>et al.</i> , 2017; Wilcox, 2015	P2	P2	Ab1
Grapevine fleck virus	GFKV	<i>Tymoviridae</i>	<i>Maculavirus</i>	CISEH, 2018b; Kanuya, <i>et al.</i> , 2012; Meng <i>et al.</i> , 2017; Mikus and Goodman, 1999; Naidu and Mekuria, 2010; Poojari <i>et al.</i> , 2016	P2	P2	Ab1
Grapevine geminivirus A	GGVA	<i>Geminiviridae</i>	Unassigned	Al Rwahnih <i>et al.</i> , 2016a	Ab1	P2	Ab1
Grapevine labile rod-shaped virus	GLRSV	Unclassified	Unassigned	Basso <i>et al.</i> , 2017; Faggioli <i>et al.</i> , 1992	Ab1	Ab1	Ab1
Grapevine leafroll-associated virus 1	GLRaV-1	<i>Closteroviridae</i>	<i>Ampelovirus</i>	Borges <i>et al.</i> , 2020; Fuchs <i>et al.</i> , 2009; Meng <i>et al.</i> , 2017; MacKenzie <i>et al.</i> , 1996; Martin <i>et al.</i> , 2005; Sharma <i>et al.</i> , 2011	P2	P2	Ab1
Grapevine leafroll-associated virus 2	GLRaV-2	<i>Closteroviridae</i>	<i>Closterovirus</i>	Meng <i>et al.</i> , 2017; Fuchs <i>et al.</i> , 2009; Martin <i>et al.</i> , 2005; Borges <i>et al.</i> , 2020; Sharma <i>et al.</i> , 2011	P2	P2	Ab1
Grapevine leafroll-associated virus 3	GLRaV-3	<i>Closteroviridae</i>	<i>Ampelovirus</i>	Borges <i>et al.</i> , 2020; Fuchs <i>et al.</i> , 2009; Hoffman <i>et al.</i> , 2020; MacKenzie <i>et al.</i> , 1996; Martin <i>et al.</i> , 2005; Meng <i>et al.</i> , 2017; Mikus and Goodman,	P2	P2	Ab1

PEST	ABBREVIATION	FAMILY	GENUS	REFERENCES	PRESENCE/ABSENCE		
					CAN	USA	MEX
				1999; Sharma <i>et al.</i> , 2011			
Grapevine leafroll-associated virus 4	GLRaV-4	<i>Closteroviridae</i>	<i>Ampelovirus</i>	CISEH, 2018c; Sharma <i>et al.</i> , 2011; Wilcox, 2015	P2	P2	Ab1
Grapevine leafroll-associated virus 7	GLRaV-7	<i>Closteroviridae</i>	<i>Velarivirus</i>	Al Rwahnih <i>et al.</i> , 2012a; Morales and Monis, 2007; Wilcox, 2015	Ab1	P2	Ab1
Grapevine leafroll-associated virus 13	GLRaV-13	<i>Closteroviridae</i>	<i>Ampelovirus</i>	Ito and Nakaune, 2016	Ab1	Ab1	Ab1
Grapevine line pattern virus	GLPV	<i>Bromoviridae</i>	<i>Ilarvirus</i>	Wilcox, 2015	Ab1	Ab1	Ab1
Grapevine Pinot Gris virus	GPGV	<i>Betaflexiviridae</i>	<i>Trichovirus</i>	Al Rwahnih, 2016c; Giampetruzzi <i>et al.</i> , 2012	P2	P2	Ab1
Grapevine red blotch virus	GRBV	<i>Geminiviridae</i>	<i>Grabovirus</i>	Al Rwahnih <i>et al.</i> , 2013; Wilcox, 2015	P2	P2	P2
Grapevine red globe virus	GRGV	<i>Tymoviridae</i>	<i>Maculavirus</i>	Wilcox, 2015	Ab1	Ab3	Ab1
Grapevine roditis leaf discoloration-associated virus	GRLDaV	<i>Caulimoviridae</i>	<i>Badnavirus</i>	Maliogka <i>et al.</i> , 2015	Ab1	Ab1	Ab1
Grapevine rupestris vein feathering virus	GRVfV	<i>Tymoviridae</i>	<i>Marafivirus</i>	Chingandu <i>et al.</i> , 2020; Giampetruzzi <i>et al.</i> , 2012; Xiao and Meng, 2016	P2	P2	Ab1
Grapevine rupestris stem pitting-associated virus	RSPaV	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	Al Rwahnih <i>et al.</i> , 2009; CISEH, 2018d; Martin <i>et al.</i> , 2005; Meng <i>et al.</i> , 2017	P2	P2	Ab1
Grapevine Tunisian ringspot virus	GTRV	<i>Secoviridae</i>	<i>Nepovirus</i>	Basso <i>et al.</i> , 2017; Quertani <i>et al.</i> , 1992	Ab1	Ab1	Ab1
Grapevine stunt virus	GSV	Unclassified	Unassigned	Meng <i>et al.</i> , 2017	Ab1	Ab1	Ab1
Grapevine vein-clearing virus	GVCV	<i>Caulimoviridae</i>	<i>Badnavirus</i>	Jones <i>et al.</i> , 2015; Zhang <i>et al.</i> , 2011	Ab1	P2	Ab1
Grapevine virus A	GVA	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	CISEH, 2018e; Jones <i>et al.</i> , 2015	P2	P2	Ab1
Grapevine virus B	GVB	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Basso <i>et al.</i> , 2017; CISEH, 2018f	P2	P2	Ab1
Grapevine virus D	GVD	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Abou-Ghanem <i>et al.</i> , 1997; Basso <i>et al.</i> , 2017; CISEH, 2018g	Ab1	Ab3	Ab1

PEST	ABBREVIATION	FAMILY	GENUS	REFERENCES	PRESENCE/ABSENCE		
					CAN	USA	MEX
Grapevine virus E	GVE	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Alabi <i>et al.</i> , 2013; Basso <i>et al.</i> , 2017;	Ab1	P2	Ab1
Grapevine virus F	GVF	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Al Rwahnih <i>et al.</i> , 2012b	Ab1	P2	Ab1
Grapevine virus G	GVG	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Blouin <i>et al.</i> , 2018b; Diaz-Lara <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Grapevine virus H	GVH	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Candresse <i>et al.</i> , 2018; Diaz-Lara <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Grapevine virus I	GVI	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Blouin <i>et al.</i> , 2018a; Diaz-Lara <i>et al.</i> , 2018; Diaz-Lara <i>et al.</i> , 2019	Ab1	Ab1	Ab1
Grapevine virus J	GVJ	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Diaz-Lara <i>et al.</i> , 2018	Ab1	Ab1	Ab1
Grapevine virus L	GVL	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Debat <i>et al.</i> , 2019	Ab1	P2	Ab1
Grapevine virus M	GVM	<i>Betaflexiviridae</i>	<i>Vitivirus</i>	Alabi <i>et al.</i> , 2019	Ab1	P2	Ab1
Grapevine virus T	GVT	<i>Betaflexiviridae</i>	<i>Foveavirus</i>	Glasa <i>et al.</i> , 2018	Ab1	Ab1	Ab1
Grapevine Syrah virus-1	GSyV-1	<i>Tymoviridae</i>	<i>Marafivirus</i>	Al Rwahnih <i>et al.</i> , 2009	P2	P2	Ab1
Peach rosette mosaic virus	PRMV	<i>Secoviridae</i>	<i>Nepovirus</i>	Meng <i>et al.</i> , 2017	P7	P2	Ab1
Petunia asteroid mosaic virus	PAMV	<i>Tombusviridae</i>	<i>Tombusvirus</i>	Basso <i>et al.</i> , 2017	P7	Ab1	Ab1
Potato virus X	PVX	<i>Alphaflexiviridae</i>	<i>Potexvirus</i>	Wilcox, 2015	P7	P2	Ab3
Raspberry bushy dwarf virus	RBDV	Unclassified	<i>Ideovirus</i>	Wilcox, 2015	P7	P2	Ab1
Raspberry ringspot virus	RpRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	Martelli, 2014; Wilcox, 2015	Ab1	Ab1	Ab1
Sowbane mosaic virus	SoMV	<i>Solemoviridae</i>	<i>Sobemovirus</i>	Cesati and Van Regenmortel, 1969; Wilcox, 2015	P7	P2	Ab1
Strawberry latent ringspot virus	SLRSV	<i>Secoviridae</i>	Unassigned	Meng <i>et al.</i> , 2017	P7	P2	Ab3
Summer grape latent virus, syn. Grapevine Cabernet Sauvignon reovirus	SGLV; GCSV	<i>Reoviridae</i>	Unassigned	Meng <i>et al.</i> , 2017	Ab1	P2	Ab1
Tobacco mosaic virus	TMV	<i>Virgaviridae</i>	<i>Tobamovirus</i>	Wilcox, 2015	P7	P2	P2
Tobacco necrosis virus D	TNV-D	<i>Tombusviridae</i>	<i>Betanecrovirus</i>	Basso <i>et al.</i> , 2017	P7	Ab1	Ab1
Tobacco ringspot	TRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	Borges <i>et al.</i> , 2020;	P7	P2	P2

PEST	ABBREVIATION	FAMILY	GENUS	REFERENCES	PRESENCE/ABSENCE		
					CAN	USA	MEX
virus				Meng <i>et al.</i> , 2017			
Tomato black ring virus	TBRV	<i>Secoviridae</i>	<i>Nepovirus</i>	Meng <i>et al.</i> , 2017; Wilcox, 2015	P7	Ab1	Ab1
Tomato mosaic virus	ToMV	<i>Virgaviridae</i>	<i>Tobamovirus</i>	Basso <i>et al.</i> , 2017	P7	P2	Ab3
Tomato ringspot virus	ToRSV	<i>Secoviridae</i>	<i>Nepovirus</i>	Borges <i>et al.</i> , 2020; Meng <i>et al.</i> , 2017; Wilcox, 2015;	P2	P2	Ab3

References

- Abou-Ghanem, N., Saldarelli, P., Minafra, A., Buzkan, N., Castellano, M. A. & Martelli, G. P.** 1997. Properties of grapevine virus D, a novel putative trichovirus. *Journal of Plant Pathology*, 79(1):15-25.
- Al Rwahnih, M., Alabi, O. J., Westrick, N. M., Golino, D. & Rowhani, A.** 2016a. Description of a novel monopartite geminivirus and its defective subviral genome in grapevine. *Phytopathology*, 107(2):240-251.
- Al Rwahnih, M., Alabi, O.J., Westrick, N. M., Golino, D. & Rowhani, A.** 2016b. Near-complete genome sequence of grapevine fabavirus, a novel putative member of the genus *Fabavirus*. *Genome Announcements*, 4(4):1.
- Al Rwahnih, M., Daubert, S., Golino, D. & Rowhani, A.** 2009. Deep sequencing analysis of RNAs from a grapevine showing syrah decline symptoms reveals a multiple virus infection that includes a novel virus. *Virology*, 387(2):395-401.
- Al Rwahnih, M., Dave, A., Anderson, M. M., Rowhani, A., Uyemoto, J. K. & Sudarshana, M. R.** 2013. Association of a DNA virus with grapevines affected by red blotch disease in California. *Phytopathology*, 103(10):1069-1076.
- Al Rwahnih, M., Dolja, V. V., Daubert, S., Koonin, E. V. & Rowhani, A.** 2012a. Genomic and biological analysis of grapevine leafroll-associated virus 7 reveals a possible new genus within the family Closteroviridae. *Virus Research*, 163(1): 302-309.
- Al Rwahnih, M., Golino, D. & Rowhani, A.** 2016c. First report of grapevine pinot gris virus infecting grapevine in the United States. *Plant Disease*, 100(5):1030-1030.
- Al Rwahnih, M., Sudarshana, M. R., Uyemoto, J. K. & Rowhani, A.** 2012b. Complete genome sequence of a novel vitivirus isolated from grapevine. *Journal of Virology*, DOI: 10.1128/JVI.01444-12
- Alabi, O. J., McBride, S., Appel, D. N, Al Rwahnih, M. & Pontasch, F. N.** 2019. Grapevine virus M, a novel vitivirus discovered in the American hybrid bunch grape cultivar Blanc du Bois in Texas. *Archives of Virology*, 164:1739-1741.
- Alabi, O. J., Poojari, S., Sarver, K., Martin, R. R. & Naidu, R. A.** 2013. Complete genome sequence analysis of an american isolate of grapevine virus E. *Virus Genes*, 46(3):563-566.
- Basso, M. F., Fajardo, T. V. M. & Saldarelli, P.** 2017. Grapevine virus diseases: economic impact and current advances in viral prospection and management. *Revista Brasileira de Fruticultura*, 39(1):1-22.
- Blouin, A. G., Chooi, K. M., Warren, B., Napier, K. R., Barrero, R. A. & MacDiarmid, R. M.** 2018a. Grapevine virus I, a putative new vitivirus detected in co-infection with grapevine virus G in New Zealand. *Archives of Virology*, 163(5):1371-1374.
- Blouin, A. G., Keenan, S., Napier, K. R., Barrero, R. A. & MacDiarmid, R. M.** 2018b. Identification of a novel vitivirus from grapevines in New Zealand. *Archives of Virology*, 163(1):281-284.
- Borges, D. F., Preising, S., de Queiroz Ambrosio, M. M. & da Silva, W. L.** 2020. Detection of multiple grapevine viruses in New England vineyards. *Crop Protection*, 132:105143.
- Bovey, R.** 1985. Novel viruses & virus diseases of grapevine, new data on known diseases and their agents. *Phytopathologia Mediterranea*, 24(1/2):8-11.
- Candresse, T., Theil, S., Faure, C. I. & Marais, A.** 2018. Determination of the complete genomic sequence of grapevine virus H, a novel vitivirus infecting grapevine. *Archives of Virology*, 163(1):277-280.
- Castrovilli, S., Savino, V., Castellano, M. A. & Engelbrecht, D. J.** 1985. Characterization of a grapevine isolate of broad bean wilt virus. *Phytopathologia Mediterranea*, 24(1/2):35-40.
- Cesati, R. R., & Van Regenmortel, M. H. V.** 1969. Serological detection of a strain of tobacco necrosis virus in grapevine leaves. *Phytopathologische Zeitschrift*, 64(4):362-366.
- Chingandu, N., Jarugula, S., Adiputra, J., Bagewadi, B., Adegbola, R., Thammina, C. & Naidu, R.** 2020. First report of Grapevine rupestris vein feathering virus in grapevines from Washington State. *Plant Disease*. DOI: 10.1094/pdis-08-20-1662-pdn.
- CISEH.** 2018a. Widely Prevalent Viruses of the United States: Grapevine fanleaf virus (GFLV).

Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=20559>.

- CISEH.** 2018b. Widely Prevalent Viruses of the United States: Grapevine fleck virus (GFkV). Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=56425>.
- CISEH.** 2018c. Widely Prevalent Viruses of the United States: Grapevine Leafroll-associated virus 4 (GLRaV-4). Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=58464>.
- CISEH.** 2018d. Widely Prevalent Viruses of the United States: Grapevine rupestris stem pitting-associated virus. Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=56426>.
- CISEH.** 2018e. Widely Prevalent Viruses of the United States: Grapevine virus A (GVA). Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=56415>.
- CISEH.** 2018f. Widely Prevalent Viruses of the United States: Grapevine virus B (GVB). Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=56416>.
- CISEH.** 2018g. Widely Prevalent Viruses of the United States: Grapevine virus D (GVD). Center for Invasive Species and Ecosystem Health.
<https://www.prevalentviruses.org/subject.cfm?id=56417>.
- Debat, H., Zavallo, D., Brisbane, R. S., Voncina, D., Almeida, R. P. P., Blouin, A. G., Al Rwahnih, M., Gomez-Talquenca, W. & Asurmendi, S.** 2019. Grapevine virus L: a novel vitivirus in grapevine. *European Journal of Plant Pathology*, 155:319–328.
- Diaz-Lara, A., Brisbane, R.S., Aram, K., Golino, D. & Al Rwahnih, M.** 2019. Detection of new vitiviruses infecting grapevine in California. *Archives of Virology*, 164:2573–2580.
- Diaz-Lara, A., Golino, D. & Al Rwahnih, M.** 2018. Genomic characterization of grapevine virus J, a novel virus identified in grapevine. *Archives of Virology*, 163(7):1965-1967.
- Faggioli, F., Riccioni, L., Mazzei, M. & Barba, M.** 1992. Purification and characterization of a new virus found in grapevine. *Phytopathologia Mediterranea*, 31(1):37-40.
- Fuchs, M., Martinson, T., Loeb, G. & Hoch, H.** 2009. Survey for the three major leafroll disease-associated viruses in Finger Lakes vineyards in New York. *Plant Disease*, 93(4):395-401.
- Giampetruzzi, A., Roumi, V., Roberto, R., Malossini, U., Yoshikawa, N., La Notte, P., Terlizzi, F., Credi, R. & Saldarelli, P.** 2012. A new grapevine virus discovered by deep sequencing of virus-and viroid-derived small RNAs in Cv Pinot gris. *Virus Research*, 163(1):262-268.
- Girgis, S. M., Bem, F. P., Dovas, C. I., Sclavounos, A., Avgelis, A. D., Tsagris, M., Katis, N. & Kyriakopoulou, P. E.** 2009. Characterisation of a novel ilarvirus causing grapevine angular mosaic disease. *European Journal of Plant Pathology*, 125(2):203-211.
- Glasa, M., Predajňa, L., Sihelská, N., Šoltys, K., Ruiz-García, A. B., Olmos, A., Wetzels, T. & Sabanadzovic, S.** 2018. Grapevine virus T is relatively widespread in Slovakia and Czech Republic and genetically diverse. *Virus Genes*, 54(5):737-741.
- Hajizadeh, M., Sokhandan-Bashir, N. & Elbeaino, T.** 2012. First report of grapevine deformation virus and grapevine Anatolian ring spot virus in Iran. *Journal of Plant Pathology*, 94(4).
- Hoffmann, M., Talton, W., Nita, M., Jones, T., Al Rwahnih, M., Sudarshana, M. & Almeyda, C.** 2020. First report of grapevine leafroll-associated virus 3 in *Vitis vinifera* in North Carolina. *Journal of Plant Pathology*, 03(1):385-386.
- Ito, T. & Nakaune, R.** 2016. Molecular characterization of a novel putative ampelovirus tentatively named grapevine leafroll-associated virus 13. *Archives of Virology*, 161(9):2555-2559.
- Jankulova, M., Savino, V., Gallitelli, D., Quacquarelli, A. & Martelli, G. P.** 1976. Isolation of artichoke Italian latent virus from the grapevine in Bulgaria. In Proceedings of the 6th meeting of the International Committee on the Taxonomy of Viruses. Cordoba, Spain. Monografias INIA. 18. 143-148.
- Jones, T. J., Rayapati, N. A. & Nita, M.** 2015. Occurrence of grapevine leafroll associated virus-2,-3 and grapevine fleck virus in Virginia, USA, and factors affecting virus infected vines.

European Journal of Plant Pathology, 142(2):209-222.

- Kanuya, E., Clayton, I. A., Naidu, R. A. & Karasev, A. V.** 2012. First report of grapevine fleck virus in Idaho grapevines. *Plant Disease*, 96(11):1705-1705.
- MacKenzie, D. J., Johnson, R. C. & Warner, C.** 1996. Incidence of four important viral pathogens in Canadian vineyards. *Plant Disease*, 80(8):955-958.
- Maliogka, V.I., Olmos, A., Pappi, P.G., Lotos, L., Efthimiou, K., Grammatikaki, G., Candresse, T., Katis, N.I. & Avgelis, A. D.** 2015. A novel grapevine badnavirus is associated with the roditis leaf discoloration disease. *Virus Research*, 203: 47-55.
- Martelli, G. P.** 2014. Directory of virus and virus-like diseases of grapevine and their agents. *Journal of Plant Pathology*, 96(1S):1-136.
- Martin, R. R., Eastwell, K. C., Lamprecht, S. & Tzanetakis I. E.** 2005. Survey for viruses of grapevine in Oregon and Washington. *Plant Disease*, 89:763-766.
- Meng, B., Martelli, G. P., Golino, D. A. & Fuch, M. (eds.).** 2017. *Grapevine Viruses: Molecular Biology, Diagnostics and Management*. Springer, Cham, Switzerland. 698 pp.
- Milkus, B. N., & Goodman, R. N.** 1999. A survey of Missouri vineyards for the presence of five grape viruses. *American Journal of Enology and Viticulture*, 50(1):133-134.
- Morales, R. Z., & Monis, J.** 2007. First detection of grapevine leafroll associated virus-7 in California vineyards. *Plant Disease*, 91(4):465-465.
- Naidu, R. A., & Mekuria, T. A.** 2010. First report of grapevine fleck virus from Washington vineyards. *Plant Disease*, 94(6):784-784.
- Ouertani, R., Savino, V., Minafra, A., Boscia, D., Castellano, M. A., Martelli, G. P. & Greco, N.** 1992. Properties of a previously undescribed grapevine nepovirus from Tunisia. *Archives of Virology*, 126:107-117.
- Pearson, R. C. & Goheen, A. C.** 1988. *Compendium of Grape Diseases*. American Phytopathological Society, St. Paul, MN, USA. 93 pp.
- Poojari, S., Lowery, T., Rott, M., Schmidt, A.-M., Delury, N., Boule, J. & Urbez-Torres, J. R.** 2016. First report and prevalence of grapevine fleck virus in grapevines (*Vitis vinifera*) in Canada. *Plant Disease*, 100(5):1028.
- Sharma, A. M., Wang, J., Duffy, S., Zhang, S., Wong, M. K., Rashed, A., Cooper, M .L., Daane, K. M., & Almeida, R. P. P.** 2011. Occurrence of grapevine leafroll-associated virus complex in Napa Valley. *PLOS ONE*, 6(10): e26227.
- Silva, J. M. F., Al Rwahnih, M., Blawid, R., Nagata, T. & Fajardo, T. V. M.** 2017. Discovery and molecular characterization of a novel enamovirus, grapevine enamovirus-1. *Virus Genes*, 53(4):667-671.
- Uyemoto, J. K., Taschenberg, E. F. & Hummer, D. K.** 1977. Isolation and identification of a strain of grapevine Bulgarian latent virus in Concord grapevine in New York State. *Plant Disease Reporter*, 61(11):949-953.
- Wilcox, W. F., Gubler, W. D. & Uyemoto, J. K.** 2015. *Compendium of Grape Diseases, Disorders, and Pests*. 2nd Edition. American Phytopathological Society, St. Paul, MN, USA. 232 pp.
- Xiao, H., & Meng, B.** 2016. First report of grapevine asteroid mosaic-associated virus and grapevine rupestris vein feathering virus in grapevines in Canada. *Plant Disease*, 100(10):2175.
- Zhang, Y., Singh, K., Kaur, R. & Qiu, W.** 2011. Association of a novel DNA virus with the grapevine vein-clearing and vine decline syndrome. *Phytopathology*, 101(9):1081-1090.

Table 2: Fungal pathogens (incl. Chromista) of grapevine

Last updated December 2021.

PEST	DISEASE	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
<i>Botryosphaeria dothidea</i> (Moug.: Fr.) Ces. and De Not. Syn.: <i>Fusicoccum aesculi</i> Corda	Botryosphaeria dieback	Ammad <i>et al.</i> , 2014; Smith and Stanosz, 2001; Wilcox <i>et al.</i> , 2015	P2	P2	Ab3
<i>Cadophora luteo-olivacea</i> (J.F.H. Beyma) T.C. Harr. and McNew	Grapevine trunk disease	Gramaje <i>et al.</i> , 2011; Navarrete <i>et al.</i> , 2011; Raimondo <i>et al.</i> , 2019; Travadon <i>et al.</i> , 2015	P2	P2	Ab1
<i>Colletotrichum acutatum</i> J.H. Simmonds Syn.: <i>Glomerella acutata</i> Guerber and J.C. Correll	Anthraxnose, bitter rot; blackspot	Haviland <i>et al.</i> , 2019; Hong <i>et al.</i> , 2008; Wilcox <i>et al.</i> , 2015	P2	P2	P2
<i>Coniella diplodiella</i> (Speg.) Petr. and Syd. Syn.: <i>Coniothyrium diplodiella</i> (Speg.) Sacc., <i>Pilidiella diplodiella</i> (Speg.) Crous and Van Niekerk	Dieback, white rot	Blake and Williamson, 2015; Locci and Quaroni, 1972; Wilcox <i>et al.</i> , 2015	P2	P2	Ab1
<i>Coniella vitis</i> Chethana, J.Y. Yan, X.H. Li and K.D. Hyde	White rot	Chethana <i>et al.</i> , 2017	Ab1	Ab1	Ab1
<i>Cryptovalsa ampelina</i> (Nitschke) Fuckel	Dieback	Díaz <i>et al.</i> , 2011; Luque <i>et al.</i> , 2006; Trouillas and Gubler, 2010	P2	P2	Ab1
<i>Cytospora chrysosperma</i> (Pers.: Fr.) Fr. Syn.: <i>Valsa sordida</i> Nitschke	Perennial canker	Arzanlou and Narmani, 2015; Lawrence <i>et al.</i> , 2018	P7	P2	Ab3
<i>Cytospora vinacea</i> D.P. Lawr., Travadon and Pouzoulet	Canker	Lawrence <i>et al.</i> , 2017b	Ab1	P2	Ab1
<i>Cytospora viticola</i> D.P. Lawr., Travadon and Pouzoulet	Canker	Lawrence <i>et al.</i> , 2017b; Oksal <i>et al.</i> , 2020	P2	P2	Ab1
<i>Dactylonectria macrodidyma</i> (Halleen, Schroers and Crous) L. Lombard and Crous, <i>Cylindrocarpon macrodidymum</i> Schroers, Halleen and Crous	Black foot	Petit <i>et al.</i> , 2011; Probst <i>et al.</i> , 2019; Úrbez-Torres <i>et al.</i> , 2012; Wilcox <i>et al.</i> , 2015	P2	P2	Ab1

PEST	DISEASE	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
<i>Diaporthe ampelina</i> (Berk and M.A. Curtis) R.R. Gomes, C. Glienke and Crous Syn.: <i>Phomopsis viticola</i> (Sacc.) Sacc.	Phomopsis cane and leaf spot	Chen <i>et al.</i> , 2014; Pscheidt and Ocamb, 2020; Wilcox <i>et al.</i> , 2015	P3	P2	Ab1
<i>Diplodia corticola</i> A.J.L. Phillips, A. Alves and J. Luque Syn.: <i>Botryosphaeria corticola</i> A.J.L. Phillips, A. Alves and J. Luque	Canker	Reed <i>et al.</i> , 2018; Úrbez-Torres <i>et al.</i> , 2010b; Varela <i>et al.</i> , 2011	Ab1	P2	Ab3
<i>Elsinoe ampelina</i> Shear	Anthraco nose	Santos <i>et al.</i> , 2018; Wilcox <i>et al.</i> , 2015; Yun <i>et al.</i> , 2007	P2	P2	Ab3
<i>Eutypa laevata</i> (Nitschke) Sacc.	Eutypa dieback	Rolshausen <i>et al.</i> , 2014	P2	P2	Ab1
<i>Eutypa leptoplaca</i> (Mont.) Rappaz	Eutypa dieback	Pscheidt and Ocamb, 2020; Trouillas and Gubler, 2004; Trouillas and Gubler, 2010	Ab1	P2	Ab1
<i>Ilyonectria destructans</i> (Zinssm.) Rossman, L. Lombard and Crous Syn.: <i>Cylindrocarpon destructans</i> (Zinssm.) Scholten, <i>Neonectria radicularis</i> (Gerlach and L. Nilsson) Mantiri and Samuels	Black foot	Khorasani, 2013; Petit and Gubler, 2005; Pscheidt and Ocamb, 2020; Wilcox <i>et al.</i> , 2015	P2	P2	Ab3
<i>Neofusicoccum luteum</i> (Pennycook and Samuels) Crous, Slippers and A.J.L. Phillips Syn.: <i>Botryosphaeria lutea</i> A.J.L. Phillips	Canker and dieback	Chebil <i>et al.</i> , 2013; Savocchia <i>et al.</i> , 2007; Úrbez-Torres <i>et al.</i> , 2006	Ab1	P2	Ab3
<i>Neofusicoccum mediterraneum</i> Crous, M.J. Wingf. and A.J.L. Phillips	Cankers and dieback	Martin <i>et al.</i> , 2011; Úrbez-Torres <i>et al.</i> , 2010a; Varela <i>et al.</i> , 2011	Ab1	P2	Ab1
<i>Neonectria obtusispora</i> (Cooke and Harkn.) Rossman, L. Lombard and Crous Syn.: <i>Cylindrocarpon obtusisporum</i> (Cooke and Harkn.) Wollenw.	Black foot	Petit <i>et al.</i> , 2011; Pscheidt and Ocamb, 2020; Scheck <i>et al.</i> , 1998a; Wilcox <i>et al.</i> , 2015	P7	P2	Ab1
<i>Phaeoacremonium angustius</i> W. Gams, Crous and M.J. Wingf.	Esca and Petri disease	Aroca and Raposo, 2009; Chicau <i>et al.</i> , 2000	P2	P2	Ab1
<i>Phaeoacremonium fraxinopennsylvanicum</i> (T.E. Hinds) D. Gramaje, L. Mostert and Crous	Esca and Petri disease	Eskalen <i>et al.</i> , 2005	P2	P2	Ab1

PEST	DISEASE	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
Syn.: <i>Phaeoacremonium mortoniae</i> Crous and W. Gams, <i>Togninia fraxinopennsylvanica</i> (T.E. Hinds) Hausner, Eyjólfsdóttir and J. Reid					
<i>Phaeoacremonium inflatipes</i> W. Gams, Crous and M.J. Wingf.	Esca and Petri disease	Aroca and Raposo, 2009; Scheck <i>et al.</i> , 1998b	Ab1	P2	Ab3
<i>Phaeoacremonium krajdinii</i> L. Mostert, Summerb. and Crous Syn.: <i>Togninia krajdinii</i> L. Mostert, W. Gams and Crous	Petri disease	Gramaje <i>et al.</i> , 2011a	Ab1	P2	Ab3
<i>Phaeoacremonium minimum</i> (Tul. and C. Tul.) D. Gramaje, L. Mostert and Crous Syn.: <i>Phaeoacremonium aleophilum</i> W. Gams, Crous, M.J. Wingf. and Mugnai, <i>Togninia minima</i> (Tul. and C. Tul.) Berl., <i>Togninia alnicola</i> (Ellis and Everh.) Berl.	Leaf stripe, Black measles	Haviland <i>et al.</i> , 2019a; Urbez-Torres <i>et al.</i> , 2012; Wilcox <i>et al.</i> , 2015	P2	P2	Ab1
<i>Phaeoacremonium parasiticum</i> (Ajello, Georg and C.J.K. Wang) W. Gams, Crous and M.J. Wingf. Syn.: <i>Phialophora parasitica</i> Ajello, Georg and Wang	Dieback	Aroca and Raposo, 2009; Groenewald <i>et al.</i> , 2001; OSU, 2020; Romero-Rivas <i>et al.</i> , 2009	Ab1	P2	Ab1
<i>Phaeomoniella chlamydospora</i> (W. Gams, Crous, M.J. Wingf. and Mugnai) Crous and W. Gams Syn.: <i>Phaeoacremonium chlamydosporum</i> W. Gams, Crous, M.J. Wingf. and Mugnai	Black measles	Haviland <i>et al.</i> , 2019a; Scheck <i>et al.</i> , 1998b; Wilcox <i>et al.</i> , 2015	P2	P2	Ab1
<i>Phymatotrichopsis omnivora</i> (Shear) Hennebert Syn.: <i>Phymatotrichum omnivorum</i> (Shear) Duggar	Texas root rot	Davis <i>et al.</i> , 2017; Smith, 2019; Wilcox <i>et al.</i> , 2015	Ab1	P2	Ab3
<i>Phytophthora cinnamomi</i> Rands	Phytophthora crown and root rot	Latorre <i>et al.</i> , 1997; Nouri <i>et al.</i> , 2017; Wilcox <i>et al.</i> , 2015	P7	P2	P2
<i>Phytophthora citricola</i> Sawada.	Root rot	Erwin and Ribeiro, 1996; Schwingle <i>et al.</i> , 2007	P7	P2	Ab3
<i>Phytophthora cryptogea</i> Pethybr. and Laff.	Phytophthora crown and root rot	Koike <i>et al.</i> , 2019; Latorre <i>et al.</i> , 1997; McKeever and Chastagner,	P7	P2	P2

PEST	DISEASE	REFERENCES	PRESENCE / ABSENCE		
			CAN	USA	MEX
		2016; Wilcox <i>et al.</i> , 2015			
<i>Phytophthora drechsleri</i> Tucker	Phytophthora crown and root rot	Latorre <i>et al.</i> , 1997; Olson and Benson, 2011; Olson <i>et al.</i> , 2016; Wilcox <i>et al.</i> , 2015	P7	P2	P2
<i>Roesleria subterranea</i> (Weinm.) Redhead	Grapevine root rot	EPPO, 2012; Miles and Schilder, 2009; Neuhauser <i>et al.</i> , 2011	P2	P2	Ab1
<i>Rosellinia necatrix</i> Prill. Syn.: <i>Dematophora necatrix</i> Hartig	Dematophora root rot	Mansoori and Dorostkar, 2008; Wilcox <i>et al.</i> , 2015; Windbiel-Rojas <i>et al.</i> , 2020	Ab1	P2	P2
<i>Seimatosporium botan</i> Sat. Hatak. and Y. Harada	Trunk canker	Díaz <i>et al.</i> , 2012; Díaz <i>et al.</i> , 2013; Lawrence <i>et al.</i> , 2017	Ab1	Ab1	Ab1
<i>Verticillium dahliae</i> Kleb.	Verticillium wilt	Gubler <i>et al.</i> , 2004; Wilcox <i>et al.</i> , 2015; Zhang <i>et al.</i> , 2009	P2	P2	P2

References

- Ammad, F., Benchabane, M., Toumi, M., Belkacem, N., Guesmi, A., Ameer, C., Lecomte, P. & Merah, O. 2014. Occurrence of *Botryosphaeriaceae* species associated with grapevine dieback in Algeria. *Turkish Journal of Agriculture and Forestry* 38:865-876.
- Aroca, A., & Raposo, R. 2009. Pathogenicity of *Phaeoacremonium* species on grapevines. *Journal of Phytopathology* 157(7-8):413-419.
- Arzanlou, M. & Narmani, A. 2015. ITS sequence data and morphology differentiate *Cytospora chrysosperma* associated with trunk disease of grapevine in northern Iran. *Journal of Plant Protection Research* 55(2):117-125.
- Blake, J. H. & Williamson, M. 2015. Index of plant diseases in South Carolina (Third Edition). Clemson Cooperative Extension.
- Chebil, S., Fersi, R., Yakoub, A., Chenenaoui, S., Chattaoui, M., Melki, I., Zemni, H., Rhouma, A., Durante, G., Zacchi, E. & Mliki, A. 2013. First report of *Botryosphaeria dothidea*, *Diplodia seriata*, and *Neofusicoccum luteum* associated with canker and dieback of grapevines in Tunisia. *Plant Disease* 98(3):420-420.
- Chen, S. F., Morgan, D. P. & Michailides, T. J. 2014. *Botryosphaeriaceae* and *Diaporthaceae* associated with panicle and shoot blight of pistachio in California, USA. *Fungal Diversity* 67(1):157-179.
- Chethana, K. W. T., Zhou, Y., Zhang, W., Liu, M., Xing, Q. K., Li, X. H., Yan, J. Y. & Hyde, K. D. 2017. *Coniella vitis* sp. nov. is the common pathogen of white rot in Chinese vineyards. *Plant Disease* 101(12):2123-2136.
- Chicau, G., Aboim-Inglez, M. Cabral, S. & Cabral J. P. S. 2000. *Phaeoacremonium chlamydosporum* and *Phaeoacremonium angustius* associated with esca and grapevine decline in Vinho Verde grapevines in northwest Portugal. *Phytopathologia Mediterranea* 39(1):80-86.
- Davis, R. M., Frate, C. A. & Putnam, D. H. 2017. UC IPM Pest Management Guidelines: Alfalfa Phymatotrichopsis root rot (Texas Root Rot) (UC ANR Publication 3430). Agriculture and Natural Resources, University of California.
- Díaz, G. A., Prehn, D. & B. A. Latorre, B. A. 2011. First report of *Cryptovalsa ampelina* and *Eutypella leprosa* associated with grapevine trunk diseases in Chile. *Plant Disease* 95(4):490-490.
- Díaz, G. A., Auger, J., Besoain, X., Bordeu, E. & Latorre, B. A. 2013. Prevalence and pathogenicity of fungi associated with grapevine trunk diseases in Chilean vineyards. *Ciencia e Investigacion Agraria* 40(2):327-339.
- Díaz, G. A., Elfar, K., & Latorre, B. A. 2012. First report of *Seimatosporium botan* associated with trunk disease of Grapevine (*Vitis vinifera*) in Chile. *Plant Disease* 96(11):1696.
- EPPO. 2012. *Roesleria subterranea*: an emerging disease on grapevine? (2012/190). European and Mediterranean Plant Protection Organization. 7-8 pp.
- Erwin, D. C. & Ribeiro, O. K. 1996. *Phytophthora* Diseases Worldwide. The American Phytopathological Society, St. Paul, MN, USA. 562 pp.
- Eskalen, A., Rooney-Latham, S. & Gubler, W. D. 2005. Occurrence of *Togninia fraxinopennsylvanica* on esca-diseased grapevines (*Vitis vinifera*) and declining ash trees (*Fraxinus latifolia*) in California. *Plant Disease* 89(5):528-528.
- Gramaje, D., Mostert, L. & Armengol, J. 2011. Characterization of *Cadophora luteo-olivacea* and *C. melinii* isolates obtained from grapevines and environmental samples from grapevine nurseries in Spain. *Phytopathologia Mediterranea* 50: S112-S126.
- Groenewald, M., Kang, J. -C., Crous, P. W. & Gams, W. 2001. ITS and β -tubulin phylogeny of *Phaeoacremonium* and *Phaeomoniella* species. *Mycological Research* 105(6):651-657.
- Gubler, W. D., Baumgartner, K., Browne, G. T., Eskalen, A., Rooney Latham, S., Petit, E. & Bayramian, L. A. 2004. Root diseases of grapevines in California and their control. *Australasian Plant Pathology* 33:157-165.
- Haviland, D. R., Symmes, E. J., Adaskaveg, J. E., Duncan, R. A., Roncoroni, J. A., Gubler, W.

- D., Hanson, B., Hembree, K. J., Holtz, B., Stapleton, J. J., Tollerup, K. E., Trouillas, F. P. & Zalom, F. G.** 2019. UC IPM Pest Management Guidelines: Almond (UC ANR Publication 3431). UC Statewide Integrated Pest Management Program, Oakland, CA.
- Hong, S. -K., Kim, W. -G., Yun, H. -K. & Choi K. -J.** 2008. Morphological variations, genetic diversity and pathogenicity of *Colletotrichum* species causing grape ripe rot in Korea. *The Plant Pathology Journal* 24(3):269-278.
- Khorasani, M.** 2013. *Cylindrocarpon* species in Pacific Northwest Douglas-fir nurseries: diversity and effects of temperature and fungicides on mycelial growth. Master Thesis, University of Washington, Ann Arbor.
- Koike, S. T., Stanghellini, H., Mauzey, S. J., Burkhardt, A. & Stanghellini M. S.** 2019. First report of *Phytophthora* root and bulb rot caused by *Phytophthora cryptogea* on shallot in the United States. *Plant Disease* 103(6):1436-1436.
- Latorre, B. A., Wjlcx, W. F. & Banaos, M. P.** 1997. Crown and root rots of table grapes caused by *Phytophthora* spp. in Chile. *Vitis* 36(4):195-197.
- Lawrence, D. P., Holland, L. A., Nouri, M. T., Travadon, R., Abramians, A., Michailides, T. J. & Trouillas, F. P.** 2018. Molecular phylogeny of *Cytospora* species associated with canker diseases of fruit and nut crops in California, with the descriptions of ten new species and one new combination. *IMA Fungus* 9(2):330-370.
- Lawrence, D. P., Travadon, R. & Baumgartner, K.** 2017. Novel *Seimatosporium* species from grapevine in Northern California and their interactions with fungal pathogens involved in the trunk-disease complex. *Plant Disease* 102(6):1081-1092.
- Lawrence, D. P., Travadon, R., Pouzoulet, J., Rolshausen, P. E., Wilcox, W. F. & Baumgartner, K.** 2017b. Characterization of *Cytospora* isolates from wood cankers of declining grapevine in North America, with the descriptions of two new *Cytospora* species. *Plant Pathology* 66(5):713-725.
- Locci, R. & Quaroni, S.** 1972. Studies on *Coniothyrium diplodiella* I. Isolation, cultivation and identification of the fungus. *Rivista di Patologia Vegetale* 8(1):59-82.
- Luque, J., Sierra, D., Torres, E. & Garcia, F.** 2006. *Cryptovalsa ampelina* on grapevines in N.E. Spain: identification and pathogenicity. *Phytopathologia Mediterranea* 45:S101–S109.
- Mansoori, B. & Dorostkar M.** 2008. Reactions of some grape cultivars to *Dematophora necatrix*. *Vitis* 47(4):231-233.
- Martin, M. T., Martin, L. & Cuesta, M. J.** 2011. First report of *Neofusicoccum mediterraneum* and *N. australe* causing decay in *Vitis vinifera* in Castilla y León, Spain. *Plant Disease* 95(7):876-876.
- McKeever, K. M. & Chastagner, G. A.** 2016. A survey of *Phytophthora* spp. associated with *Abies* in U.S. christmas tree farms. *Plant Disease* 100(6):1161-1169.
- Miles, T. D. & Schilder, A. M. C.** 2009. First report of grape root rot caused by *Roesleria subterranea* in Michigan. *Plant Disease* 93(7):765-765.
- Navarrete, F., Abreo, E., Martínez, S., Bettucci, L. & Lupo, S.** 2011. Pathogenicity and molecular detection of Uruguayan isolates of *Greeneria uvicola* and *Cadophora luteo-olivacea* associated with grapevine trunk diseases. *Phytopathologia Mediterranea* 50:S166-S175.
- Neuhauser, S., Huber, L. & Kirchmair, M.** 2011. Is *Roesleria subterranea* a primary pathogen or a minor parasite of grapevines? Risk assessment and a diagnostic decision scheme. *European Journal of Plant Pathology* 130(4):503-510.
- Nouri, M. T., Holland, L. A., Yaghmour, M. A., Doll, D. A., Browne, G. T. & Trouillas, F. P.** 2017. First report of *Phytophthora cinnamomi* causing trunk canker of almond in California. *Plant Disease* 102(1):253.
- Oksal, E., Çelik, Y. & Özer, G.** 2020. First report of canker and dieback caused by *Cytospora viticola* on grapevine in Turkey. *Journal of Plant Pathology* 102(1):239-239.
- Olson, H. A. & Benson, D. M.** 2011. Characterization of *Phytophthora* spp. on floriculture crops in North Carolina. *Plant Disease* 95(8):1013-1020.
- Olson, J. D., Damicone, J. P. & Kahn, B. A.** 2016. Identification and characterization of isolates of *Pythium* and *Phytophthora* spp. from snap beans with Cottony Leak. *Plant Disease*

100(7):1446-1453.

- OSU.** 2020. Grapevine trunk diseases in California and Oregon. Oregon State University. Last accessed 4/7/2020, <https://bpp.oregonstate.edu/main/grapevine-trunk-diseases>.
- Petit, E., Barriault, E., Baumgartner, K., Wilcox, W. F. & Rolshausen, P. E.** 2011. *Cylindrocarpon* species associated with black-foot of grapevine in Northeastern United States and Southeastern Canada. *American Society for Enology and Viticulture* 62(2):177-183.
- Petit, E. & Gubler, W.D.** 2005. Characterization of *Cylindrocarpon* species, the cause of black foot disease of grapevine in California. *Plant Disease* 89(10):1051-1059.
- Probst, C. M., Ridgway, H. J., Jaspers, M. V. & Eirian Jones, E.** 2019. Pathogenicity of *Ilyonectria liriiodendri* and *Dactylonectria macrodidyma* propagules in grapevines. *European Journal of Plant Pathology* 154(2):405-421.
- Pscheidt, J. W. & Ocamb, C. M.** 2020. Pacific Northwest Plant Disease Management Handbook. Oregon State University. <https://pnwhandbooks.org/>.
- Raimondo, M. L., Carlucci, A., Cicca-Rone, C., Sadallah, A. & Lops, F.** 2019. Identification and pathogenicity of *lignicolous* fungi associated with grapevine trunk diseases in southern Italy. *Phytopathologia Mediterranea* 58(3):639-662.
- Reed, S. E., English, J. T., Lalk, S. R. & Tosie, K. M.** 2018. First report of *Diplodia corticola* causing stem and branch cankers on white oak (*Quercus alba*) in Missouri, U.S.A. *Plant Disease* 102(12):2657.
- Rolshausen, P. E., K. Baumgartner, R. Travadon, P. Fujiyoshi, J. Pouzoulet, & W. F. Wilcox.** 2014. Identification of *Eutypa* spp. causing eutypa dieback of grapevine in Eastern North America. *Plant Disease* 98:483-491.
- Romero-Rivas, L. C., Álvarez, L.A., Gramaje, D., Armengol, J. & Cadenas-Giraldo, C.** 2009. First report of *Phaeoacremonium parasiticum* causing petri disease of grapevine in Peru. *Plant Disease* 93(2):200-200.
- Santos, R. F., Ciampi-Guillardi, M., Amorim, L., Massola Junior, N. S. & Sposito, M. B.** 2018. Aetiology of anthracnose on grapevine shoots in Brazil. *Plant Pathology* 67:692-706.
- Savocchia, S., Steel, C. C., Stodart, B. J. & Somers, A.** 2007. Pathogenicity of *Botryosphaeria* species isolated from declining grapevines in sub-tropical regions of Eastern Australia. *Vitis* 46(1):27-32.
- Scheck, H. J., Vasquez, S.J., Gubler, W. D. & Fogle, D.** 1998a. First report of black-foot disease, caused by *Cylindrocarpon obtusisporum*, of grapevine in California. *Plant Disease* 82(4):448-448.
- Scheck, H. J., Vasquez, S.J., Gubler, W. D. & Fogle, D.** 1998b. First report of three *Phaeoacremonium* spp. causing young grapevine decline in California. *Plant Disease* 82(5):590-590
- Schwingle, B. W., Smith, J. A. & Blanchette, R. A.** 2007. *Phytophthora* species associated with diseased woody ornamentals in Minnesota nurseries. *Plant Disease* 91(1):97-102.
- Smith, D.** 2019. *Phymatotrichum* root rot, cotton root rot, Texas root rot. Oklahoma State University. <https://grapes.extension.org/phymatotrichum-root-rot-cotton-root-rot-texas-root-rot/>.
- Smith, D. R. & Stanosz, G. R.** 2001. Molecular and morphological differentiation of *Botryosphaeria dothidea* (anamorph *Fusicoccum aesculi*) from some other fungi with *Fusicoccum* anamorphs. *Mycologia* 93(3):505-515.
- Travadon, R., Lawrence, D. P., Rooney-Latham, S., Gubler, W. D., Wilcox, W. F., Rolshausen, P. E. & Baumgartner, K.** 2015. Cadophora species associated with wood-decay of grapevine in North America. *Fungal Biology* 119(1):53-66.
- Trouillas, F. P. & Gubler, W. D.** 2004. Identification and characterization of *Eutypa leptoplaca*, a new pathogen of grapevine in Northern California. *Mycological Research* 108(10):1195-1204.
- Trouillas, F. P. & Gubler, W. D.** 2010. Pathogenicity of *Diatrypaceae* species in grapevines in California. *Plant Disease* 94(7):867-872.
- Úrbez-Torres, J. R., Leavitt, G. M., Voegel, T. M. & Gubler, W. D.** 2006. Identification and distribution of *Botryosphaeria* spp. associated with grapevine cankers in California. *Plant Disease* 90(12):1490-1503.

- Úrbez-Torres, J. R., Peduto, F. & Gubler, W. D.** 2010a. First report of grapevine cankers caused by *Lasiodiplodia crassispora* and *Neofusicoccum mediterraneum* in California. *Plant Disease* 94(6):785-785.
- Úrbez-Torres, J. R., Peduto, F. & Gubler, W. D.** 2012. First report of *Ilyonectria macrodidyma* causing root rot of olive trees (*Olea europaea*) in California. *Plant Disease* 96(9):1378-1378.
- Úrbez-Torres, J. R., Peduto, F., Rooney-Latham, S. & Gubler, W. D.** 2010b. First report of *Diplodia corticola* causing grapevine (*Vitis vinifera*) cankers and trunk cankers and dieback of canyon live oak (*Quercus chrysolepis*) in California. *Plant Disease* 94(6):785-785.
- Varela, C. P., Fernández, V. R., Casal, O. A. & Vázquez, J. P. M.** 2011. First report of cankers and dieback caused by *Neofusicoccum mediterraneum* and *Diplodia corticola* on grapevine in Spain. *Plant Disease* 95(10):1315-1315.
- Wilcox, W. F., Gubler, W. D. & Uyemoto, J. K.** 2015. Compendium of Grape Diseases, Disorders, and Pests (Second Edition). American Phytopathological Society, St. Paul, MN, USA. 232 pp.
- Windbiel-Rojas, K., Laning, C., Blecker, L. & Alfaro, M.** 2020. Pests in Gardens and Landscapes. Agriculture and Natural Resources, University of California. <http://ipm.ucanr.edu/PMG/>.
- Yun, H. K. Louime, C. & Lu, J.** 2007. First report of anthracnose caused by *Elsinoe ampelina* on muscadine grapes (*Vitis rotundifolia*) in Northern Florida. *Plant Disease* 91(7):905-905.
- Zhang, L., Zhang, G. L., Qian, X. & Li, G. Y.** 2009. First report of *Verticillium* wilt of grapevine (*Vitis vinifera*) caused by *Verticillium dahliae* in China. *Plant Disease* 93(8):841-841.

Table 3. Bacterial pathogens of grapevine

Last updated December 2021.

PEST	DISEASE	REFERENCES	PRESENCE/ABSENCE		
			CAN	US	MEX
<i>Candidatus Phytoplasma asteris</i> Lee <i>et al.</i> (16SrI-A, -B, -C)	Grapevine yellows/aster yellows	Angelini <i>et al.</i> , 2018; Davis <i>et al.</i> , 2018	P2	P2	P2
<i>Candidatus Phytoplasma aurantifolia</i> (16SrII-B)	Grapevine yellows	Bertaccini, 2018	Ab1	Ab1	Ab3
<i>Candidatus Phytoplasma australasia</i> (16SrII-A, -D)	Australian grapevine yellows	Angelini, <i>et al.</i> , 2018; Bertaccini, 2018	Ab1	Ab1	P2
<i>Candidatus Phytoplasma brasiliense</i> (16SrXV-A)	Grapevine yellows	Bertaccini, A. 2018	Ab1	Ab1	Ab1
<i>Candidatus Phytoplasma fraxini</i> Griffiths <i>et al.</i> (16SrVII-A)	Ash/elm yellows	Gajardo <i>et al.</i> , 2009; Ghayeb Zamharir <i>et al.</i> , 2017	P7	P2	Ab1
<i>Candidatus Phytoplasma phoenicium</i> (16SrIX,-B)	Grapevine yellows	Bertaccini, 2018	Ab1	Ab1	Ab1
<i>Candidatus Phytoplasma prunorum</i> (16SrX-B)	Grapevine yellows	Bertaccini, 2018	Ab1	Ab1	Ab1
<i>Candidatus Phytoplasma pruni</i> Davis <i>et al.</i> (16SrIII,-A)	Grapevine yellows	Davis <i>et al.</i> , 2019	P7	P2	P2
<i>Candidatus Phytoplasma solani</i> Quaglino <i>et al.</i> (16SrXII-A)	Grapevine yellows	Angelini <i>et al.</i> , 2018; Landi <i>et al.</i> , 2019	Ab5	Ab1	Ab1
<i>Candidatus Phytoplasma trifolii</i> Hiruki and Wang (16SrVI-A)	Grapevine yellows	Bertaccini, 2018; Flower <i>et al.</i> , 2017; Jacobs <i>et al.</i> , 2003	P7	P2	P2
<i>Candidatus Phytoplasma ulmi</i> (16SrV-A)	Grapevine yellows	Bertaccini, 2018	P7	Ab1	Ab1
<i>Candidatus Phytoplasma vitis</i> Marzorati <i>et al.</i> (16SrV-A, -B, -C, -D)	Flavescence dorée, Grapevine flavescence dorée, Palatinate grapevine yellows	Angelini <i>et al.</i> , 2018; Eveillard <i>et al.</i> , 2016	Ab1	Ab1	Ab1
<i>Pseudomonas syringae</i> pv. <i>Syringae</i> Van Hall	Bacterial Inflorescence Rot	Hall <i>et al.</i> 2016	P7	P2	Ab3
<i>Rhizobium vitis</i> (Ophel and Kerr 1990) Young <i>et al.</i> ; Syn.: <i>Agrobacterium vitis</i> (Ophel and Kerr)	Crown Gall	Kawaguchi <i>et al.</i> , 2019; Voegel and Nelson, 2018	P2	P2	Ab3
<i>Xanthomonas citri</i> pv. <i>viticola</i> (Nayudu 1972) Dye, Syn.:	Bacterial Canker	Chand and Kishun, 1990; Naue <i>et al.</i> , 2014	Ab1	Ab1	P2

PEST	DISEASE	REFERENCES	PRESENCE/ABSENCE		
			CAN	US	MEX
<i>Xanthomonas campestris</i> pv. <i>viticola</i> (Nayudu 1972) Dye					
<i>Xylella fastidiosa</i> Wells <i>et al.</i>	Pierce's Disease	Hill and Purcell, 1995, Kirkpatrick, B. C. 2015; Wilcox <i>et al.</i> , 2015	P7	P2	P3
<i>Xylophilus ampelinus</i> (Panagopoulos Willems <i>et al.</i>)	Bacterial blight of grapevine, Canker of grapevine	Hand, F. P. 2015; Komatsu and Kondo, 2015; Soto-Giron <i>et al.</i> , 2016; Wilcox <i>et al.</i> , 2015	Ab1	P2	Ab1

References

- Angelini E., Constable F., Duduk B., Fiore N., Quaglino F. & Bertaccini, A. 2018. Grapevine Phytoplasmas. In: Rao G., Bertaccini A., Fiore N., Liefting L. (eds) *Phytoplasmas: Plant Pathogenic Bacteria - I*. Springer, Singapore.
- Chand, R. I. & Kishun, R. 1990. Outbreak of grapevine bacterial canker disease in India. *Vitis* 29(3):183-188.
- Bertaccini, A. 2018. Grapevine Phytoplasmas a 2014-2018 update in: Proceedings of the 19th Congress of ICVG. Presented at the International Council for the Study of Virus and Virus-Like Diseases of the Grapevine, Santiago, Chile, pp. 83-89.
- Davis, R. E., Dally, E. L., Webb, C. & Appel, J. A. 2019. First report of North American Grapevine Yellows (NAGY) in Kansas: Detection of '*Candidatus* Phytoplasma pruni'- and '*Ca.* Phytoplasma asteris'-related strains in diseased plants. *Plant Disease* 103(4):759.
- Davis, R. E., Dally, E. L., Zhao, Y. Wolf, T. K. & Smith, A. H. 2018. Genotyping points to divergent evolution of '*Candidatus* Phytoplasma asteris' strains causing North American Grapevine Yellows and strains causing Aster Yellows. *Plant Disease* 102(9):1696-1702.
- Eveillard, S., Jollard, C., Labroussaa, F., Khalil, D., Perrin, M., Desqué, D., Salar, P., Razan, F., Hévin, C., Bordenave, L., Foissac, X., Masson, J. E. & Malembic-Maher, S. 2016. Contrasting susceptibilities to Flavescence Dorée in *Vitis vinifera*, rootstocks and wild *Vitis* species. *frontier in Plant Science* 7(1762).
- Flower, C. E., N. Hayes-Plazolles, J. M. Slavicek, & C. Rosa. 2017. First report of '*Candidatus* Phytoplasma trifolii'-related strain of 16SrVI-A phytoplasma subgroup, associated with elm yellows disease in american elm (*Ulmus americana* L.) in Ohio, U.S.A. *Plant Disease* 102(2):438.
- Landi, L., Murolo, S. & Romanazzi, G. 2019. Detection of '*Candidatus* Phytoplasma solani' in roots from Bois noir symptomatic and recovered grapevines. *Scientific Reports* 9(2013):1-12.
- Gajardo, A., Fiore, N., Prodan, S., Paltrinieri, S., Botti, S., Pino, A. M., Zamorano, A., Montealegre, J. & Bertaccini A. 2009. Phytoplasmas associated with grapevine yellows disease in Chile. *Plant Disease* 93(8):789-796.
- Ghayeb Zamharir, M., Paltrinieri, S., Hajivand, S., Taheri, M. & Bertaccini, A. 2017. Molecular identification of diverse '*Candidatus* Phytoplasma' species associated with grapevine decline in Iran. *Journal of Phytopathology* 165:407-413.
- Hall, S. J., Dry, I. B., Blanchard, C. L., & Whitelaw-Weckert, M. A. 2016. Phylogenetic relationships of *Pseudomonas syringae* pv. *syringae* isolates associated with bacterial inflorescence rot in grapevine. *Plant Disease* 100(3): 607-619.
- Hand, F. P. 2015. Disease Caused by Bacteria: Bacterial Blight. In Gubler, W. D., Uyemoto, J. K., Wilcox, W. F. (Eds.), *Compendium of Grape Diseases, Disorders, and Pests* (Vol. 2, pp. 94-95). St. Paul, Minnesota: American Pathological Society.
- Hill, B. L., & Purcell, A. H. 1995. Multiplication and movement of *Xylella fastidiosa* within grapevine and four other plants. *Phytopathology* 85(11):1368-1372.
- Jacobs, K. A., Lee, I. M., Griffiths, H. M., Miller, F. D. & Bottner, K. D. 2003. A new member of the Clover proliferation phytoplasma group (16SrVI) associated with Elm Yellows in Illinois *Plant Disease* 87(3):241-246.
- Kirkpatrick, B. C. 2015. Disease Caused by Bacteria: Pierce's Disease. In Gubler, W. D., Uyemoto, J. K., Wilcox, W. F. (Eds.), *Compendium of Grape Diseases, Disorders, and Pests* (Vol. 2, pp. 98-103). St. Paul, Minnesota: American Pathological Society.
- Kawaguchi, A., Nita, M., Ishii, T., Watanabe, M. & Noutoshi, Y. 2019. Biological control agent *Rhizobium* (= *Agrobacterium*) *vitis* strain ARK-1 suppresses expression of the essential and non-essential *vir* genes of tumorigenic *R. vitis*. *BMC Research Notes* 12(1):1-6.
- Komatsu, T. & Kondo, N. 2015. Winter habitat of *Xylophilus ampelinus*, the cause of bacterial blight of grapevine, in Japan. *Journal of General Plant Pathology* 81:237-242.
- Landi, L., Murolo, S. & Romanazzi, G. 2019. Detection of '*Candidatus* Phytoplasma solani' in roots from Bois noir symptomatic and recovered grapevines. *Scientific Reports* 9(2013):1-12.

- Naue, C. R., Costa, V.S.O., Barbosa, M. A. G., Batista, D. C., Souza, E. B., & Mariano, R. L. R.** 2014. *Xanthomonas campestris* pv. *viticola* on grapevine cutting tools and water: survival and disinfection. *Journal of Plant Pathology* 96(3): 451-458.
- Soto-Giron, M. J., L. M. Rodriguez-R, C. Luo, M. Elk, H. Ryu, J. Hoelle, J. W. Santo Domingo, & K. T. Konstantinidis.** 2016. Biofilms on hospital shower hoses: characterization and implications for nosocomial infections. *Applied and Environmental Microbiology* 82(9):2872-2883.
- Voegel, T. M. & Nelson, L. M.** 2018. Quantification of *Agrobacterium vitis* from grapevine nursery stock and vineyard soil using droplet digital PCR. *Plant Disease* 102(11):2136-2141.
- Wilcox, W. F., Gubler, W. F. & Uyemoto, J. K.** 2015. *Compendium of Grape Diseases, Disorders, and Pests*. Second Edition. American Phytopathological Society, St. Paul, MN, USA. 232 pp.

Table 4: Nematode pests of grapevine

Last updated December 2021.

PEST	MODE of PARASITISM	REFERENCES	PRESENCE/ABSENCE		
			CAN	USA	MEX
<i>Gracilacus mirus</i> n. sp.	Migratory endo and ectoparasite	Raski, 1962	Ab1	Ab1	Ab1
<i>Helicotylenchus pseudorobustus</i> (Steiner) Golden	Migratory ecto and semi endoparasite	Dong <i>et al.</i> , 2007; Esser, 1982; Ravichandra, 2008; Siddiqui <i>et al.</i> , 1973; Subbotin <i>et al.</i> , 2015	P7	P2	Ab1
<i>Hemicriconemoides californianus</i> Pinochet and Raski	Ectoparasite	Esser, 1982; Pinochet and Raski, 1975	Ab1	P2	Ab1
<i>Hoplolaimus seinhorsti</i> Luc	Migratory endo and ectoparasite	Brown, <i>et al.</i> , 1993; Catalano <i>et al.</i> , 1992	Ab1	P2	Ab1
<i>Longidorus africanus</i> Merny	Root tip ectoparasite	Andres <i>et al.</i> , 1991; Raski, 1988, Wilcox <i>et al.</i> , 2015	Ab1	P2	Ab1
<i>Longidorus attenuatus</i>	Root tip ectoparasite	Arias and Andres, 1989; CABI 2021 edition; Griffiths and Robertson, 1984; Raski, 1988	Ab1	Ab1	Ab1
<i>Longidorus coespiticola</i> Hooper. [<i>Longidorus coespiticola</i> (maybe a variant spelling of the species name)]	Root tip ectoparasite	Arias and Andres, 1989; Ravichandra, 2008	Ab1	Ab1	Ab1
<i>Longidorus cretensis</i> n.sp	Root tip ectoparasite	Tzortzakakis <i>et al.</i> , 2001	Ab1	Ab1	Ab1
<i>Longidorus diadecturus</i> Eveleigh and Allen	Root tip ectoparasite	Robbins and Brown, 1991	P2	P2	Ab1
<i>Longidorus elongatus</i> (deMan) Thorne and Swanger	Root tip ectoparasite	Arias and Andres, 1989; CABI 2021 edition; Griffiths and Robertson, 1984; Raski, 1988	P7	P2	Ab1
<i>Longidorus euonymus</i> Mali and Hooper	Root tip ectoparasite	Barsi, 1994a; Choleva-Abadzhieva, 1975; Lone <i>et al.</i> , 2018	Ab1	Ab1	Ab1
<i>Longidorus fasciatus</i> Roca and Lamberti	Root tip ectoparasite	Brown <i>et al.</i> , 1993; Brown <i>et al.</i> , 1997	Ab1	Ab1	Ab1
<i>Longidorus juvenilis</i> Dalmasso	Root tip ectoparasite	Coiro <i>et al.</i> , 1992; Kleynhans <i>et al.</i> , 1966	Ab1	Ab1	Ab1
<i>Longidorus macrosoma</i> Hooper	Root tip ectoparasite	Andres <i>et al.</i> , 1991; Arias and Andres, 1989; Raski, 1988	Ab1	Ab1	Ab1
<i>Longidorus magnus</i> Lamberti,	Root tip ectoparasite	Lamberti <i>et al.</i> , 1982	Ab1	Ab1	Ab1

PEST	MODE of PARASITISM	REFERENCES	PRESENCE/ABSENCE		
			CAN	USA	MEX
Bleve-Zacheo and Arias					
<i>Meloidogyne arenaria</i> (Neal) Chitwood	Sedentary endoparasite	CABI 2021 edition; Cid del Prado <i>et al.</i> , 2001. Hugo and Storey, 2017; Powers <i>et al.</i> , 2005; Raski, 1988	P7	P2	P2
<i>Meloidogyne hapla</i> Chitwood	Sedentary endoparasite	CABI 2021 edition; Ogawa <i>et al.</i> , 1995; Powers <i>et al.</i> , 2005; Raski, 1988	P1	P2	P2
<i>Meloidogyne incognita</i> (Kofoid) Chitwood, syn. <i>Meloidogyne acrita</i> Chitwood	Sedentary endoparasite	Martínez, 1989; Ogawa <i>et al.</i> , 1995; Powers <i>et al.</i> , 2005; Raski, 1988	P7	P2	P2
<i>Meloidogyne javanica</i> (Treub) Chitwood	Sedentary endoparasite	Cid del Prado <i>et al.</i> , 2001; Ogawa <i>et al.</i> , 1995; Powers <i>et al.</i> , 2005; Raski, 1988	P7	P2	P2
<i>Meloidogyne mali</i> Itoh, Ohshima and Ichinoe	Sedentary endoparasite	Bridge and Starr, 2007; Itoh <i>et al.</i> , 1969	Ab1	P2	Ab1
<i>Meloidogyne nataliei</i> Golden, Rose and Bird	Sedentary endoparasite	Bird <i>et al.</i> , 1994; Raski, 1988	Ab1	P2	Ab1
<i>Merlinius brevidens</i> [Allen] Siddiqi, syn. <i>Geocenamus brevidens</i> (Allen) Siddiqi, (<i>Geocenamus brevidens</i> (possibly misspelled (Allen) Brzeski) (accepted name)	Root tip ectoparasite	Dong <i>et al.</i> , 2007; Esser 1982; McKenry and Roberts, 1985	P7	P2	Ab1
<i>Mesocriconema rusticum</i> (Micoletzky) Loof and De Grisse, syn. <i>Macroposthonia rusticum</i>	Ectoparasite	Siddiqui <i>et al.</i> , 1973	Ab1	P2	Ab1
<i>Mesocriconema xenoplax</i> (Raski) Loof and De Grisse, syn. <i>Criconemoides xenoplax</i> Raski, <i>Macroposthonia xenoplax</i> (Raski) De Grisse and Loof, <i>Criconemella xenoplax</i> (Raski) Luc and Raski	Ectoparasite	Dong <i>et al.</i> , 2007; Ferris <i>et al.</i> , 2004; Ogawa <i>et al.</i> , 1995; Raski, 1952	P2	P2	P2
<i>Paralongidorus maximus</i> (Butschli) Siddiqi	Root tip ectoparasite	Mc Elroy <i>et al.</i> , 1977	Ab1	Ab1	Ab1

PEST	MODE of PARASITISM	REFERENCES	PRESENCE/ABSENCE		
			CAN	USA	MEX
<i>Paratrichodorus pachydermus</i> (Seinhorst) Siddiqi	Ectoparasite	Kumari, 2010; Ravichandra, 2008	P7	P2	Ab1
<i>Pratylenchus brachyurus</i> (Godfrey) Filipjev and Schuurm. Stekh.	Migratory endo and ectoparasite	Cepeda and Hernández, 1991; Dong <i>et al.</i> , 2007; McKenry and Roberts, 1985; Oliveira <i>et al.</i> , 1999; Siddiqui <i>et al.</i> , 1973	Ab3	P2	P2
<i>Pratylenchus coffeae</i> Zimmerman Filipjev and Schuurmans Stekhoven	Migratory endo and ectoparasite	Hafez <i>et al.</i> , 1992; Hafez <i>et al.</i> , 2010; Silva and Inomoto, 2002	Ab1	P2	Ab3
<i>Pratylenchus crenatus</i> Loof	Migratory endo and ectoparasite	Brown <i>et al.</i> , 1980; Hafez <i>et al.</i> , 1992; Hafez <i>et al.</i> , 2010; Siddiqui <i>et al.</i> , 1973	P7	P2	Ab1
<i>Paratylenchus hamatus</i> Thorne and Allen	Migratory endo and ectoparasite	Dong <i>et al.</i> , 2007; Raski, 1952; Ravichandra, 2008; Siddiqui <i>et al.</i> , 1973; Van den Berg <i>et al.</i> , 2014	P7	P2	Ab1
<i>Pratylenchus hexincisus</i> Jenkins and Taylor	Migratory endo and ectoparasite	Carta <i>et al.</i> , 2001; Dong <i>et al.</i> , 2007	P7	P2	Ab1
<i>Pratylenchus neglectus</i> (Rensch) Filipjev and S. Stekhoven	Migratory endo and ectoparasite	Carta <i>et al.</i> , 2001; Hafez <i>et al.</i> , 1992; Hafez <i>et al.</i> , 2010; Siddiqui <i>et al.</i> , 1973; Subbotin <i>et al.</i> , 2008	P7	P2	Ab1
<i>Paratylenchus neoamblycephanus</i>	Migratory endo and ectoparasite	Dong <i>et al.</i> , 2007; McKenry and Roberts, 1985; Ravichandra, 2008	Ab1	P2	Ab1
<i>Pratylenchus penetrans</i> Cobb), Filipjev and Schwermans-Stekhoven	Migratory endo and ectoparasite	Carta <i>et al.</i> , 2001; Ogawa <i>et al.</i> , 1995; Potter <i>et al.</i> , 1984; Subbotin <i>et al.</i> , 2008; Villalobos <i>et al.</i> , 1980	P2	P2	Ab3
<i>Pratylenchus pratensis</i> (de Man) Filipjev	Migratory endo and ectoparasite	Handoo and Morgan, 1989. Hugo and Storey <i>et al.</i> , 2017	P7	P7	Ab3
<i>Pratylenchus vulnus</i> Allen and Jensen	Migratory endo and ectoparasite	Chitambar and Raski, 1984; Handoo and Morgan, 1989; Storey <i>et al.</i> , 2017; Ogawa <i>et al.</i> , 1995	Ab4	P2	Ab3
<i>Quinisulcius acutus</i> (Allen) Siddiqi (accepted name) syn. <i>Tylenchorhynchus acutus</i> Allen	Ectoparasite	Siddiqui <i>et al.</i> , 1973	P7	P2	Ab1
<i>Rotylenchulus reniformis</i> Linford and Oliveria	Semi endoparasite	Ravichandra, 2008	Ab1	P2	Ab1
<i>Rotylenchulus robustus</i> (de Man) Filip'ev syn.	Semi endoparasite	Cantalapiedra-Navarrete <i>et al.</i> , 2013; Dong <i>et al.</i> , 2007; Siddiqui <i>et al.</i> , 1973	P7	P2	Ab1

PEST	MODE of PARASITISM	REFERENCES	PRESENCE/ABSENCE		
			CAN	USA	MEX
<i>Rotylenchus fallorobustus</i> Sher					
<i>Tylenchorhynchus capitatus</i> Allen	Ectoparasite	Allen, 1955; Siddiqui <i>et al.</i> , 1973	P7	P2	Ab3
<i>Tylenchorhynchus clarus</i> Allen	Ectoparasite	Handoo <i>et al.</i> , 2014; McKenry and Roberts, 1985; Siddiqui <i>et al.</i> , 1973	Ab1	P2	Ab3
<i>Tylenchorhynchus claytoni</i> Steiner	Ectoparasite	Siddiqui <i>et al.</i> , 1973	P2	P2	Ab3
<i>Tylenchorhynchus cylindricus</i> Cobb	Ectoparasite	Siddiqui <i>et al.</i> , 1973	Ab1	P2	Ab3
<i>Tylenchorhynchus mashhoodi</i> Siddiqi and Basir	Ectoparasite	Dong <i>et al.</i> , 2007	Ab1	P2	Ab1
<i>Tylenchulus semipenetrans</i> Cobb	Semi endoparasite	Dong <i>et al.</i> , 2007; Edwards, M. 1988	Ab1	P2	P2
<i>Xiphinema americanum</i> Cobb	Root tip ectoparasite	Allen <i>et al.</i> , 1984; Ebsary <i>et al.</i> , 1984; Lone <i>et al.</i> , 2018; Ogawa <i>et al.</i> , 1995; Ramírez and Jiménez, 1987; Vrain and Rouselle, 1980; Weimin <i>et al.</i> , 2004	P2	P2	P2
<i>Xiphinema brevicolle</i> Bordello and DaCosta	Root tip ectoparasite	Bridge and Starr, 2007; Olivera <i>et al.</i> , 2004	Ab1	P2	Ab1
<i>Xiphinema bricolensis</i> Ebsary, Vrain and Graham	Root tip ectoparasite	Vrain, T. C. 1993	P2	P2	Ab1
<i>Xiphinema browni</i> Lazarova, Peneva and Kumari, <i>syn. Xiphinema mediterraneum</i> Martelli and Lamberti, <i>X. pachtaicum</i> (Tulaganov) Kirjanova	Root tip ectoparasite	Roca <i>et al.</i> , 1991; Esser. 1982	Ab1	P2	Ab1
<i>Xiphinema californicum</i> Lamberti and Bleve-Zacheo	Root tip ectoparasite	Georgi, 1988, Bridge and Starr, 2007	Ab1	P2	Ab1
<i>Xiphinema diversicaudatum</i> (Micoletzky) Thorne	Root tip ectoparasite	Hugo and Storey, 2017; Lone <i>et al.</i> , 2018; Weimin <i>et al.</i> , 2004	P7	P2	Ab1
<i>Xiphinema index</i> Thorne and Allen	Root tip ectoparasite	Lone <i>et al.</i> , 2018; Weimin <i>et al.</i> , 2004; Téliz and Goheen, 1968	Ab1	P2	Ab3
<i>Xiphinema insigne</i> Loos; <i>X. indicum</i> Siddiqi, <i>syn. X. neodimorphicaudatum</i> Khan,	Root tip ectoparasite	Lambert <i>et al.</i> , 1997; Lone <i>et al.</i> , 2018; Luc and Southey, 1980	Ab1	P2	Ab1

PEST	MODE of PARASITISM	REFERENCES	PRESENCE/ABSENCE		
			CAN	USA	MEX
<i>X. tugewai</i> Darekar and Khan					
<i>Xiphinema italiae</i> Meyl	Root tip ectoparasite	Weimin <i>et al.</i> , 2004	Ab1	Ab1	Ab1
<i>Xiphinema melitense</i> n. sp.	Root tip ectoparasite	Roca <i>et al.</i> , 1991	Ab1	Ab1	Ab1
<i>Xiphinema monohysterum</i> Brown	Root tip ectoparasite	McLeod and Khair, 1971	Ab1	Ab1	Ab1
<i>Xiphinema occiduum</i> n.sp.	Root tip ectoparasite	Ebsary <i>et al.</i> , 1984	P2	Ab1	Ab1
<i>Xiphinema pachtaicum</i> Tulaganov	Root tip ectoparasite	Roca <i>et al.</i> , 1991	Ab1	Ab1	Ab1
<i>Xiphinema pacificum</i> Ebsary, Vrain and Graham	Root tip ectoparasite	Vrain, T. C. 1993	P2	P2	Ab1
<i>Xiphinema rivesi</i> Dalmasso	Root tip ectoparasite	Akinbade <i>et al.</i> , 2014; Ebsary <i>et al.</i> , 1984; Georgi, 1988	P2	P2	Ab1
<i>Xiphinema simile</i> Lamberti, Choleva and Agostinelli	Root tip ectoparasite	Barsi, 1994	Ab1	Ab1	Ab1
<i>Xiphinema taylori</i> Lamberti, Ciancio, Agostinelli and Coiro	Root tip ectoparasite	Weimin <i>et al.</i> , 2004	Ab1	Ab1	Ab1
<i>Xiphinema vuittenezi</i> Luc, Lima, Weischer and Flegg	Root tip ectoparasite	Bridge and Starr, 2007; Weimin <i>et al.</i> , 2004	Ab1	P2	Ab1
<i>Zygotylenchus guevarai</i> Tobar Jiminez	Migratory endo and ectoparasite	Siddiqui, 1975	Ab1	Ab3	Ab1

References

- Akinbade, S. A., Mojtahedi, H., Guerra, L. Eastwell, K., Villamor, D. E. V., Handoo Z. A. & Skantar, A. M.** 2014. First Report of *Xiphinema rivesi* (Nematoda, Longidoridae) in Washington State. *Plant Disease*, 98: 1018.
- Allen, M. W.** 1955. A review of the nematode genus *Tylenchorhynchus*. University of California Publications in Zoology, 61, 129–166.
- Allen, W. R., Van Schagen, J. G., & Ebsary, B. A.** 1984. Comparative transmission of the peach rosette mosaic virus by Ontario populations of *Longidorus diadecturus* and *Xiphinema americanum* (Nematoda: Longidoridae). *Canadian Journal of Plant Pathology*, 6, 29–32.
- Arias, M. & Andres, M. F.** 1989. Virus-vector nematodes in cereals and fruit crops in Spain. EPPO (European and Mediterranean Plant Protection Organization) Bulletin, 19:625-632.
- Andres, M., Arias, M. & Bello, A.** 1991. Distribución ecológica del género *Longidorus* (Micoletzky) Filipjev en la región central de España. *Nematopica*, 21: 79-87.
- Barsi, L.** 1994. Bivulval females of *Longidorus euonymus*, *Xiphinema diversicaudatum* and *X. vuittenezi* (Nematoda: Dorylaimida). *Nematologia Mediterranea*, 22: 271-272.
- Bird, G., Diamond, C., Warner, F. & Davenport, J.** 1994. Distribution and Regulation of *Meloidogyne nataliei*. *Journal of Nematology*, 26: 727-730.
- Bridge, J. & Starr, J. L.** 2007. *Plant Nematodes of Agricultural Importance: A Color Handbook*. Academic Press Burlington, MA. 152 pp.
- Brown, D. J. F., Dalamasso, A. & Trudgill, D. L.** 1993. Nematode Pest of Soft Fruits and Vines. In K. Evans. *et al.* eds., *Plant parasitic nematodes in temperate agriculture*. Wallingford, UK, CABI, 427- 462 pp.
- Brown, D. J. F, Kyriakopoulou, P. E. & Robertson, W. M.** 1997. Frequency of transmission of Artichoke Italian latent nepovirus by *Longidorus fasciatus* (Nematoda: Longidoridae) from artichoke fields in the Iria and Kandia areas of Argolis in northeast Peloponnesus, Greece. *European Journal of Plant Pathology*, 103: 501-506.
- Brown, M. J., Riedel, R. M. & Rowe R. C.** 1980. Species of *Pratylenchus* Associated with *Solanum tuberosum* cv Superior in Ohio. *Journal of Nematology*, 12: 189-192.
- CABI.** 2021. Crop Protection Compendium. CAB International. <https://www.cabi.org/cpc/>.
- Cantalapiedra-Navarrete, C., Navas-Cortés, J. A., Liébanas, G., Vovlas, N., Subbotin, S. A., Palomares-Rius, J. E., & Castillo, P.** 2013. Comparative molecular and morphological characterisations in the nematode genus *Rotylenchus*: *Rotylenchus paravitis* n. sp., an example of cryptic speciation. *Zoologischer Anzeiger*, 252: 246–268.
- Carta, K. L, Skantar, M. A. & Handoo, A. Z.** 2001. Molecular, morphological, and thermal characters of 19 *Pratylenchus* spp. and relatives using the D3 segments of the nuclear LSU rDNA gene. *Nematopica*, 31: 193-208.
- Catalano, L., Savino, V. & Lamberti, F.** 1992. Presence of grapevine fanleaf nepovirus in population of longidorid nematodes and their vectoring capacity. *Nematologia Mediterranea*, 20: 67-70.
- Cepeda, S. M., & Hernández, B. J. R.** 1991. Control químico del nematodo de la lesión *Pratylenchus brachyurus* en el cultivo del manzano (*Pyrus malus* L.) en Arteaga, Coahuila. *Memorias XVIII Congreso Nacional de Fitopatología*. Resumen 214. Puebla, Pue., Mexico.
- Chitambar, J. J. & Raski, D. J.** 1984. Reactions of grape rootstocks to *Pratylenchus vulnus* and *Meloidogyne* spp. *Journal of Nematology*, 16: 166-170.
- Choleva-Abadzhieva, B.** 1975. Study of the species composition and spread of nematodes of the Family Longidoridae (Nematoda: Dorylaimoidea) on vine in Bulgaria. *Acta Zoologica Bulgaria*, 3:19-30.
- Cid del Prado, V.I., Tovar, S.A., & Hernández, J.A.** 2001. Distribución de especies y razas de *Meloidogyne* en México. *Revista Mexicana de Fitopatología*, 19: 32–39.

- Coiro, M.I., Agostinelli, A. & Lamberti, F.** 1992. Longidoridae (Nematoda) in the vineyards of the province of Verona. *Nematologia Mediterranea*, 20: 87-95.
- Dong, K., Chitambar, J., Subbotin, S., Alzubaidy, M., Luque-Williams, M., Romero, J., Kosta, K. & Luna, R.** 2007. Significant records in Nematology: California statewide nematode survey project for 2006. California Plant Pest and Damage Report July 2005 through December 2006. 23: 45-71.
- Ebsary, B. A., Potter, J. W., & Allen, W. R.** 1984. Redescription and distribution of *Xiphinema rivesi* Dalmasso, 1969 and *Xiphinema americanum* Cobb, 1913 in Canada with a description of *Xiphinema occiduum* n.sp. (Nematoda: Longidoridae). *Canadian Journal of Zoology*, 62: 1696–1702.
- Esser, R. P.** 1982. Nematodes associated with grapevines. Nematology Circular; Florida Department of Agriculture & Consumer Services, Division of Plant Industry No. 90:1-2.
- Edwards, M.** 1988. Effect of type of rootstock on yield of Carina grapevines (*Vitis vinifera*) and levels of citrus nematode (*Tylenchulus semipenetrans* Cobb). *Australian Journal of Experimental Agriculture*, 28: 283-286.
- Ferris, H., McKenry, M. V., Jaffee, B. A., Anderson, C. E. & Juurma, A.** 2004. Population characteristics and dosage trajectory analysis for *Mesocriconema xenoplax* in California Prunus orchards. *Journal of Nematology*. 36: 505-516.
- Georgi, L. L.** 1988. Morphological variation in *Xiphinema* spp. from New York orchards. *Journal of Nematology*, 20: 47-57.
- Griffiths, B. S. & Robertson, W. M.** 1984. Morphological and histochemical changes occurring during the lifespan of root-tip galls on *Lolium perenne* induced by *Longidorus elongatus*. *Journal of Nematology*, 16: 223-229.
- Hafez, S., Golden, A. M., Rashid, F. & Handoo, Z.** 1992. Plant parasitic nematodes associated with crops in Idaho and Eastern Oregon. *Nematropica*, 22: 193–204.
- Hafez, S. L., Sundararaj, P., Handoo, Z. A. & Siddiqi, M. R.** 2010. Occurrence and distribution of nematodes in Idaho crops. *International Journal of Nematology*, 20(1): 91-98.
- Handoo, A. Z. & Morgan A. G.** 1989. A key and diagnostic compendium to the species of the genus *Pratylenchus* Filipjev, 1936 (lesion nematodes). *Journal of Nematology*, 21: 202-218.
- Handoo, Z. A., Palomares-Rius, J.E., Cantalapiedra-Navarrete, C., Liebanas, G., Subbotin, S.A. & Castillo, P.** 2014. Integrative taxonomy of the stunt nematodes of the genera *Bitylenchus* and *Tylenchorhynchus* (Nematoda, Telotylenchidae) with description of two new species and a molecular phylogeny. *Zoological Journal of the Linnean Society*, 172: 231–264.
- Hugo, H. J. & Storey, S. G.** 2017. Nematode pests in deciduous fruit. In Fourie, H., Spaul, V.W., Jones, R.K., Daneel, M.S. & D. De Waele. eds. *Nematology in South Africa: A view from the 21st century*. Chapter 15. Springer International, Switzerland.
- Itoh, Y., Ohshima, Y. & Ichinohe, M.** 1969. A root-knot nematode, *Meloidogyne mali* n. sp. on apple-tree from Japan (Tylenchida: Heteroderidae). *Applied Entomology and Zoology*, 4:194-202.
- Kleynhans, K. P. N., Van den Berg, E., Swart, A., Marais, M. & Buckley, N. H.** 1996. Plant nematodes in South Africa. ARC-Plant Protection Research Institute. Pretoria, South Africa. 165 pp.
- Kumari, S.** 2010. Research Note: Description of *Paratrichodorus pachydermus* (Nematoda: Trichodoridae) from the Czech Republic. *Helminthologia*, 47(3): 196–198.
- Lambert, F., Lovev, T., Choleva, B., Brown, D. G. F., Agostinelli, A. & Radicci, V.** 1997. Morphometric variation and juvenile stages of some Longidorid nematodes from Bulgaria with comments on the number of juveniles stages of *Longidorus africanus*, *L. closelongatus* and *Xiphinema santos*. *Nematologia Mediterranea*, 26: 213-237.
- Lamberti, F., Bleve Zacheo, T., & Arias, M.** 1982 The Longidoridae of the Maltese Islands with

- the description of *Longidorus magnus* sp. n. and *Xiphinema melitense* sp. n. *Nematologia Mediterranea* 10: 183-200.
- Lone G. M., Zaki, F. A., Pathania S. S. & Mohi, S.** 2018. Population abundance and distribution of ectoparasitic plant nematodes associated with apple trees (*Malus domestica*) in apple belts of Kashmir, India. *International Journal of Current Microbiology and Applied Sciences*, 7(6): 3817–824.
- Luc, M. & Southey, J. F.** 1980. Study of biometrical variability in *Xiphinema insigne* Loos, 1949, and *X. elongatum* Schuurmans Stekhoven & Teunissen, 1938; description of *X. savanicola* n. sp. (Nematoda: Longidoridae) and comments on thelytokous species. *Revue de Nematologie*, 3(2):243–269.
- Martínez, G. M.** 1989. Observaciones sobre la distribución espacial en el suelo de *Xiphinema americanum* Cobb y *Meloidogyne incognita* Chitwood, en viñedos en la costa de Hermosillo. Tesis. Escuela de Agricultura y Ganadería, Universidad de Sonora.
- Mc Elroy, F. D., Brown, D. J. F. & Boag, B.** 1977. The Virus-vector and damage potential, morphometrics and distribution of *Paralongidorus maximus*. *Journal of Nematology*, 9: 122-130
- McKenry, M. V. & Roberts, P. A.** 1985. Phytonematology study guide. Publications, Division of Agriculture and Natural Resources, University of California.
- McLeod R. W. & Khair G. T.** 1971. *Xiphinema australiae* n. sp., its host range, observations on *X. radicum* Goodey, 1936 and *X. monohysterum* Brown, 1968 and a key to Monodelphic *Xiphinema* spp. (Nematoda: Longidoridae). *Nematologica*, 17: 58-68.
- Ogawa, J. M., Zehr, E. I., Bird, G. W., Ritchie, D. F., Uriu, K., & Uyemoto, J. K.** 1995. Compendium of stone fruit diseases. American Phytopathological Society, St. Paul, MN, USA. 61–64 pp.
- Oliveira, C. M. G., Inomoto, M. M., Vieira, A. M. C. & Monteiro, A. R.** 1999. Efeito de densidades populacionais de *Pratylenchus brachyurus* no crescimento de plantas de *Coffea arabica* cv. Mundo Novo e *C. canephora* cv. Apoata. *Nematropica*, 29: 215–221.
- Pinochet, J., & Raski, D. J.** 1975. Four new species of the genus *Hemicriconemoides* (Nematoda: Criconematidae). *Journal of Nematology*, 7: 263–270.
- Potter, J. W., Dirks, V. A., Johnson, P. W., Olthof, T. H. A., Layne, R. E. C., & McDonnell, M. M.** 1984. Response of peach seedlings to infection by the root lesion nematode *Pratylenchus penetrans* under controlled conditions. *Journal of Nematology* 16: 317-322.
- Powers, T. O., Mullin, P. G., Harris, T. S., Sutton, L. A. & Higgins, R. S.** 2005. Incorporating molecular identification of *Meloidogyne* spp. into a large-scale regional nematode survey. *Journal of Nematology*, 37: 226–235.
- Ramírez, A. J. A. & Jiménez, L. M.** 1987. Identificación y cuantificación de nematodos fitoparásitos asociados a la vid en la costa de Hermosillo, Sonora. Resumen 134. Memorias del XIV Congreso Nacional de Fitopatología. Morelia, Michoacán, Mexico.
- Raski, D. J.** 1952. On the morphology of *Criconemoides* Taylor, 1936, with descriptions of six new species (Nematoda: Criconematidae). *Proceedings of the Helminthological Society of Washington*, 19:85–99 pp.
- Raski, D. J.** 1962. Paratylenchidae n. fam. with descriptions of five new species of *Gracilacus* n. g. and an emendation of *Cacopaurus* Thorne, 1943, *Paratylenchus* Mikoletzky, 1922 and Criconematidae Thorne, 1943. *Proceedings of the Helminthological Society of Washington*, 29:189–207 pp.
- Raski, D. J.** 1988. Nematode parasites of grape. In Pearron. R.C & A.C. Austin eds. Compendium of grape diseases. American Phytopathological Society, St. Paul, MN, USA. 55–59 pp.
- Ravichandra, N. G.** 2008. Plant Nematology. I.K. International Publishing House Pvt. Ltd., New Delhi, India. 693 pp.
- Robbins, R. T., & Brown, D. J. F.** 1991. Comments on the Taxonomy, occurrence, and distribution of Longidoridae (Nematoda) in North America. *Nematologica* 37: 395-419.

- Roca, F., Lamberti, F. & Elia, F.** 1991. Longidoridae (Nematoda, Dorilaimida) Delle Regioni Italiane XII. Lumbria. *Nematologia Mediterranea*, 19: 279-289.
- Siddiqui, I. A., Sher, S. A., & French, A. M.** 1973. Distribution of plant parasitic nematodes in California. State of California Department of Food and Agriculture, Division of Plant Industry. 324 pp.
- Siddiqui, I. A.** 1975. *Zygotylenchus guevarai*. CIH description of plant parasitic nematodes. St. Albans, UK: Commonwealth Institute of Helminthology. Set 5, No. 65.
- Silva, R. A. & Inomoto, N. M.** 2002. Host-range characterization of two *Pratylenchus coffeae* isolates from Brazil. *Journal of Nematology*, 34:135–139.
- Storey, S. G., Malan, A. P., & Hugo, H. J.** 2017. Nematode pests of grapevine. Pages 325-343 in *Nematology in South Africa: A view from the 21st century*. Springer.
- Subbotin, S. A., Ragsdale, E. J., Mullens, T., Roberts, P. A., Mundo-Ocampo, M. & Baldwin, J. G.** 2008. A phylogenetic framework for root lesion nematodes of the genus *Pratylenchus* (Nematoda): Evidence from 18S and D2–D3 expansion segments of 28S ribosomal RNA genes and morphological characters. *Molecular Phylogenetics and Evolution*, 48(2): 491–505.
- Subbotin, S. A., Vovlas, N., Yeates, G. W., Hallmann, J., Kiewnick, S., Chizhov, V. N., Manzanilla-Lopez, R. H., Inserra, R. N., & Castillo, P.** 2015. Morphological and molecular characterization of *Helicotylenchus pseudorobustus* (Steiner, 1914) Golden, 1956 and related species (Tylenchida: Hoplolaimidae) with phylogeny of the genus. *Nematology*, 17: 27–52.
- Téliz, O. D. & Goheen, A. C.** 1968. Diseases of grapevines in Mexico. *Plant Disease Report*, 52: 372–373.
- Tzortzakakis E. A., Peneva V., Terzakis M., Neilson R. & Brown D. J. F.** 2001. *Longidorus cretensis* n. sp. (Nematoda: Longidoridae) from a vineyard infected with a foliar 'yellow mosaic' on Crete, Greece. *Systematic Parasitology*, 48: 131–139.
- Van den Berg, E., Tiedt, L. R. & Subbotin, S. A.** 2014. Morphological and molecular characterisation of several *Paratylenchus* Micoletzky, 1922 (Tylenchida: Paratylenchidae) species from South Africa and USA, together with some taxonomic notes. *Nematology*, 16(3): 323–358.
- Villalobos, B. O.** 1980. Efecto residual de Namacur 10G y Furadan 10G sobre las poblaciones de nematodos y el rendimiento en dos variedades de vid en la P. P. La Enramada, Municipio de Matamoros, Coahuila. Tesis de Licenciatura, Universidad Juárez del Estado de Durango, Gómez Palacio, México, 78 p.
- Vrain, T. C. & Rousselle, G. L.** 1980. Distribution of plant parasitic nematodes in Quebec apple orchards. *Plant Disease*, 64: 582–583.
- Vrain, T. C.** 1993. Restriction Fragment Length Polymorphism Separates Species of the *Xiphinema americanum* Group. *Journal of Nematology*, 25: 361-364.
- Weimin, Y., Szalanski, A. & Robbins, R. T.** 2004. Phylogenetic relationships and genetic variation in *Longidorus* and *Xiphinema*. (Nematoda: Longidoridae) using ITS1 sequence of nuclear ribosomal DNA. *Journal of Nematology* 36:220-231
- Wilcox, W. F., Gubler, W. F. & Uyemoto, J.K.** 2015. *Compendium of Grape Diseases, Disorders, and Pests (Second Edition)*. American Phytopathological Society, St. Paul, MN, USA. 232 pp.

Table 5: Arthropod pests of grapevines

Context: The pests named in this table are considered vectors, which promote the introduction (entry and establishment) and spread of the grapevine diseases listed in this standard.

VECTOR NAME	REFERENCES	PRESENCE / ABSENCE		
		CAN	USA	MEX
Acari				
Trombidiformes: Eriophyidae				
<i>Colomerus vitis</i> (Pagenstecher)	PMC, 2016; Peischmann & James, 2005; Roy <i>et al.</i> , 2009	P2	P2	Ab1
Insecta				
Hemiptera: Cicadellidae				
<i>Erythroneura ziczac</i> Walsh Vector status unconfirmed	Ávila-Rodríguez <i>et al.</i> , 2010; Maw <i>et al.</i> , 2000; Metcalf, 1968; Pinto & Viggiani, 1987	P1	P2	Ab3
<i>Homalodisca vitripennis</i> (Germar)	Martin <i>et al.</i> , 2014; Maw <i>et al.</i> , 2000; Saguez <i>et al.</i> , 2014	Ab1	P1	P2
<i>Oncopsis alni</i> Schrank	Maw <i>et al.</i> , 2000; Saguez <i>et al.</i> , 2014	Ab1	Ab1	Ab1
<i>Scaphoideus titanus</i> Ball	Maw <i>et al.</i> , 2000; Metcalf, 1967; Saguez <i>et al.</i> , 2014	P2	P1	Ab1
Hemiptera: Cixiidae				
<i>Hyalesthes obsoletus</i> Signoret	Maw <i>et al.</i> , 2000	Ab1	Ab1	Ab1
<i>Oliarus atkinsonii</i> (Myers)	Maw <i>et al.</i> , 2000	Ab1	Ab1	Ab1
Hemiptera: Coccidae				
<i>Parthenolecanium corni</i> (Bouché)	Cockerell, 1899; Emond & Cerezke, 1990; García Morales <i>et al.</i> , 2016; King 1901; Maw <i>et al.</i> , 2000	P1	P1	Ab3
<i>Pulvinaria innumerabilis</i> (Rathvon)	Maw <i>et al.</i> , 2000	P2	P1	Ab1

<i>Pulvinaria vitis</i> (L.)	Maw <i>et al.</i> , 2000; Maw, 2019	P2	P2	Ab1
Hemiptera: Membracidae				
<i>Spissistilus festinus</i> (Say)	Beyer <i>et al.</i> , 2017; Caldwell 1949; Cockerell, 1899; Deitz & Wallace, 2012; Maw <i>et al.</i> , 2000; Metcalf & Wade, 1965; Wildermuth, 1915	P2	P1	Ab3
Hemiptera: Pseudococcidae				
<i>Heliococcus adenostomae</i> McKenzie	García Morales <i>et al.</i> , 2016	Ab1	P2	Ab1
<i>Phenacoccus aceris</i> (Signoret)	Maw <i>et al.</i> , 2000; García Morales <i>et al.</i> , 2016	P2	P2	Ab1
<i>Planococcus citri</i> (Risso)	García Morales <i>et al.</i> , 2016	Ab1	P1	P2
<i>Planococcus ficus</i> (Signoret)	García Morales <i>et al.</i> , 2016	Ab1	P2	P3
<i>Pseudococcus calceolariae</i> (Maskell)	Ben-Dov, 1994; García Morales <i>et al.</i> , 2016	Ab1	P2	Ab3
<i>Pseudococcus comstocki</i> (Kuwana)	Ben-Dov, 1994; García Morales <i>et al.</i> , 2016; Maw <i>et al.</i> , 2000; Willians & Granara, 1992	P2	P1	Ab3
<i>Pseudococcus longispinus</i> (Targioni Tozzetti)	García Morales <i>et al.</i> , 2016; Paiero & Marshall, 2003; Salazar-Torres & Solís-Aguilar, 1990; Urias-López & Carrillo-Sánchez, 1983	P2	P1	Ab3
<i>Pseudococcus maritimus</i> (Ehrhorn)	Maw <i>et al.</i> , 2000	P2	P2	P2
<i>Pseudococcus viburni</i> (Signoret)	Arriola Padilla, 2009; García Morales <i>et al.</i> , 2016; Maw <i>et al.</i> , 2000, as <i>P. affinis</i> . In 2003 <i>P. viburni</i> was found in the Biodome in Quebec (Maw, 2019); McKenzie 1967, as <i>P. obscurus</i> ; Miller, 1996	P2	P1	Ab3

References

- Arriola Padilla, V. J.** 2009. La familia Pseudococcidae (Hemiptera: Coccoidea) en México. Ph.D. Thesis. Colegio de Postgraduados. México. 150 pp.
- Ávila-Rodríguez, V., González-Hernández, A., Alvarado-Gómez, O. G., Nava-Camberos, U., & Cortéz-Mondaca E.** 2010. Géneros de Trichogrammatidae en México asociados a cultivos agrícolas y áreas naturales aledañas. *Southwestern Entomologist*, 35(2):177–191.
- Ben-Dov, Y.** 1994. A systematic catalogue of the mealybugs of the world (Insecta: Homoptera: Coccoidea: Pseudococcidae and Putoidae) with data on geographical distribution, host plants, biology, and economic importance. Intercept Limited, Andover, UK. 686 pp.
- Beyer, B. A., Srinivasan, R., Roberts, P. M. & Abney, M. R.** 2017. Biology and management of the three-cornered alfalfa hopper (Hemiptera: Membracidae) in alfalfa, soybean, and peanut. *Journal of Integrated Pest Management* 8(1):doi: 10.1093/jipm/pmx003.
- Caldwell, J. S.** 1949. A generic revision of the treehoppers of the tribe Ceresini in America north of Mexico based on a study of the male genitalia. *Proceedings of the United States National Museum* 98(3234):491-521 +6 plates.
- Cockerell, T. D. A.** 1899. Insecta. Rhynchota, Hemiptera-Homoptera. [Aleurodidae and Coccidae]. Vol. II. Part. 2. *Biologia Centrali-Americana*.
- Deitz, L. L. & Wallace, M. S.** 2012. Richness of the Nearctic treehopper fauna (Hemiptera: Aetalionidae and Membracidae). *Zootaxa* 3423:1-26.
- Emond, F. I. & Cerezke, H. F.** 1990 Forest insect and disease conditions in Alberta, Saskatchewan, Manitoba, and the Northwest Territories in 1989 and predictions for 1990. Information report NOR-X-313. Forestry Canada, Northwest Region, Northern Forestry Centre, Edmonton, Alberta.
- García Morales, M., Denno, B. D., Miller, D. R., Miller, G. L., Ben-Dov, Y. & Hardy, N. B.** 2016. ScaleNet: A literature-based model of scale insect biology and systematics. Database. doi: 10.1093/database/bav118. <http://scalenet.info>. [Online].
- King, G. B.** 1901. The Coccidae of British North America. *The Canadian Entomologist* 33(7):193-200.
- Martin, K.W., A.C. Hodges, & N. C. Leppla.** 2012. Citrus pests, glassy-winged sharpshooter. Pdf here: <https://idtools.org/id/citrus/pests/factsheet.php?name=Glassy-winged+sharpshooter>
- Maw, E.** 2019. Personal Communication.
- Maw, E., Foottit, R. G., Hamilton, K. G. A. & Scudder, G. G. E.** 2000. Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa. 220 pp.
- McKenzie, H.L.** 1967. Mealybugs of California with taxonomy, biology, and control of North American species (Homoptera: Coccoidea: Pseudococcidae). University of California Press, Berkeley, California.
- Metcalf, Z.** 1967. General Catalogue of Homoptera. Fasc6. Cicadelloidea. Part 10. Euscelidae. USDA, ARS. 2695 pp. PDF here : <https://www.biodiversitylibrary.org/item/107309#page/1/mode/1up>
- Metcalf, Z.,** 1968. General catalogue of Homoptera. Fasc6. Cicadelloidea. Part 17. Cicadellidae. USDA, ARS. 1513 pp. Pdf available here <https://www.biodiversitylibrary.org/item/107146#page/1195/mode/1up>
- Metcalf, Z. P. & V. Wade.** 1965. General Catalogue of the Homoptera a Supplement to Fascicle I- Membracidae of the General Catalogue of Hemiptera Section II. North Carolina State University. 1552 pp. PDF available here: <https://www.biodiversitylibrary.org/item/194618#page/1/mode/1up>.
- Miller, D. R.** 1996. Checklist of the scale insects (Coccoidea: Homoptera) of Mexico.

- Proceedings of the Entomological Society of Washington, 98(1):68-86.
- Paiero, S. M., Marshall, S. A. & Hamilton, K.G.A.** 2003. New records of Hemiptera from Canada and Ontario. *Journal of the Entomological Society of Ontario* 134:115-129.
- Prischmann, D. A. and James, D. G.** 2005. New mite records (Acari: Eryophyidae, Tetranychidae) from grapevines in Oregon and Washington State. *International Journal of Acarology* 31(3): 289-291.
- Pinto, J. D. & Viggiani, G.** 1987. Two new Trichogrammatidae (Hymenoptera) from North America: *Ittysella lagunera* Pinto & Viggiani (n. gen., n. sp.) and *Epoligosita mexicana* Viggiani (n. sp.). *The Pan-Pacific entomologist*. 63(4):371-376.
- PMC.**, 2016. Crop profile for grape in Canada. Pesticide Risk Reduction Program, Agriculture and Agri-Food Canada, Ottawa.
- Roy, M., Fortin, C. & Mainguy, J.** 2009. Ravageurs galligènes de la vigne au Québec. Laboratoire de diagnostic en phytoprotection, MAPAQ, Sainte-Foy, Quebec.
- Salazar-Torres, J. C. & Solis-Aguilar, J. F.** 1990. Escamas (Homoptera: Coccoidea) presentes en cuatro especies frutales de la familia Rosaceae en Zacatlán, Puebla. *Revista Chapingo*, 15(67-68):135-137.
- Saguez, J., Olivier, C., Hamilton, A., Lowery, T., Stobbs, L., Lasnier, J., Galka, B., Chen, X., Mauffette, Y. & Vincent, C.** 2014. Diversity and abundance of leafhoppers in Canadian vineyards. *Journal of insect science (Online)* 14:73-73.
- Urias-López, M. A. & Carrillo-Sánchez, J. L.** 1983. Plagas principales de la yuca, *Manihot esculenta* Crantz, en la sabana de Huimanguillo, Tabasco. *Agricultura Técnica en México*, 9(1):65-83.
- Wildermuth, V. L.** 1915. Three-cornered alfalfa hopper. *Journal of Agricultural Research* 3: 343-364.
- Williams, D. J. & Granara, W., M. C.** 1992. Mealybugs of central and South America. Wallingford. CAB International. United Kingdom. 635 pp.